

CLASS-COMBINATORIAL MODEL CATEGORIES

B. CHORNY AND J. ROSICKÝ*

ABSTRACT. We develop an extension of the framework of combinatorial model categories. The category of small presheaves over large indexing categories and ind-categories are the basic examples of non-combinatorial model categories embraced by the new machinery called class-combinatorial model categories.

The definition of the new class of model categories is based on the corresponding extension of the theory of locally presentable and accessible categories developed in the companion paper [11], where we introduced the concepts of locally class-presentable and class-accessible categories.

In this work we prove that the category of weak equivalences of a nice class-combinatorial model category is class-accessible. Our extension of J. Smith's localization theorem depends on the verification of a cosolution-set condition. The deepest result is that the (left Bousfield) localization of a class-combinatorial model category with respect to a strongly class-accessible localization functor is class-combinatorial again.

1. INTRODUCTION

The theory of combinatorial model categories pioneered by J. Smith in the end of '90s has become a standard framework for abstract homotopy theory. The foundations of the subject may be found in [2] and [14, 15]; a concise exposition has appeared in [25, A.2.6].

A model category \mathcal{M} is combinatorial if it satisfies two conditions. The first condition requires that the underlying category \mathcal{M} is locally presentable (see, e.g., [1] for the definition and an introduction to the subject). The second condition demands that the model structure be cofibrantly generated (see, e.g., [20] for the definition and discussion).

Several interesting examples of non-combinatorial model categories without Quillen equivalent combinatorial models appeared over the past decade. For example the categories of pro-spaces and ind-spaces

Key words and phrases. class-combinatorial model category, Bousfield localization.

* Supported by MSM 0021622409 and GAČR 201/11/0528. The hospitality of the Australian National University is gratefully acknowledged.

were applied in new contexts in homotopy theory [12, 21] resulting in non-cofibrantly generated model structures constructed on non-locally presentable categories. The maturation of the calculus of homotopy functors [17] stimulated the development of the abstract homotopy theory of small functors over large categories [10] resulting in formulation of the basic ideas of Goodwillie calculus in the language of model categories [3]. The model categories used for this purpose are also not cofibrantly generated and the underlying category of small functors from spaces to spaces (or spectra) is not a locally presentable category.

However, all the model categories from the examples above are *class-cofibrantly* generated (except for the pro-categories, which are *class-fibrantly* generated). This extension of the classical definition was introduced in [8], which in turn developed the ideas by E. Dror Farjoun originated in the equivariant homotopy theory [16].

The purpose of the current paper is to develop a framework extending J. Smith's combinatorial model categories, so that the model categories of small presheaves on large categories and ind-categories of model categories (the opposite categories of pro-categories) would become the examples of the newly defined *class-combinatorial* model categories. The definition of a class-combinatorial model category consists, similarly to that of a combinatorial model category, of two conditions: the underlying category is required to be *class-locally presentable* and the model structure must be class-cofibrantly generated. As we mentioned above, the second condition was studied in the earlier work [8], while the first condition relies on a concept of class-locally presentable category. This was introduced and studied in the companion project [11], which is a prerequisite for reading this paper. Informally, a *class-accessible* category is a category closed under sufficiently large filtered colimits and equipped with a class of generating objects with uniformly bounded presentability ranks, such that every object in the category is a sufficiently large filtered colimit of generating objects. A class-accessible category is called class-locally presentable if in addition we require existence of all limits and colimits. See [11, Def. 2.1] for the formal introduction and discussion of these concepts.

The main results of our paper generalize the corresponding results about the combinatorial model categories. First, in Theorem 2.11 we prove that the levelwise weak equivalences in the category of small presheaves form a class-accessible category (see [11] for the definition). In Remark 2.13 we formulate mild conditions which guarantee that a class-combinatorial model category has a class-accessible subcategory of weak equivalences. Such class-combinatorial model categories are called nice in this paper. Then we prove a class-version of Smith's

theorem on existence of combinatorial model structures [2, 1.7] in Theorem 2.14.

The central result of J. Smith's theory is the localization theorem, stating the existence of the (left Bousfield) localization of any combinatorial model category with respect to any *set* of maps. After a brief discussion of construction of localization functors with respect to cone-coreflective *classes* of cofibrations with bounded presentability ranks of domains and codomains, we prove in Theorem 3.10 a variant of a localization theorem for nice class-combinatorial model categories with respect to strongly class-accessible homotopy localization functors (i.e., localization functors preserving λ -filtered colimits and λ -presentable objects for some cardinal λ). Although an application of our localization theorem depends on the verification of a cosolution-set condition for the class of intended generating trivial cofibrations, we are able to check this condition in many interesting situations. In the last Theorem 3.14 we prove that in the cases where the localization with respect to a strongly class-accessible functor exists, the localized model category is class-combinatorial again.

We conclude the paper by several examples of localized model categories. Using Theorem 3.14 we show that the n -polynomial model category constructed in [3] is class-combinatorial (Example 3.16). On the other hand, there is a model category constructed in [9] as a localization of a class-combinatorial model category with respect to an inaccessible localization functor that happens to be non-class-cofibrantly generated (Example 3.17).

2. CLASS-COMBINATORIAL MODEL CATEGORIES

Recall that a weak factorization system was introduced in [11, Def. 4.7]. We repeat this definition here, since it is of central importance for this section.

Definition 2.1. A weak factorization system $(\mathcal{L}, \mathcal{R})$ in a locally class- λ -presentable category \mathcal{K} was called *cofibrantly class- λ -generated* if $\mathcal{L} = \text{cof}(\mathcal{C})$ for a cone-coreflective class \mathcal{C} of morphisms such that

- (1) morphisms from \mathcal{C} have λ -presentable domains and codomains and
- (2) any morphism between λ -presentable objects has a weak factorization with the middle object λ -presentable.

To be *cone-coreflective* means for each f there is a subset \mathcal{C}_f of \mathcal{C} such that each morphism $g \rightarrow f$ in \mathcal{K}^\rightarrow with $g \in \mathcal{C}$ factorizes as

$$g \rightarrow h \rightarrow f$$

with $h \in \mathcal{C}_f$.

If the weak factorization is functorial, a cofibrantly class- λ -generated weak factorization system is cofibrantly class- μ -generated for each regular cardinal $\mu \geq \lambda$. Without functoriality, condition (2) does not need to go up to μ and thus we will make it a part of the following definition.

Definition 2.2. Let \mathcal{K} be a class locally- λ -presentable model category. We say that \mathcal{K} is *class- λ -combinatorial* if both (cofibrations, trivial fibrations) and (trivial cofibrations, fibrations) are cofibrantly class- μ -generated weak factorization systems for every regular cardinal $\mu \geq \lambda$.

It is called *class-combinatorial* if it is class- λ -combinatorial for some regular cardinal λ .

Any combinatorial model category is class-combinatorial. The reason is that weak factorizations are functorial and, moreover, the resulting functors are accessible. Thus they are strongly accessible and this property goes up for $\lambda \triangleleft \mu$ (cf. [27]).

Example 2.3. Let \mathbf{SSet} denote the category of simplicial sets. Given a simplicial category \mathcal{A} , we denote by $\mathcal{P}(\mathcal{A})$ the category of *small simplicial presheaves* on \mathcal{A} . The objects are functors $\mathcal{A}^{\text{op}} \rightarrow \mathbf{SSet}$ which are small weighted colimits of simplicial representable functors. Although in [11] we used this notation for small set-valued presheaves, in this paper we reserve the notation $\mathcal{P}(\mathcal{A})$ only for small simplicial presheaves in order to avoid confusion. If we will happen to use the category of ordinary presheaves on \mathcal{A} , then we will use the full notation: $\mathbf{SSet}^{\mathcal{A}^{\text{op}}}$.

The simplicial category $\mathcal{P}(\mathcal{A})$ is complete provided that \mathcal{A} is complete (see [13]); completeness is meant in the enriched sense (see, e.g., [4] or [23]).

The category $\mathcal{P}(\mathcal{A})$ is always class-finitely-accessible, because each small simplicial presheaf is a conical colimit of presheaves from $\mathcal{G} = \{\text{hom}_{\mathcal{A}}(-, A) \otimes K \mid A \in \mathcal{A}, K \in \mathbf{SSet}\}$. Therefore each small presheaf is a filtered colimit of finite colimits of elements of \mathcal{G} . The elements of \mathcal{G} are, in turn, filtered colimits of the elements of $\mathcal{G}_{\text{fin}} = \{\text{hom}(-, A) \otimes L \mid A \in \mathcal{A}, L \in \mathbf{SSet}_{\text{fin}}\}$ where $\mathbf{SSet}_{\text{fin}}$ denotes the full subcategory of \mathbf{SSet} consisting of finitely presentable simplicial sets. Therefore, every small presheaf is a filtered colimit of finite colimits of the elements of \mathcal{G}_{fin} .

We are going to show that, for a complete simplicial category \mathcal{A} , the category $\mathcal{P}(\mathcal{A})$ equipped with the projective model structure is class-combinatorial. We will need the following result.

Lemma 2.4. *Let \mathcal{K} be a class locally λ -presentable simplicial category. Then μ -presentable objects in \mathcal{K} are closed under finite weighted limits for every uncountable regular cardinal $\mu \geq \lambda$.*

Proof. A weight $W: \mathcal{D} \rightarrow \mathbf{SSet}$ is finite if \mathcal{D} has finitely many objects and all its hom-objects $\mathcal{D}(c, d)$ and all values $W(d)$ are finitely presentable simplicial sets (see [24, 4.1]) Following [24, 4.3], finite weighted limits can be constructed from finite conical limits and cotensors with finitely presentable simplicial sets. Thus we have to show that μ -presentable objects in \mathcal{K} are closed under these limits.

Finite conical limits in \mathcal{K} coincide with finite limits in the underlying category \mathcal{K}_0 since their definition has nothing to do with the enrichment. We intend to use a similar statement proved earlier in presheaf categories. For this purpose we apply the limit preserving embedding constructed in [11, 2.6] $E: \mathcal{K} \rightarrow \mathbf{Set}^{\mathcal{A}^{\text{op}}}$, where $\mathcal{A} = \text{pres}_{\lambda} \mathcal{K}$.

Let $F: \mathcal{D} \rightarrow \mathcal{K}$ be a finite diagram of μ -presentable objects. If we denote $P = \lim_{D \in \mathcal{D}} F(D)$, then $EP = \lim_{D \in \mathcal{D}} EF(D)$, where $EF(D)$ are representable functors, since $F(D) \in \mathcal{A}$, hence EF is a diagram of μ -presentable objects, since E is *strongly* λ -accessible, hence strongly μ -accessible by [11, 2.8], because $\mu \geq \lambda$.

Consider the category $\mu\text{-colex}(\mathcal{A})$, which is a free completion of \mathcal{A} under μ -small colimits, as a full subcategory of μ -presentable objects of the free completion of \mathcal{A} under colimits $\mathbf{Set}^{\mathcal{A}^{\text{op}}}$ (by a simple observation that every μ -small colimit of representables is μ -small and a more technical converse statement proved in [26, pp. 35-37]. Note that every limit existing in $\mu\text{-colex}(\mathcal{A})$ remains a limit in $\mathbf{Set}^{\mathcal{A}^{\text{op}}}$, in other words, the full embedding constructed above preserves limits. We argue as follows. If L is an inverse limit of a diagram in $\mu\text{-colex}(\mathcal{A})$ and L' is a possible candidate for an inverse limit of the same diagram in $\mathbf{Set}^{\mathcal{A}^{\text{op}}}$, then L' may be represented as a μ -filtered colimit of μ -presentable presheaves, i.e., elements of $\mu\text{-colex}(\mathcal{A})$: $L' = \text{colim}_i P_i$. The natural map $L \rightarrow L'$ has an inverse, since everyone of P_i 's admits a natural map into L , because it forms a cone over the original diagram through the map $P_i \rightarrow L'$, and together these maps define a natural map $L' \rightarrow L$. We conclude that $L \cong L'$ by naturality considerations.

The category $\mu\text{-colex}(\mathcal{A})$ is closed under finite limits by [22, 4.9], hence μ -presentable objects in $\mathbf{Set}^{\mathcal{A}^{\text{op}}}$ are closed under finite limits. In other words we have shown that EP is μ -presentable, but we need to show that P is μ -presentable. Let \mathcal{K} be a μ -filtered colimit $K =$

$\operatorname{colim}_i K_i$, then

$$\begin{aligned} \operatorname{hom}(P, K) &= \operatorname{hom}(EP, EK) = \operatorname{hom}(EP, \operatorname{colim}_i E(K_i)) = \\ &= \operatorname{colim}_i \operatorname{hom}(EP, EK_i) = \operatorname{colim}_i \operatorname{hom}(P, K_i) \end{aligned}$$

It remains to show that the μ -presentable objects are closed under cotensors with finitely presentable simplicial sets. The latter are finite conical colimits of Δ_n , $n = 0, 1, 2, \dots$. Thus we have to show that μ -presentable objects in $\mathcal{P}(\mathcal{A})$ are closed under cotensors with Δ_n 's.

Let H be μ -presentable in $\mathcal{P}(\mathcal{A})$. Since H is a μ -small colimit of tensors H_i of finitely presentable simplicial sets with representables and both colimits and cotensors in $\mathcal{P}(\mathcal{A})$ are pointwise, we have

$$\begin{aligned} H^{\Delta_n}(A) &= (\operatorname{colim} H_i)^{\Delta_n}(A) = (\operatorname{colim} H_i(A))^{\Delta_n} \\ &= \operatorname{hom}(\Delta_n, \operatorname{colim} H_i(A)) \cong \operatorname{colim} \operatorname{hom}(\Delta_n, H_i(A)) \\ &= \operatorname{colim} H_i(A)^{\Delta_n} = (\operatorname{colim} H_i^{\Delta_n})(A) \end{aligned}$$

for each A in \mathcal{A} . Hence

$$H^{\Delta_n} \cong \operatorname{colim} H_i^{\Delta_n}$$

and thus it suffices to show that each $H_i^{\Delta_n}$ is μ -presentable. Since each H_i is equal to $V \otimes \operatorname{hom}(-, B)$ for some finitely presentable simplicial set V and B in \mathcal{A} , we get for the same reasons as above

$$\begin{aligned} \operatorname{hom}(\Delta_n, V \otimes \operatorname{hom}(A, B)) &= \operatorname{hom}(\Delta_n, V \times \operatorname{hom}(A, B)) \cong \\ &\cong \operatorname{hom}(\Delta_n, V) \times \operatorname{hom}(\Delta_n, \operatorname{hom}(A, B)) \cong V^{\Delta_n} \times \operatorname{hom}(A, B^{\Delta_n}) = \\ &= (V^{\Delta_n} \otimes \operatorname{hom}(-, B^{\Delta_n}))(A) \end{aligned}$$

and thus

$$(V \otimes \operatorname{hom}(-, B))^{\Delta_n} \cong V^{\Delta_n} \otimes \operatorname{hom}(-, B^{\Delta_n}).$$

The latter objects are μ -presentable. □

Proposition 2.5. *Let \mathcal{A} be a complete simplicial category. Then $\mathcal{P}(\mathcal{A})$ is class- λ -combinatorial with respect to the projective model structure for each uncountable regular cardinal λ .*

Proof. Following [10], $\mathcal{P}(\mathcal{A})$ is a model category where the generating classes \mathcal{I} and \mathcal{J} of cofibrations and trivial cofibrations are coreflective and satisfy condition 2.1(1) for any regular cardinal λ . In fact, \mathcal{I} consists of morphisms

$$\partial \Delta_n \otimes \operatorname{hom}(-, A) \rightarrow \Delta_n \otimes \operatorname{hom}(-, A)$$

and \mathcal{J} of morphisms

$$\Lambda_n^k \otimes \text{hom}(-, A) \rightarrow \Delta_n \otimes \text{hom}(-, A),$$

and all involved domains and codomains are finitely presentable. We have to show that they satisfy 2.1(2) as well, i.e., that they are bounded. Let λ be uncountable and $f : G \rightarrow H$ be a morphism between λ -presentable objects and consider a morphism $g \rightarrow f$ where $g \in \mathcal{I}$. Following the proof of 3.7 in [10], this morphism corresponds to a morphism $\text{hom}(-, A) \rightarrow P$ where P is the pullback

$$G^{\partial \Delta_n} \times_{H^{\partial \Delta_n}} H^{\Delta_n}.$$

Since P is λ -presentable (see 2.4), there is a choice of a set \mathcal{T}_f from [11] 4.8 (2) whose cardinality does not exceed λ . Since all morphisms from \mathcal{I} have finitely presentable domains and codomains, the factorization of f stops at ω . Thus the cardinality of \mathcal{T}_f^* is smaller than λ . Following [11] 4.8 (2), condition [11] 3.7 (2) is satisfied. The argument for \mathcal{J} is the same. \square

Remark 2.6. A very useful property of a combinatorial model categories is that the class of weak equivalences is an accessible and accessibly embedded subcategory of the category of morphisms \mathcal{K}^\rightarrow (see [28, 4.1] or [25, A2.6.6]). Together with Smith's theorem [2] this implies the localization theorem for combinatorial model categories with respect to sets of maps. It would be natural to expect that a similar property holds in the class-combinatorial situation. Unfortunately we were unable to prove it in this generality. But in many interesting situations we are able to prove that the class of weak equivalences is a class-accessible subcategory of the category of morphisms.

Lemma 2.7. *Let \mathcal{A} be a complete simplicial category. Then $\mathcal{P}(\mathcal{A})$ admits a strongly class-accessible fibrant replacement functor.*

Proof. The functor $\text{Ex}^\infty : \mathbf{SSet} \rightarrow \mathbf{SSet}$ is a finitely accessible fibrant replacement simplicial functor (see [18]). For a small simplicial functor $F : \mathcal{A}^{\text{op}} \rightarrow \mathbf{SSet}$, let $\text{Fib}(F)$ be the composition

$$\mathcal{A}^{\text{op}} \xrightarrow{F} \mathbf{SSet} \xrightarrow{\text{Ex}^\infty} \mathbf{SSet}.$$

We will show that this composition is small.

The category of finitely accessible simplicial functors $\mathbf{SSet} \rightarrow \mathbf{SSet}$ is equivalent to the category of simplicial functors $\mathbf{SSet}^{\mathbf{SSet}_{\text{fin}}}$. This equivalence sends a finitely accessible functor $\mathbf{SSet} \rightarrow \mathbf{SSet}$ to its restriction to $\mathbf{SSet}_{\text{fin}}$. Thus hom-functors $\text{hom}(S, -) : \mathbf{SSet} \rightarrow \mathbf{SSet}$ with S finitely presentable correspond to hom-functors $\text{hom}(S, -) : \mathbf{SSet}_{\text{fin}} \rightarrow \mathbf{SSet}$. Since every simplicial functor $\mathbf{SSet}_{\text{fin}} \rightarrow \mathbf{SSet}$ is a

weighted colimit of hom-functors, every finitely accessible simplicial functor $\mathbf{S}\mathbf{Set} \rightarrow \mathbf{S}\mathbf{Set}$ is a weighted colimit of hom-functors $\mathrm{hom}(S, -)$ with S finitely presentable. Thus the composition $\mathrm{Ex}^\infty F$ is a weighted colimit of functors $\mathrm{hom}(S, -)F$ with S finitely presentable. But the functor $\mathrm{hom}(S, -)F$ is small because it is isomorphic to the cotensor F^S . The reason is that natural transformations

$$\mathcal{A}(-, A) \rightarrow \mathrm{hom}(S, -)F = \mathrm{hom}(S, F-)$$

correspond to morphisms $S \rightarrow FA$, i.e., to morphisms

$$S \rightarrow \mathcal{P}(\mathcal{A})(\mathcal{A}(-, A), F)$$

which, by the definition of the cotensor, correspond to morphisms $\mathcal{A}(-, A) \rightarrow F^S$. Consequently, $\mathrm{Ex}^\infty F$ is small as a weighted colimit of small functors.

We have obtained the functor $\mathrm{Fib} : \mathcal{P}(\mathcal{A}) \rightarrow \mathcal{P}(\mathcal{A})$ which clearly has fibrant values. Moreover, the pointwise trivial cofibration

$$\mathrm{Id}_{\mathbf{S}\mathbf{Set}} \rightarrow \mathrm{Ex}^\infty$$

yields a weak equivalence $\mathrm{Id}_{\mathcal{P}(\mathcal{A})} \rightarrow \mathrm{Fib}$. Thus Fib is a fibrant replacement functor on $\mathcal{P}(\mathcal{A})$. Since Ex^∞ is finitely accessible, so is Fib . We know that Ex^∞ is a weighted colimit of hom-functors $\mathrm{hom}(S, -)$ with S finitely presentable. The corresponding weight is λ -small for an uncountable regular cardinal λ . Let F be λ -presentable in $\mathcal{P}(\mathcal{A})$. Then $\mathrm{Fib}(F)$ is a λ -small weighted colimit of $\mathrm{hom}(S, -)F \cong F^S$ and the latter functors are λ -presentable following 2.4. Hence $\mathrm{Fib}(F)$ is λ -presentable (the argument is analogous to [24], 4.14). Thus Fib is strongly class- λ -accessible. \square

Definition 2.8. Let \mathcal{A} be a complete simplicial category and $f : A \rightarrow B$ be a morphism in $\mathcal{P}(\mathcal{A})$. The *mapping cocylinder* on f is the object $S(f)$ of $\mathcal{P}(\mathcal{A})$ defined as a pullback

$$\begin{array}{ccc} S(f) & \xrightarrow{r} & B^{\Delta_1} \\ q \downarrow & & \downarrow B^j \\ B & \xrightarrow{f} & B \end{array}$$

where $j : \Delta_0 \rightarrow \Delta_1$ sends 0 to 0.

Remark 2.9. Mapping cocylinders were used in the PhD thesis of J.P. Serre in order to replace an arbitrary map of topological spaces by a fibration. We are going to use them basically for the same purpose in $\mathcal{P}(\mathcal{A})$. The advantage over the modern methods of factorization is the functoriality of $S(-)$.

Lemma 2.10. *Let \mathcal{A} be a complete simplicial category and $f : A \rightarrow B$ a morphism of fibrant objects in $\mathcal{P}(\mathcal{A})$. Then there exists a factorization*

$$A \xrightarrow{i} S(f) \xrightarrow{p} B$$

of f where i is a weak equivalence and p is a fibration.

Proof. The pullback in 2.8 may be split in two pullbacks

$$\begin{array}{ccc} S(f) & \xrightarrow{r} & B^{\Delta_1} \\ q_1 \downarrow & & \downarrow B^u \\ A \times B & \xrightarrow{f \times \text{id}_B} & B \times B \\ q_2 \downarrow & & \downarrow B^v \\ A & \xrightarrow{f} & B \end{array}$$

where

$$\Delta_0 \xrightarrow{v} \Delta_0 + \Delta_0 \xrightarrow{u} \Delta_1$$

is the factorization of j . Since both u and v are cofibrations and B is fibrant, the vertical morphisms B^u and B^v are fibrations (see [19] 9.3.9 (2a)). Moreover, since j is a trivial cofibration, B^j is a trivial fibration. Thus q_1 and q_2 are fibrations and $q = q_2 q_1$ is a trivial fibration.

Let t denote the unique morphism $\Delta_1 \rightarrow \Delta_0$. Since,

$$B^j f^{\Delta_1} A^t = f A^j A^t = f,$$

there is a unique morphism $i : A \rightarrow S(f)$ such that $qi = \text{id}_A$ and $ri = f^{\Delta_1} A^t$. Since q is a trivial fibration, i is a weak equivalence. Since $B^v : B \times B \rightarrow B$ is the first projection of the product, $q_2 : A \times B \rightarrow A$ is the first projection as well. Let $p_2 : A \times B \rightarrow B$, $\bar{p}_2 : B \times B \rightarrow B$ be the second projections and $v' : \Delta_0 \rightarrow \Delta_0 + \Delta_0$ be the second injection of the coproduct. Then $\bar{p}_2 = B^{v'}$ and

$$\begin{aligned} p_2 q_1 i &= \bar{p}_2 (f \times \text{id}_B) q_1 i = \bar{p}_2 B^u r i = \bar{p}_2 B^u f^{\Delta_1} A^t = B^{v'} B^u f^{\Delta_1} A^t \\ &= B^{uv'} f^{\Delta_1} A^t = f A^{uv'} A^t = f. \end{aligned}$$

Since B is fibrant, p_2 is a fibration and thus $p = p_2 q_1$ is a fibration. We have $f = pi$. \square

Theorem 2.11. *Let \mathcal{A} be a complete simplicial category and denote by \mathcal{W} the class of weak equivalences in the projective model structure on $\mathcal{P}(\mathcal{A})$. Then \mathcal{W} is a class-accessible category strongly accessibly embedded in $\mathcal{P}(\mathcal{A})^\rightarrow$.*

Proof. Let $\text{Fib} : \mathcal{P}(\mathcal{A}) \rightarrow \mathcal{P}(\mathcal{A})$ be the strongly class-accessible fibrant replacement functor constructed in 2.7. Consider the functor

$$R : \mathcal{P}(\mathcal{A})^{\rightarrow} \rightarrow \mathcal{P}(\mathcal{A})^{\rightarrow}$$

assigning to a morphism $f : A \rightarrow B$ the fibration $p : S(\text{Fib}(f)) \rightarrow \text{Fib}(B)$ from 2.10. Since the construction of $S(f)$ uses only finite limits, the functor $S(-) : \mathcal{P}(\mathcal{A})^{\rightarrow} \rightarrow \mathcal{P}(\mathcal{A})$ is strongly class-accessible by 2.4. Therefore the functor $R(-)$ is also strongly class-accessible. A morphism $\alpha : F \rightarrow G$ in $\mathcal{P}(\mathcal{A})$ is a weak equivalence if and only if $\text{Fib}(f)$ is a weak equivalence, hence if and only if $R(f)$ is a trivial fibration.

Let \mathcal{F}_0 denote the full subcategory of $\mathcal{P}(\mathcal{A})^{\rightarrow}$ consisting of trivial fibrations. Following 2.5 and [11] 4.9, \mathcal{F}_0 is class- λ -accessible and strongly class- λ -accessibly embedded in $\mathcal{P}(\mathcal{A})^{\rightarrow}$ for every uncountable regular cardinal λ . Since \mathcal{W} is given by the pullback

$$\begin{array}{ccc} \mathcal{K}^{\rightarrow} & \xrightarrow{R} & \mathcal{K}^{\rightarrow} \\ \uparrow & & \uparrow \\ \mathcal{W} & \longrightarrow & \mathcal{F}_0 \end{array}$$

whose vertical leg on the right is transportable, \mathcal{W} is equivalent to the pseudopullback (see [11] 3.2). Thus [11] 3.1 implies that \mathcal{W} is a class-accessible subcategory of $\mathcal{P}(\mathcal{A})^{\rightarrow}$. \square

Definition 2.12. A class-combinatorial model category \mathcal{K} is *nice* if the class of weak equivalences \mathcal{W} is a class-accessible, strongly class-accessibly embedded subcategory of $\mathcal{K}^{\rightarrow}$.

Remark 2.13. We have just proved that $\mathcal{P}(\mathcal{A})$ equipped with the projective model structure is a nice model category for any complete simplicial category \mathcal{A} . The same argument applies to every simplicial class-combinatorial model category which is equipped with a strongly class-accessible fibrant replacement functor and whose μ -presentable objects are closed under finite weighted limits for each $\mu \geq \lambda$ (where λ is a cardinal). We are not aware of any example of a class-combinatorial model category, which fails to be nice.

We now prove a version of Smith's theorem [2, 1.7] in the class-combinatorial case.

Theorem 2.14. *Let \mathcal{K} be a class-locally λ -presentable category, \mathcal{I} a λ -bounded class of morphisms and \mathcal{W} a class of morphisms of \mathcal{K} such that*

- (1) \mathcal{W} is a class- λ -accessible and strongly class- λ -accessibly embedded subcategory of \mathcal{K}^\rightarrow with the 2-out-of-3 property,
- (2) $\mathcal{I}^\square \subseteq \mathcal{W}$, and
- (3) $\text{cof}(\mathcal{I}) \cap \mathcal{W}$ is closed under pushout and transfinite composition and cone-coreflective in \mathcal{K}^\rightarrow .

Then, taking $\text{cof}(\mathcal{I})$ for cofibrations and \mathcal{W} for weak equivalences, we get a model category structure on \mathcal{K} .

Proof. Since \mathcal{I} is λ -bounded, $(\text{cof}(\mathcal{I}), \mathcal{I}^\square)$ is a cofibrantly class- λ -generated weak factorization system. For every λ -presentable $w \in \mathcal{W}$, we construct a factorization in \mathcal{K} into a cofibration j followed by a trivial fibration. By 2-out-of-3 property for \mathcal{W} and (2), j is in \mathcal{W} . Let \mathcal{J} be the class of these morphisms j for all λ -presentable $w \in \mathcal{W}$.

We will check now the conditions of Lemma [2, 1.8]. We have to show that for every morphism $i \rightarrow w$ in \mathcal{K}^\rightarrow with $i \in \mathcal{I}$ and $w \in \mathcal{W}$ there exists $j \in \mathcal{J}$ that factors it $i \rightarrow j \rightarrow w$. First note that there exists a λ -presentable $w' \in \mathcal{W}$, which factors the original morphism, since every w is a λ -filtered colimit of λ -presentable objects \mathcal{W} and every $i \in \mathcal{I}$ is λ -presentable in \mathcal{K}^\rightarrow ; we used here that the inclusion of \mathcal{W} to \mathcal{K}^\rightarrow preserves λ -presentable objects. Next, decompose that morphism w' into a cofibration $j \in \mathcal{J}$ followed by a trivial fibration. The lifting axiom in \mathcal{K} finishes this argument.

Lemma [2, 1.8] implies that $\text{cof } \mathcal{J} = \text{cof } \mathcal{I} \cap \mathcal{W}$. The requirement that $\text{cof } \mathcal{I} \cap \mathcal{W}$ is cone-coreflective in \mathcal{K}^\rightarrow ensures that \mathcal{J} is cone-coreflective as well (by the same argument as above). By construction, the domains of all the elements in \mathcal{J} are λ -presentable. Hence \mathcal{J} satisfies the assumptions of [11] 4.3 and thus $(\text{cof}(\mathcal{J}), \mathcal{J}^\square)$ is a weak factorization system. Since \mathcal{W} is closed under retracts in \mathcal{K}^\rightarrow (cf. [1] 2.4 and 2.5), we get a model category structure on \mathcal{K} . \square

Remark 2.15. Let \mathcal{K} be a locally presentable category, \mathcal{I} a set of morphisms and \mathcal{W} a class of morphisms of \mathcal{K} such that

- (1) \mathcal{W} has the 2-out-of-3 property and is closed under retracts in \mathcal{K}^\rightarrow ,
- (2) $\mathcal{I}^\square \subseteq \mathcal{W}$, and
- (3) $\text{cof}(\mathcal{I}) \cap \mathcal{W}$ is closed under pushout and transfinite composition.

Then, taking $\text{cof}(\mathcal{I})$ for cofibrations and \mathcal{W} for weak equivalences, we get a combinatorial model category if and only if the inclusion of \mathcal{W} in \mathcal{K}^\rightarrow is accessible. This is the content of Smith's theorem (see [2] for sufficiency and [25] or [28] for necessity).

We do not know whether this can be generalized to class-accessible setting and 2.14 is what we are able to do. The question is whether

cone-coreflectivity of $\text{cof}(\mathcal{I}) \cap \mathcal{W}$ follows from the other assumptions. We also do not know whether the model category in 2.14 is class-combinatorial. Indeed, we only know that the class \mathcal{J} satisfies 2.1(1).

Homotopy equivalences can be defined in any category \mathcal{K} with finite coproducts which is equipped with a weak factorization system $(\mathcal{L}, \mathcal{R})$ (see [28]). Recall that a *cylinder object* $C(K)$ of an object K is given by an $(\mathcal{L}, \mathcal{R})$ factorization of the codiagonal

$$\nabla : K + K \xrightarrow{\gamma_K} C(K) \xrightarrow{\sigma_K} K$$

We denote by

$$\gamma_{1K}, \gamma_{2K} : K \rightarrow C(K)$$

the compositions of γ_K with the coproduct injections.

Then, as usual, we say that morphisms $f, g : K \rightarrow L$ are *homotopic*, and write $f \sim g$, if there is a morphism $h : C(K) \rightarrow L$ such that the following diagram commutes

$$\begin{array}{ccc} K + K & \xrightarrow{(f,g)} & L \\ & \searrow \gamma_K \quad \nearrow h & \\ & C(K) & \end{array}$$

Here, (f, g) is induced by f and g . The homotopy relation \sim is clearly reflexive, symmetric, compatible with the composition and does not depend on the choice of a cylinder object. But, it is not transitive in general and we will denote its transitive hull by \approx . We get the quotient functor

$$Q : \mathcal{K} \rightarrow \mathcal{K}/\approx.$$

A morphism $f : K \rightarrow L$ is called a *homotopy equivalence* if Qf is an isomorphism, i.e., if there exists $g : L \rightarrow K$ such that both $fg \approx \text{id}_L$ and $gf \approx \text{id}_K$. The full subcategory of \mathcal{K}^\rightarrow consisting of homotopy equivalences with respect to a weak factorization system $(\mathcal{L}, \mathcal{R})$ will be denoted by $\mathcal{H}_{\mathcal{L}}$. The following result generalizes [28], 3.8.

Proposition 2.16. *Let \mathcal{K} be a locally class-presentable category and $(\mathcal{L}, \mathcal{R})$ be a weak factorization system with a strongly class-accessible cylinder functor. Then $\mathcal{H}_{\mathcal{L}}$ is a full image of a strongly class-accessible functor into \mathcal{K}^\rightarrow .*

Proof. Given $n < \omega$, let \mathcal{M}_n be the category whose objects are $(4n+2)$ -tuples

$$\tau = (f, g, a_1, \dots, a_n, b_1, \dots, b_n, h_1, \dots, h_n, k_1, \dots, k_n)$$

of morphisms $f : A \rightarrow B$, $g : B \rightarrow A$, $a_1, \dots, a_n : A \rightarrow A$, $b_1, \dots, b_n : B \rightarrow B$, $h_1, \dots, h_n : C(A) \rightarrow A$ and $k_1, \dots, k_n : C(B) \rightarrow B$. Morphisms are pairs (u, v) of morphisms $u : A \rightarrow A'$ and $v : B \rightarrow B'$ such that $f'u = vf$, $g'v = ug$, $uh_i = h'_i C(u)$ and $vk_i = k'_i C(v)$ for $i = 1, \dots, n$. This category is obtained by an inserter construction inserting our $n + 2$ morphisms among Id and C . Since C is strongly class-accessible, the procedure of the proof of [11] 3.9 yields that \mathcal{M}_n is a class-accessible category.

Let $\overline{\mathcal{M}}_n$ be the full subcategory of \mathcal{M}_n such that $h_1\gamma_A = (gf, a_1)$, $h_i\gamma_A = (a_i, a_{i+1})$, $h_n\gamma_n = (a_n, \text{id}_A)$, $k_1\gamma_A = (fg, b_1)$, $k_i\gamma_A = (b_i, b_{i+1})$ and $k_n\gamma_n = (b_n, \text{id}_B)$ where $1 < i < n$. This category is obtained from \mathcal{M}_n by an equifier construction and, by the same reason as above, the procedure of the proof of [11] 3.7 yields that $\overline{\mathcal{M}}_n$ is class-accessible. Moreover, its inclusion into \mathcal{M}_n is strongly accessible.

We have full embeddings

$$M_{m,n} : \mathcal{M}_m \rightarrow \mathcal{M}_n,$$

for $m < n$, which take the missing a_i, b_i, h_i, k_i as the identities. The union \mathcal{M} of all \mathcal{M}_n 's is a class-accessible category. Since all $\overline{\mathcal{M}}_n$'s are strongly accessibly embedded to \mathcal{M}_n , $\overline{\mathcal{M}}$ is strongly accessibly embedded by to $\overline{\mathcal{M}}$. Let

$$F : \overline{\mathcal{M}} \rightarrow \mathcal{K}^{\rightarrow}$$

sends each $(4n + 2)$ -tuple above to f . This is a strongly class-accessible functor whose image is $\mathcal{H}_{\mathcal{L}}$. \square

3. LEFT BOUSFIELD LOCALIZATIONS

Recall that \tilde{h} is a *cofibrant approximation* of h if there is a commutative square

$$\begin{array}{ccc} \tilde{A} & \xrightarrow{v} & A \\ \tilde{h} \downarrow & & \downarrow h \\ \tilde{B} & \xrightarrow{w} & B \end{array}$$

where v and w are weak equivalences and \tilde{A}, \tilde{B} are cofibrant.

Definition 3.1. Let \mathcal{K} be a class-combinatorial simplicial model category and \mathcal{F} a class of morphisms of \mathcal{K} . Assume that \mathcal{F} contains only cofibrations between cofibrant objects. An object K in \mathcal{K} is called *\mathcal{F} -local* if it is fibrant and

$$\text{hom}(f, K) : \text{hom}(B, K) \rightarrow \text{hom}(A, K)$$

is a weak equivalence of simplicial sets for each $f: A \rightarrow B$ in \mathcal{F} .

A morphism h of \mathcal{K} is called an \mathcal{F} -local equivalence if $\text{hom}(\tilde{h}, K)$ is a weak equivalence for each \mathcal{F} -local object K ; here, \tilde{h} is a cofibrant approximation of h .

The full subcategory of \mathcal{K} consisting of \mathcal{F} -local objects is denoted $\text{Loc}(\mathcal{F})$ and the full subcategory of \mathcal{K}^\rightarrow consisting of \mathcal{F} -local equivalences is denoted $\text{LEq}(\mathcal{F})$.

We say that there exists a *left Bousfield localization* of \mathcal{F} if cofibrations in \mathcal{K} and \mathcal{F} -local equivalences form a model category structure on \mathcal{K} .

Remark 3.2. (1) It is easy to see that the definition of a local \mathcal{F} -equivalence does not depend on the choice of a cofibrant approximation.

(2) Following [19], 9.3.3 (2), any weak equivalence in \mathcal{K} is an \mathcal{F} -local equivalence. On the other hand, every \mathcal{F} -local equivalence between \mathcal{F} -local objects is a weak equivalence in \mathcal{K} (cf. [18] X.2.1. 2)).

(3) If \mathcal{K} is left proper then the intersection of cofibrations and \mathcal{F} -local equivalences is closed under pushout and transfinite composition (see [19], 13.3.10, 17.9.4 for a non-trivial part of the proof); the trivial part is that $\text{hom}(-, K)$ sends colimits to limits and cofibrations to fibrations. It is also closed under retracts in \mathcal{K}^\rightarrow of course.

Given a morphism f , $\{f\}$ -local objects are called f -local and analogously for f -local equivalences. The corresponding categories are called $\text{Loc}(f)$ and $\text{LEq}(f)$.

Proposition 3.3. *Let \mathcal{K} be a class-combinatorial simplicial model category and \mathcal{F} a set of cofibrations between cofibrant objects of \mathcal{K} . Then $\text{Loc}(\mathcal{F})$ is a class-accessible category strongly accessibly embedded in \mathcal{K} .*

Proof.

$$\begin{array}{ccc} \mathcal{K} & \xrightarrow{\text{hom}(f, -)} & \mathbf{SSet}^\rightarrow \\ \uparrow & & \uparrow \\ \text{Loc}(f) & \longrightarrow & \mathcal{W} \end{array}$$

is a pullback where \mathcal{W} denotes weak equivalences in \mathbf{SSet} . Since the vertical leg on the right is transportable, $\text{Loc}(f)$ is a pseudopullback and thus it is class-accessible and its inclusion to \mathcal{K} is strongly class-accessible (see [11] 3.1, 3.2 and 2.11). Since

$$\text{Loc}(\mathcal{F}) = \bigcap_{f \in \mathcal{F}} \text{Loc}(f),$$

the result follows from [11] 3.3. \square

Definition 3.4. Let \mathcal{K} be a simplicial model category and $f : A \rightarrow B$ a cofibration of cofibrant objects. Consider a pushout

$$\begin{array}{ccc} \partial\Delta_n \otimes A & \xrightarrow{\text{id} \otimes f} & \partial\Delta_n \otimes B \\ \downarrow i_n \otimes \text{id} & & \downarrow p_{n1} \\ \Delta_n \otimes A & \xrightarrow{p_{n2}} & P_{f,n} \end{array}$$

where $i_n : \partial\Delta_n \rightarrow \Delta_n$ is the inclusion of the boundary into a simplex. Let $h_{f,n} : P_{f,n} \rightarrow \Delta_n \otimes B$ be the canonical morphism, which is a cofibration since \mathcal{K} is simplicial.

Cofibrations $h_{f,n}$, $n = 0, 1, \dots$ are called *f-horns*. If \mathcal{F} is a class of cofibrations, then we denote by $\text{Hor}(\mathcal{F})$ the collection of all *f-horns*, for all $f \in \mathcal{F}$.

Remark 3.5. Every $h_{f,n} \in \text{Hor}(\mathcal{F})$ is an \mathcal{F} -local equivalence because the morphism

$$\text{hom}(h_{f,n}, K) : \text{hom}(\Delta_n \otimes B, K) \rightarrow \text{hom}(P_{f,n}, K)$$

is a weak equivalence for every \mathcal{F} -local object K . In fact, the morphism

$$\text{hom}(\text{id} \otimes f, K) : \text{hom}(\Delta_n \otimes B, K) \rightarrow \text{hom}(\Delta_n \otimes A, K)$$

is a weak equivalence because K is \mathcal{F} -local and $\text{hom}(p_{n,2}, K)$ is a trivial fibration as a pullback of the trivial fibration

$$\text{hom}(\text{id} \otimes f, K) : \text{hom}(\partial\Delta_n \otimes B, K) \rightarrow \text{hom}(\partial\Delta_n \otimes A, K).$$

Thus it suffices to use the 2-out-of-3 property.

We used the fact that *f-horns* are cofibrations between cofibrant objects and that the definition of an \mathcal{F} -local equivalence does not depend on the choice of a cofibrant approximation.

Lemma 3.6. *Let \mathcal{K} be a simplicial model category and \mathcal{F} a class of cofibrations between cofibrant objects of \mathcal{K} . Then a fibrant object K of \mathcal{K} is \mathcal{F} -local if and only if it is injective to all *f-horns* for $f \in \mathcal{F}$.*

Proof. Since each $f \in \mathcal{F}$ is a cofibration, $\text{hom}(f, K)$ is a fibration for each fibrant object K . Thus a fibrant object K is \mathcal{F} -local if and only if $\text{hom}(f, K)$ is a trivial fibration for each $f \in \mathcal{F}$. This is the same as having the right lifting property with respect to each inclusion $i_n : \partial\Delta_n \rightarrow \Delta_n$. The latter is clearly equivalent to being injective with respect to *f-horns* $h_{f,n}$ for all $f \in \mathcal{F}$. \square

Lemma 3.7. *Let \mathcal{F} be a cone-coreflective class of cofibrations between λ -presentable cofibrant objects. Then $\text{Hor}(\mathcal{F})$ is a cone-coreflective class of morphisms between λ -presentable objects.*

Proof. Since $\partial\Delta_n \otimes B$ and $P_{f,n}$ are λ -presentable provided that A and B are λ -presentable, we have to prove that $\text{Hor}(\mathcal{F})$ is cone-coreflective. Let $f: A \rightarrow B$ be an element of \mathcal{F} . Given a commutative square

$$\begin{array}{ccc} P_{f,n} & \xrightarrow{\quad} & X \\ h_{f,n} \downarrow & & \downarrow g \\ \Delta_n \otimes B & \xrightarrow{\quad} & Y \end{array}$$

with $h_{f,n} \in \text{Hor}(\mathcal{F})$ and g arbitrary, we form, by adjunction, the following commutative square:

$$\begin{array}{ccc} A & \xrightarrow{\quad} & X^{\Delta^n} \\ f \downarrow & & \downarrow g' \\ B & \xrightarrow{\quad} & Q_{g,n} \end{array}$$

where $Q_{g,n} = X^{\partial\Delta^n} \times_{Y^{\partial\Delta^n}} Y^{\Delta^n}$.

Since \mathcal{F} is cone-coreflective, there exists a set of morphisms $\mathcal{F}_{g'} = \{f': A' \rightarrow B'\} \subset \mathcal{F}$ such that any morphism $f \rightarrow g'$ in \mathcal{K}^\rightarrow factors through some element $f' \in \mathcal{F}_{g'}$. Unrolling back the adjunction, we obtain the set of horns $\text{Hor}(\mathcal{F}_{g'}) = \{h_{f',n}: P_{f',n} \rightarrow \Delta^n \otimes B' \mid n \geq 0\}$ which depends entirely on g . Thus $\text{Hor}(\mathcal{F})$ is cone-coreflective. \square

Remark 3.8. (1) Let \mathcal{F} be a cone-coreflective class of cofibrations between λ -presentable cofibrant objects in a class λ -combinatorial simplicial model category \mathcal{K} . Then $\text{Loc}(\mathcal{F})$ is weakly reflective and closed under λ -filtered colimits in \mathcal{K} (following 3.7, 3.6 and [11] 4.4). Recall that a weak reflection $r_K: K \rightarrow K^*$ is obtained as a factorization

$$K \xrightarrow{r_K} K^* \longrightarrow 1.$$

in $(\text{cof}(\text{Hor}(\mathcal{F}) \cup \mathcal{C}), (\text{Hor}(\mathcal{F}) \cup \mathcal{C})^\square)$ where \mathcal{C} is a bounded class such that $\text{cof}(\mathcal{C})$ are cofibrations in \mathcal{K} . Thus r_K belongs to $\text{cof}(\text{Hor}(\mathcal{F}) \cup \mathcal{C})$. If \mathcal{K} is left proper then, following 3.2 (3), $\text{cof}(\text{Hor}(\mathcal{F}) \cup \mathcal{C}) \subseteq \text{cof}(\mathcal{C}) \cap \text{LEq}(\mathcal{F})$. Hence r_K is both a cofibration and an \mathcal{F} -local equivalence.

In order to apply the class version of Smith's theorem 2.14 we need to know that the local equivalences are class-accessible and strongly accessibly embedded. The only known technique to establish it is to

represent the local equivalences as a class of maps taken into a class-accessible strongly accessibly embedded subcategory (usually a category of weak equivalences in a nice class-combinatorial model category) and then apply [11, 3.1].

Unfortunately we do not always know that the weak reflections constructed above are functorial. But in the case that there exists a functor $L : \mathcal{K} \rightarrow \text{Loc}(\mathcal{F})$ and a natural transformation $\eta : \text{Id} \rightarrow L$ such that η_K is an \mathcal{F} -local equivalence for each K in \mathcal{K} , such a functor L is called an \mathcal{F} -localization functor.

(2) Given a model category \mathcal{K} and a functor $L : \mathcal{K} \rightarrow \mathcal{K}$, then $\text{LEq}(L)$ will denote the class of morphisms sent by L to weak equivalences. If an \mathcal{F} -localization functor L exists, then $\text{LEq}(\mathcal{F}) = \text{LEq}(L)$.

In fact h is an \mathcal{F} -local equivalence iff its cofibrant approximation \tilde{h} is an \mathcal{F} -local equivalence. Since η_K is an \mathcal{F} -local equivalence for each K , \tilde{h} is an \mathcal{F} -local equivalence iff $L(\tilde{h})$ is an \mathcal{F} -local equivalence, i.e., a weak equivalence in \mathcal{K} (see 3.2 (2)).

Proposition 3.9. *Let \mathcal{K} be a nice class-combinatorial model category and $L : \mathcal{K} \rightarrow \mathcal{K}$ be a strongly class-accessible functor. Then $\text{LEq}(L)$ is a class-accessible category strongly class-accessibly embedded in \mathcal{K}^\rightarrow .*

Proof. By assumption, the class \mathcal{W} of weak equivalences is class-accessible and strongly class-accessibly embedded in \mathcal{K}^\rightarrow . Since $\text{LEq}(L)$ is given by the pullback

$$\begin{array}{ccc} \text{LEq}(L) & \longrightarrow & \mathcal{W} \\ \downarrow & & \downarrow \\ \mathcal{K}^\rightarrow & \xrightarrow{L^\rightarrow} & \mathcal{K}^\rightarrow \end{array}$$

having the vertical leg on the right transportable, $\text{LEq}(L)$ is a pseudopullback and thus class-accessible and strongly class-accessibly embedded in \mathcal{K}^\rightarrow (see [11] 3.1 and 3.2). \square

Theorem 3.10. *Let \mathcal{K} be a nice, class-combinatorial, left proper, simplicial model category and let \mathcal{F} be a class of morphisms in \mathcal{K} . Suppose there exists a strongly class-accessible \mathcal{F} -localization functor $L : \mathcal{K} \rightarrow \mathcal{K}$. Then the left Bousfield localization of \mathcal{K} with respect to \mathcal{F} exists if and only if the intersection of $\text{LEq}(\mathcal{F})$ with the cofibrations of \mathcal{K} is a cone-coreflective class of morphisms.*

Proof. Necessity immediately follows from the existence of the (trivial cofibration, fibration) factorizations in the localized model category cf. [11] 4.2 (2)).

In order to establish sufficiency, we will verify the conditions of 2.14. By 3.8 (2) and 3.9 the subcategory $\text{LEq}(\mathcal{F})$ is class-accessible. There is a regular cardinal λ such that $\text{LEq}(\mathcal{F})$ is class- λ -accessible and \mathcal{K} is class- λ -combinatorial. In fact, $\text{LEq}(\mathcal{F})$ is class- μ -accessible and \mathcal{K} is class- ν -combinatorial, it suffices to take $\mu, \nu \triangleleft \lambda$. Let \mathcal{I} be the generating class of cofibrations in \mathcal{K} . Then $\mathcal{I}^\square \subseteq \text{LEq}(\mathcal{F})$ because $\text{LEq}(\mathcal{F})$ contains all weak equivalences. Following 3.2 (3), $\text{cof } \mathcal{I} \cap \text{LEq}(\mathcal{F})$ is closed under pushouts and transfinite compositions. \square

Example 3.11. Let \mathcal{A} be a complete simplicial category. Then $\mathcal{P}(\mathcal{A})$ equipped with the projective model structure is a class-combinatorial model category (see 2.5). Let $f : V \rightarrow W$ be a cofibration of simplicial sets. Then the class $\mathcal{F} = \{f \otimes \text{hom}(-, A) \mid A \in \mathcal{A}\}$ is bounded. The argument is the same as in the proof of 2.5. The localization of $\mathcal{P}(\mathcal{A})$ with respect to \mathcal{F} is equivalent to the levelwise localization with respect to f .

Let $L_f : \mathbf{SSet} \rightarrow \mathbf{SSet}$ be the f -localization functor, i.e., a fibrant replacement functor in the f -localized model category structure on \mathbf{SSet} . Then L_f is finitely accessible provided that V and W are finitely presentable. Moreover, L_f can always be chosen to be simplicial (see [19, 1.7] or [29, 24.2]). Similarly to Lemma 2.7, we get a strongly class-accessible simplicial functor $L : \mathcal{P}(\mathcal{A}) \rightarrow \mathcal{P}(\mathcal{A})$ assigning to F the composition $L_f F$. Since $\text{LEq}(L) = \text{LEq}(\mathcal{F})$, $\text{LEq}(\mathcal{F})$ is a class-accessible subcategory of $\mathcal{P}(\mathcal{A})^\rightarrow$ by 3.9.

In a general case, L_f is accessible and we would need an extension of 2.4 to λ -small weighted limits. This is valid but we have not burdened our paper with a proof.

Example 3.12. Let \mathcal{F} be a set of cofibrations between cofibrant objects in a nice class-combinatorial left proper model category \mathcal{K} . Suppose that the set of horns of \mathcal{F} , $\text{Hor}(\mathcal{F}) = \{A_c \rightarrow B_c \mid c \in C\}$, is a bounded set of maps (this assumption is not redundant as [11, 4.8(1)] shows). Then there is a variant of the small object argument, originated in [12], allowing to construct a strongly class-accessible localization functor L such that $\text{LEq}(L) = \text{LEq}(\mathcal{F})$. For every $X \in \mathcal{K}$ we apply the following inductive procedure. Let $X_0 = X$, and if X_{k-1} is constructed. Consider the set of all possible maps $g : A_c \rightarrow X_{k-1}$ for all $c \in C$. Define $X'_k = (\sqcup_g B_c) \coprod_{\sqcup_g A_c} X_{k-1}$ by attaching all possible maps from $\text{Hor}(\mathcal{F})$ to X_{k-1} . Then $X_k = \widehat{X'_k}$ is obtained by applying the functorial fibrant replacement in \mathcal{K} existing since \mathcal{K} is nice. We continue

this procedure while $k < \lambda$, for a fixed cardinal λ such that the domains of the maps in $\text{Hor}(\mathcal{F})$ and also the domains of the trivial cofibrations are λ -small. Put $LX = \text{colim}_{k < \lambda} X_k$. LX is fibrant, since the fibrant objects are closed under λ -filtered colimits in the class-combinatorial model category \mathcal{K} . The natural map $X \rightarrow LX$ is an \mathcal{F} -equivalence as a transfinite composition of \mathcal{F} -equivalences. One can show, similarly to Lemma 2.7, that L is a strongly class-accessible homotopy localization functor, hence Proposition 3.9 implies that the class of \mathcal{F} -equivalences is class-accessible.

The same method would be applicable to a weakly coreflective bounded class of maps \mathcal{F} if the cosolution set provided by the weak coreflectivity condition would depend functorially on the map we are about to factor. Unfortunately we do not know such examples in class-combinatorial model categories (cf. [7] for functorial examples of this kind in a non-class combinatorial model category). Of course, this condition is not necessary for a class \mathcal{F} of maps to satisfy in order to have a class accessible $\text{LEq}(\mathcal{F})$ — see Example 3.11.

Recall that if \mathcal{K} is a model category, then a functor $L : \mathcal{K} \rightarrow \mathcal{K}$ equipped with natural transformation $\eta : \text{Id}_{\mathcal{K}} \rightarrow L$ is called *homotopy idempotent* if $L\eta_K$ and η_{LK} are weak equivalences for each K in \mathcal{K} .

Definition 3.13. Let \mathcal{K} be a model category equipped with a homotopy idempotent functor $L : \mathcal{K} \rightarrow \mathcal{K}$ preserving weak equivalences. *L -fibrations* are the maps with the right lifting property with respect to cofibrations, which are also L -equivalences. A left Bousfield localization of \mathcal{K} with respect to L , or just *L -localization* of \mathcal{K} is a new model structure on \mathcal{K} such that the class of cofibrations coincides with the original class of cofibrations in \mathcal{K} and the class of weak equivalences is $\text{LEq}(L)$. New fibrations are the L -fibrations.

Theorem 3.14. *Let \mathcal{K} a nice, proper, simplicial class-combinatorial model category and $L : \mathcal{K} \rightarrow \mathcal{K}$ a strongly class-accessible homotopy idempotent functor preserving weak equivalences. Suppose additionally, that pullbacks of L -equivalences along L -fibrations are L -equivalences. Then the L -localization exists and is class-combinatorial.*

Proof. It was shown in [6, Appendix A], that the pair

$$(\text{cof}(\mathcal{I}) \cap \text{LEq}(L), (\text{cof}(\mathcal{I}) \cap \text{LEq}(L))^{\square})$$

is a weak factorization system. They argue as follows.

Take $i \in \text{cof}(\mathcal{I}) \cap \text{LEq}(L)$ and $f : X \rightarrow Y$. For any morphism $i \rightarrow f$ in \mathcal{K}^\rightarrow we perform the following construction:

$$\begin{array}{ccccc}
 A & \longrightarrow & X & \xrightarrow{\quad} & LX \\
 \downarrow i & & \downarrow f & \searrow & \downarrow \wr \\
 & & & Z & \\
 & & & \swarrow \sim & \\
 & & & P & \longrightarrow W \\
 & \nearrow & \nearrow & & \downarrow \\
 B & \longrightarrow & Y & \xrightarrow{\quad} & LY
 \end{array}$$

After applying the functor L on the morphism f we factor Lf as a trivial cofibration followed by a fibration in \mathcal{K} , obtaining the L -fibration $W \rightarrow LY$, since this is a fibration of L -local objects [6, A.8(iii)]. Then constructing $P = W \times_{LY} Y$ we obtain an L -fibration $P \rightarrow Y$ as a pull-back of an L -fibration and an L -equivalence $P \rightarrow W$ due to the additional assumption. The induced morphism $X \rightarrow P$ is an L -equivalence by the 2-out-of-3 property. Now we factor the morphism $X \rightarrow P$ into a cofibration followed by a trivial fibration in \mathcal{K} . As the composition of two L -fibrations, the morphism $Z \rightarrow Y$ is an L -fibration, hence there exists a lift $B \rightarrow Z$, showing that $\text{cof}(\mathcal{I}) \cap \text{LEq}(\mathcal{F})$ is cone-coreflective.

Like in the proof of 3.10, there exists a regular cardinal λ such that \mathcal{K} is λ -combinatorial and L strongly class- λ -accessible. Assume that X and Y are λ -presentable. Then LX, LY and W are λ -presentable. Since \mathcal{K} is locally λ -presentable simplicial category, the functor $E : \mathcal{K} \rightarrow \mathcal{P}(\mathcal{A})$ from the proof of [11] 2.6 takes values in simplicial presheaves and thus $\mathcal{P}(\mathcal{A})$ can be taken in the sense of 2.3. The functor E sends λ -presentable objects to finitely presentable ones and thus it is strongly class- λ -accessible. Since E preserves limits (see [11] 2.6), 2.4 implies that EP is λ -presentable. Since \mathcal{K} is closed under λ -filtered colimits in $\mathcal{P}(\mathcal{A})$, P is λ -presentable. Thus Z is λ -presentable. Consequently, the L -localized model category is class- λ -combinatorial. \square

Example 3.15. Take $f : V \rightarrow 1$ in Example 3.11. For such maps f -localization functor is called also V -nullification. Then the resulting class of f -equivalences satisfies the conditions of 3.14, since the nullification of spaces (i.e., the localization with respect to f for f as above) is a right proper model category (see, e.g., [5]). Hence the model category resulting from the levelwise nullification of the projective model structure on the category of small functors is class-combinatorial again.

Example 3.16. Consider the category $\mathcal{P}(\mathbf{SSet}^{\text{op}})$ of small simplicial functors from simplicial sets to simplicial sets equipped with the projective model structure (see 2.5). Consider the localization functor $L : \mathcal{P}(\mathbf{SSet}^{\text{op}}) \rightarrow \mathcal{P}(\mathbf{SSet}^{\text{op}})$, $L = P_n \circ \text{Fib}$ constructed in [3], where P_n is Goodwillie's n -th polynomial approximation [17] and $\text{Fib} : \mathcal{P}(\mathbf{SSet}^{\text{op}}) \rightarrow \mathcal{P}(\mathbf{SSet}^{\text{op}})$ is the strongly class-accessible fibrant replacement functor from 2.7. Since P_n is a countable colimit of finite homotopy limits of cubical diagrams applied on homotopy pushouts (joins with finite sets used to construct P_n in [17] may be expressed as homotopy pushouts), it is strongly class-accessible. Thus L is strongly class-accessible, hence the polynomial model structure constructed in [3] is class-combinatorial.

The condition on the localization functor to be strongly class-accessible may not be omitted in 3.14 as the following example shows.

Example 3.17. The following localization of the class-combinatorial model category $\mathcal{P}(\mathbf{SSet})$ was constructed in [9]. The localization functor $L : \mathcal{P}(\mathbf{SSet}) \rightarrow \mathcal{P}(\mathbf{SSet})$ is the composition of the evaluation functor at the one point space $\text{ev}_*(F) = F(1)$ with a fibrant replacement $\widehat{(-)}$ in simplicial sets (chosen to be a small functor like $\widehat{\text{Id}}_{\mathcal{S}}$ or Ex^∞) and the Yoneda embedding $Y : \mathbf{SSet} \rightarrow \mathcal{P}(\mathbf{SSet})$, i.e., $L(F) = \text{hom}(-, \widehat{F(1)})$. This localization functor satisfies the conditions of A6 in [6] (pullback of an L -equivalence along an L -fibration is an L -equivalence again), and hence there exists the L -local model structure on $\mathcal{P}(\mathbf{SSet})$. The fibrant objects in the localized model category are levelwise fibrant functors weakly equivalent to representable functors, but they are not closed under filtered colimits, since filtered colimit of representable functors need not be representable, no matter how large the filtered colimit is. On the other hand, in a class-cofibrantly generated model category sufficiently large filtered colimits of fibrant objects are fibrant again. In other words, we obtain a localization of $\mathcal{P}(\mathbf{SSet})$, which is not class-cofibrantly generated. The reason is that the localization functor L is not class-accessible. See [9] for more details on this model structure.

REFERENCES

- [1] J. Adámek and J. Rosický, *Locally Presentable and Accessible Categories*, Cambridge University Press 1994.
- [2] T. Beke, *Sheafifiable homotopy model categories*, Math. Proc. Cambr. Phil. Soc. 129 (2000), 447–475.
- [3] G. Biedermann, B. Chorny and O. Röndigs, *Calculus of functors and model categories*, Advances in Mathematics 214 (2007), 92–115.

- [4] F. Borceux, *Handbook of Categorical Algebra 2*, Cambridge Univ. Press 1994.
- [5] A. K. Bousfield, *On the telescopic homotopy theory of spaces*, Trans. Amer. Math. Soc. 353 (2001), 2391–2426.
- [6] A. K. Bousfield and E. M. Friedlander, *Homotopy theory of Γ -spaces, spectra, and bisimplicial sets*, In Geometric Applications of Homotopy Theory II, Lect. Notes in Math. 658, Springer-Verlag 1978.
- [7] B. Chorny, *Localization with respect to a class of maps I. Equivariant localization of diagrams of spaces*, Israel Journal of Math. 147 (2005), 99–140.
- [8] B. Chorny, *A generalization of Quillen’s small object argument*, J. Pure Appl. Alg. 204, Issue 3 (2006), 568–583.
- [9] B. Chorny, *Brown representability for space-valued functors*, Israel J. of Math., to appear.
- [10] B. Chorny and W. G. Dwyer, *Homotopy theory of small diagrams over large categories*, Forum Math. 21 (2009), 167–179.
- [11] B. Chorny and J. Rosický, *Locally class-presentable and class-accessible categories*, preprint 2011.
- [12] D. Christensen and D. Isaksen, *Duality and pro-spectra*, Algebr. Geom. Topol. 4 (2004), 781–812.
- [13] B. Day and S. Lack, *Limits of small functors*, J. Pure Appl. Alg. 210 (2007), 651–663.
- [14] D. Dugger, *Universal homotopy theories*, Adv. Math. 164 (2001), 144–176.
- [15] D. Dugger, *Combinatorial model categories have presentations*, Adv. Math. 164 (2001), 177–201.
- [16] E. Dror Farjoun, *Homotopy theories for diagrams of spaces*, Proc. Amer. Math. Soc. 101 (1987), 181–189.
- [17] T. G. Goodwillie, *Calculus III: Taylor Series*, Geometry & Topology, 7 (2003), 645–711.
- [18] P.G. Goerss and J.F. Jardine, *Simplicial Homotopy theory*, Birkhäuser 1999.
- [19] P.S. Hirschhorn, *Model Categories and their Localizations*, Amer. Math. Soc. 2003.
- [20] M. Hovey, *Model categories*, Amer. Math. Soc. 1999.
- [21] D. Isaksen, *A model structure on the category of pro-simplicial sets*, Trans. Amer. Math. Soc. 353 (2001), 2805–2841.
- [22] P. Karazeris, J. Rosický and J. Velebil, *Completeness of cocompletions*, J. Pure Appl. Alg. 196 (2005), 229–250.
- [23] G. M. Kelly, *Basic concepts of enriched category theory*, Cambridge University Press 1982.
- [24] G. M. Kelly, *Structures defined by finite limits in the enriched context, I.*, Cahiers Top. Géom. Diff. XXIII (1982), 3–42.
- [25] J. Lurie, *Higher Topos theory*, Princeton Univ. Press 2009.
- [26] M. Makkai and R. Paré, *Accessible Categories: The Foundation of Categorical Model Theory*, Cont. Math. 104, AMS 1989.

- [27] J. Rosický, *Generalized Brown representability in homotopy categories*, Th. Appl. Cat. 14 (2005), 451-479.
- [28] J. Rosický, *On combinatorial model categories*, Appl. Cat. Str. 17 (2009), 303-316.
- [29] M. Shulman. *Homotopy limits and colimits and enriched homotopy theory*, arXiv:math.AT/0610194.

B. CHORNY
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF HAIFA AT ORANIM
TIVON, ISRAEL
CHORNY@MATH.HAIFA.AC.IL

J. ROSICKÝ
DEPARTMENT OF MATHEMATICS AND STATISTICS
MASARYK UNIVERSITY, FACULTY OF SCIENCES
BRNO, CZECH REPUBLIC
ROSICKY@MATH.MUNI.CZ