

MILD MANIFOLDS AND A NON-STANDARD RIEMANN EXISTENCE THEOREM

YA'ACOV PETERZIL AND SERGEI STARCHENKO

ABSTRACT. Let \mathcal{R} be an o-minimal expansion of a real closed field R and K be the algebraic closure of R . In earlier papers we investigated the notions of \mathcal{R} -definable K -holomorphic maps, K -analytic manifolds and their K -analytic subsets. We call such a K -manifold mild if it eliminates quantifiers after endowing it with all K -analytic subsets. Examples are compact complex manifolds and non-singular algebraic curves over K .

We examine here basic properties of mild manifolds and prove that when a mild manifold M is strongly minimal and not locally modular then M is bi-holomorphic with a non-singular algebraic curve over K .

1. INTRODUCTION

For a complex-analytic manifold M , let \mathcal{M} be the first-order structure whose underlying set is M and basic relations are all complex-analytic subsets of Cartesian powers of M .

The following result is due to Zilber (see [17]).

Theorem 1.0.1. *If M is a compact complex-analytic manifold, then \mathcal{M} has elimination of quantifiers and has finite Morley rank. If \mathcal{M} is strongly minimal then it is a Zariski geometry (see [8] for a definition of Zariski geometries).*

Let \mathcal{R} be a fixed o-minimal expansion of a real closed field R , and let $K = R(\sqrt{-1})$ be its algebraic closure identified naturally with R^2 . In a series of papers [12–14] we investigated notions K -holomorphic maps and K -analytic manifolds definable in the structure \mathcal{R} . For a K -analytic manifold M let $\mathcal{A}(M)$ be the first order structure whose underlying set is M and basic relations are all K -analytic subsets of Cartesian powers of M (see Section 2.2 for definitions of K -analytic manifolds and K -analytic subsets). In paper [14] we showed that a version of the above Zilber's theorem holds in this general setting.

Date: January 30, 2008.

2000 Mathematics Subject Classification. Primary 03C64, 03H05, 32P05; Secondary 03C98, 30F99, 32C25.

Key words and phrases. O-minimality, Non-Archimedean complex analysis, Zariski geometries, Riemann existence theorem.

The second author was partially supported by the NSF.

Theorem 1.0.2 ([14, Theorem 8.3]). *If M is a definably compact K -manifold, then $\mathcal{A}(M)$ has elimination of quantifiers and has finite Morley rank. If $\mathcal{A}(M)$ is strongly minimal then it is a Zariski geometry.*

If we take \mathcal{R} to be any o-minimal expansion of \mathbb{R}_{an} , then every compact complex manifold can be viewed as an \mathcal{R} -definable object, and the structures \mathcal{M} and $\mathcal{A}(M)$ are the same (see [11] for more details). Thus Theorem 1.0.2 can be considered as a proper generalization of Zilber's theorem.

Notice that the assumption of compactness is essential in Zilber's theorem. For example, if we take $M = \mathbb{C}$ then the structure \mathcal{M} is not even stable (since the graph of complex exponential function is a complex-analytic subset of \mathbb{C}^2 , we can use it to define integers in this structure.) However if we take \mathcal{R} to be any o-minimal expansion of the field of real numbers, then only complex analytic subsets of \mathbb{C}^n definable in the structure \mathcal{R} are algebraic (see [14, Theorem 5.1]), and we see a big difference between \mathcal{M} and $\mathcal{A}(M)$ in this case. The structure \mathcal{M} is unstable, and the structure $\mathcal{A}(M)$ is strongly minimal and inderdefinable with the field of complex numbers.

Motivated by this and other examples we define a K -analytic manifold M to be *mild* if the structure $\mathcal{A}(M)$ has quantifier elimination. In particular, every definably compact K -manifold is mild, but there are also many mild manifolds that are not definably compact, e.g. any non-singular algebraic affine variety over K .

Our main goal in this paper is to prove the following theorem (see Theorem 6.0.5 below).

Theorem 1.0.3. *Let M be a strongly minimal mild K -manifold (i.e. M is mild and the structure $\mathcal{A}(M)$ is strongly minimal). Then either the structure $\mathcal{A}(M)$ is locally modular or M is K -biholomorphic with an algebraic curve over K .*

One of obstacles in proving the above theorem is that in general Riemann existence theorem does not hold for mild manifolds:¹ In paper [15] we considered a family of generalized 1-dimensional tori $\mathcal{E}_{\tilde{\tau}}$ and showed that for a sufficiently general nonstandard $\tilde{\tau}$ the structure $\mathcal{A}(\mathcal{E}_{\tilde{\tau}})$ is strongly minimal but $\mathcal{E}_{\tilde{\tau}}$ is not K -biholomorphic to an algebraic curve. Thus, by the above theorem, $\mathcal{A}(\mathcal{E}_{\tilde{\tau}})$ is locally modular.

To overcome this obstacle we prove in this paper that the second part of Riemann existence theorem holds for o-minimal expansions of real closed fields. Namely, every definable finite covering of a co-finite subset of K is definably isomorphic to an algebraic covering (see Theorem 5.1.3 below). In a semialgebraic category this theorem can be restated as follows.

Theorem 1.0.4. *Let R be a real closed field and $K = R(\sqrt{-1})$ be its algebraic closure identified with R^2 . Let $B \subseteq R^2$ be a co-finite subset, $C \subseteq R^m$ an R -semialgebraic set, and $\pi: C \rightarrow B$ an R -semialgebraic finite covering. Then there*

¹Due to a theorem of R. Moosa (see [10]) Riemann existence theorem holds in elementary extensions of \mathcal{M} for a compact complex manifold M .

is a nonsingular algebraic affine curve Γ over K , a regular morphism $\tau: \Gamma \rightarrow K$, and an R -semialgebraic homeomorphism $f: C \rightarrow \Gamma$ such that $\pi = \tau \circ f$.

1.1. The structure of the paper. In Section 2 we review K -manifolds and K -analytic subsets.

In Section 3 we consider the field of K -meromorphic functions $Mer(M)$ on a K -manifold M , and establish a connection between finite coverings and field extensions.

In Section 4 we define mild manifolds. The main result of this section is Theorem 4.3.3 stating that if M is a strongly minimal mild manifold that is not locally modular then there is a nonconstant K -meromorphic function $f: M \rightarrow K$. It implies that, up-to finite subsets, every non locally modular strongly minimal mild manifold is a finite covering of K .

In Section 5 we show that the second part of Riemann existence theorem holds in \mathfrak{o} -minimal expansions of real closed fields. Since proofs of basic properties of definable finite coverings are analogous to classical proofs we present them in Appendix.

In Section 6 we prove Theorem 6.0.5

We would like to thank Elias Baro for pointing out a gap in one of the proofs.

2. PRELIMINARIES

For the rest of this paper we fix an \mathfrak{o} -minimal expansion \mathcal{R} of a real closed field R , and will denote by K its algebraic closure $R(\sqrt{-1})$.

We briefly review the basic notions of analytic geometry with respect to the field K and refer to [12–14] for more details.

2.1. K -holomorphic maps. The theory is based on the following straightforward analogue to the classical definition of a holomorphic map: Given a definable open set $U \subseteq K$ we call a definable function $f: U \rightarrow K$ *K -holomorphic* if for every $z \in U$ the limit

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$$

exists in K (we identify K with \mathbb{R}^2 in an obvious manner just like the complex field is identified with \mathbb{R}^2).

A definable map from a definable open subset $U \subseteq K^n$ into K^m is called *K -holomorphic* if each of its coordinate functions is continuous and K -holomorphic in each variable separately.

Notice that in the classical setting, i.e. when R is the field of real numbers, the above definitions amount to saying that the function f is definable in the underlying \mathfrak{o} -minimal structure and is complex-holomorphic in the classical sense.

Let $U \subseteq K^n$ is a definable open set, $f: U \rightarrow K^m$ a K -holomorphic map, and $a \in U$. The K -differential df_a of f at a is a K -linear map from K^n into K^m . We denote its rank by $\text{Rnk}_K(f)_a$ and call it *the K -rank of f at a* (or just a rank of f at a if no confusion arises).

2.2. K -manifolds. A definable n -dimensional K -manifold is a definable set M , equipped with a finite cover of definable sets $M = \bigcup_i U_i$, each of which is in definable bijection with an open subset of K^n such that the transition maps are K -holomorphic maps between open subsets of K^n . As usual, K -holomorphic maps between K -manifolds are defined through the charts of the manifolds.

Let M, N be K -manifolds, $U \subseteq M$ be a definable open subset, $f: U \rightarrow N$ a K -holomorphic map, and $a \in U$. The K -rank of f at a is defined using the charts. By the Chain Rule, this rank is independent of charts chosen.

Let N be a K -manifold of K -dimension n . A definable subset $M \subseteq N$ is called a d -dimensional K -submanifold if for every $a \in M$ there is a definable open set $U \subseteq N$ containing a and a K -holomorphic $f: U \rightarrow K^{n-d}$ such that $M \cap U = f^{-1}(0)$ and $\text{Rnk}_K(f)_a = n - d$. As was shown in [14, Lemma 3.3], every such submanifold admits a finite atlas, which makes it into a K -manifold on its own.

For example, if $X \subseteq K^n$ is a non-singular affine variety over K then X is a K -submanifold of K^n .

2.3. K -analytic sets. A K -analytic subset of a K -manifold M is a definable set $A \subset M$, such that at every point $z \in M$, the set A is given, locally near z , as the zero set of finitely many K -holomorphic functions. Again, in the classical setting a \mathbb{C} -manifold and a \mathbb{C} -analytic subset are just a complex manifold and a complex analytic subset, respectively, which are definable in the underlying o-minimal structure over \mathbb{R} .

The K -dimension of a K -analytic set A is defined to be the maximal d such that A contains a d -dimensional K -manifold. We use $\dim_K A$ to denote the dimension of A as a K -analytic set and $\dim_{\mathcal{R}} A$ to denote its o-minimal dimension. We clearly have $\dim_{\mathcal{R}} A = 2 \dim_K A$.

Let M be a K -manifold, $A \subset M$ a K -analytic subset of M and $X \subset A$. The set X is a K -analytic subset of A if it is K -analytic in M . The complement $A \setminus X$ of a K -analytic subset $X \subset A$ is called *Zariski open in A* .

Notice that, by Theorem 2.3.2 below, for an algebraic K -variety the above notion of a Zariski open subset coincides with the standard algebraic notion, so no confusion should arise.

If A is a K -analytic subset of a K -manifold M then we denote by $\text{Reg}_K(A)$ the set of K -regular points of A , i.e. the set of all $a \in A$ such that A is a K -submanifold of M near A . As in the classical setting, the set $\text{Reg}_K(A)$ is Zariski open and dense in A (see [14, 6.1]).

A K -analytic subset $A \subseteq M$ of a K -analytic manifold M is called *irreducible* if it can not be written as $A = A_1 \cup A_2$, where A_1, A_2 are K -analytic subset of M and none of A_i 's is contained in the other. If $A \subset M$ is irreducible $X \subset A$ a proper K -analytic subset of A then $\dim_K(X) < \dim_K(A)$ and in particular the topological closure of $A \setminus X$ is A ([14, Corollary 4.10]). We also have:

Theorem 2.3.1 ([14, 4.6 and 4.12]). *Every K -analytic subset A of a K -manifold M is a finite union of irreducible K -analytic subsets A_i such that each A_i is not contained in any other. (These are the irreducible components of A .) Moreover these irreducible components are exactly the closures of the definably connected components of $\text{Reg}_K(A)$.*

While the results so far are mostly analogues of the classical results, the theorems below are false without the o-minimality assumptions (see [14]).

The first result can be considered as a generalization of the classical Chow's lemma.

Theorem 2.3.2 ([14, 5.1]). *Let A be a K -analytic subset of K^n or of $\mathbb{P}^n(K)$. Then A is an algebraic variety over K .*

The following theorem on removal of singularities will be used here extensively:

Theorem 2.3.3 ([14, 4.13]). *Let M be a K -manifold and $E \subset M$ a K -analytic subset of M . If A is a K -analytic subset of $M \setminus E$ then its topological closure in M is a K -analytic subset of M .*

The theorem below follows from Theorem 2.3.3 and the fact that $\text{Reg}_K(A)$ is Zariski open in A .

Theorem 2.3.4. *Let $A \subset M$ be an irreducible K -analytic subset of a K -manifold M . A set $X \subset \text{Reg}_K(A)$ is a K -analytic subset of the K -manifold $\text{Reg}_K(A)$ if and only if there is a K -analytic subset Y of M such that $X = Y \cap \text{Reg}_K(A)$.*

Finally, as is shown in [14, Corollary 4.14], if A is a K -analytic subset of a K -manifold then it remains K -analytic in elementary extensions of \mathcal{R} .

2.4. More on ranks of K -holomorphic maps. The following two theorems follow from properties of \mathcal{R} -differentiable maps in o-minimal structures (see for example [3, chapter7]).

Theorem 2.4.1. *Let $f: M \rightarrow N$ be a K -holomorphic map between K -manifolds. Then*

- (1) *If r is the maximum rank of f on M then $\dim_{\mathcal{R}}(f(M)) = 2r$ (we use \mathcal{R} -dimension for $f(M)$ because it is not a K -analytic set in general). Assume that f is of constant rank r on M . Then*
- (2) *If $n = \dim_K(M)$ then for each $b \in f(M)$, $f^{-1}(b)$ is a K -manifolds of dimension $n - r$;*
- (3) *Each $a \in M$ has a definable open neighborhood U such that $f(U)$ is a K -submanifold of N of dimension r , In particular, if $\dim_K(N) = r$ then $f(M)$ is an open subset of N .*

It is easy to see that if $f: M \rightarrow N$ is a K -analytic map between K -manifolds and $k \in \mathbb{N}^{>0}$ then the set of all points $a \in M$ where the rank of f is less than k is K -analytic subset of M . It follows that f has constant rank on a nonempty Zariski open subset of M .

2.5. Meromorphic maps. A K -meromorphic map from a K -manifold M into a K -manifold N , is a definable K -holomorphic map f from a Zariski open dense subset $\text{dom}(f)$ of M into N . As usual, we will identify two K -meromorphic maps f, g from M to N if $f(x) = g(x)$ for all $x \in \text{dom}(f) \cap \text{dom}(g)$.

Remark 2.5.1. In paper [14], in the definition of a K -meromorphic map, we also required that the closure of the graph of f in $M \times N$ is a K -analytic subset. However it was proved there (Corollary 9.1) that this extra assumption is not needed.

3. THE FIELD OF MEROMORPHIC FUNCTIONS

If M is a definably connected K -manifold then it is not hard to see that the set of all K -meromorphic functions $\varphi: M \rightarrow K$ form a field under point-wise addition and multiplication. We will denote this field by $\text{Mer}(M)$. Identifying K with the constant functions on M we will always consider $\text{Mer}(M)$ as a field extension of K .

Let $f: M \rightarrow N$ be a K -meromorphic map between definably connected K -manifolds such that the range of f is dense in N . It is not hard to see that for any K -meromorphic function $\varphi: N \rightarrow K$ the function $\varphi \circ f$ is a K -meromorphic function on M . Thus we have the map $\varphi \mapsto \varphi \circ f$ from $\text{Mer}(N)$ to $\text{Mer}(M)$. We will denote this map by f^* . Obviously, $f^*: \text{Mer}(N) \rightarrow \text{Mer}(M)$ is a field embedding.

3.1. K -meromorphic finite coverings. Recall that a continuous map between topological spaces $f: X \rightarrow Y$ is called an n -covering map, where $n \in \mathbb{N}^{>0}$, if every $b \in Y$ has an open neighborhood V such that $f^{-1}(V)$ consists of n pairwise disjoint sets U_1, \dots, U_n and f maps each U_i homeomorphically onto V . We call it a *finite covering map* if it is an n -covering map for some n . The following is immediate.

Claim 3.1.1. A continuous map between Hausdorff topological spaces $f: X \rightarrow Y$ is an n -covering if and only if f is a local homeomorphism and $|f^{-1}(b)| = n$ for any $b \in Y$.

Definition 3.1.2. A K -meromorphic map $f: M \rightarrow N$ between K -manifolds is a *K -meromorphic n -covering map* if there is a Zariski open dense subset U of M and a Zariski open dense subset V of N such that that $U \subset \text{dom}(f)$ and $f|_U: U \rightarrow V$ is an n -covering.

It is called a *K -meromorphic finite covering* if it is a K -meromorphic n -covering map for some $n \in \mathbb{N}^{>0}$.

Remark 3.1.3. It is not hard to see that if $f: M \rightarrow N$ is a K -meromorphic finite covering then M and N have the same dimension.

Theorem 3.1.4. Let M, N be definably connected K -manifolds. If $f: M \rightarrow N$ is a K -meromorphic n -covering then $\text{Mer}(M)$ is a finite extension of $f^*(\text{Mer}(N))$ of degree at most n .

Proof. By the Primitive Element Theorem, it is sufficient to show that for any $\alpha(x) \in \text{Mer}(M)$ there are $c_1(y), \dots, c_n(y) \in \text{Mer}(N)$ and a Zariski open dense $U \subseteq M$ such that

$$\alpha^n(x) + c_1(f(x))\alpha^{n-1}(x) + \dots + c_{n-1}(f(x))\alpha(x) + c_n(f(x)) = 0$$

for all $x \in U$. Obviously, we need to consider only non-constant α .

Let $U \subseteq M$ and $V \subseteq N$ be Zariski open dense subsets such that $f: U \rightarrow V$ is an n -covering. Decreasing U and V if needed we can assume that $\alpha(x)$ is K -holomorphic on U .

Using definable choice we can find definable $\xi_1, \dots, \xi_n: V \rightarrow U$ such that $f^{-1}(y) = \{\xi_1(y), \dots, \xi_n(y)\}$ for all $y \in V$.

For $i = 1, \dots, n$, let $s_i(t_1, \dots, t_n)$ be the i -th basic symmetric polynomial in n variables.

Consider functions c_i from V into K , defined as

$$c_i(y) = (-1)^i s_i(\alpha(\xi_1(y)), \dots, \alpha(\xi_n(y))).$$

Using the symmetry of s_i 's, it is not difficult to see that each c_i is K -holomorphic on V (see [13, bottom of p.16] for a similar argument).

For $x \in U$ and $y = f(x)$ we have $\prod_{i=1}^n (x - \alpha(\xi_i(y))) = 0$.

Thus

$$\alpha^n(x) + c_1(f(x))\alpha^{n-1}(x) + \dots + c_{n-1}(f(x))\alpha(x) + c_n(f(x)) = 0$$

on U . □

4. THE STRUCTURE $\mathcal{A}(M)$ AND MILD MANIFOLDS

Let M be a K -manifold. We denote by $\mathcal{A}(M)$ the first order structure whose universe is M and basic relations are K -analytic subsets of Cartesian powers of M .

Remark 4.0.5. The structure $\mathcal{A}(M)$ depends, by definition, on the ambient o-minimal structure \mathcal{R} (because the definition of a K -analytic set depends on that structure).

Let \mathcal{R}^* be an $|R|^+$ -saturated elementary extension of \mathcal{R} . We denote by R^* the universe of \mathcal{R}^* , by K^* its algebraic closure and for an \mathcal{R} -definable subset $X \subseteq R^n$ we denote by X^* the set of its \mathcal{R}^* -points. It is easy to see that M^* is a K^* -manifold, and as was pointed out earlier, if A is a K -analytic subset of M then A^* is K^* -analytic subset of M^* . We let $\mathcal{A}(M^*)$ denote the first order structure whose universe is M^* and basic sets are K^* -analytic subsets of Cartesian powers of M^* .

It is easy to see that there is a reduct \widetilde{M} of $\mathcal{A}(M^*)$ such that \widetilde{M} is an $|M|^+$ -saturated elementary extension of $\mathcal{A}(M)$. However, as is shown in [13], the structures $\mathcal{A}(M^*)$ and \widetilde{M} may differ, in terms of their definable sets. For more on the relationship between $\mathcal{A}(M^*)$ and \widetilde{M} , in the case where $\mathcal{R} = \mathbb{R}_{an}$, see [11].

4.1. Mild manifolds. Our notion of a mild manifold was inspired by Zilber's observation that compact complex manifolds with their analytic structure, eliminate quantifiers (see [20]).

Definition 4.1.1. We say that a K -manifold M is *mild* if the structure $\mathcal{A}(M)$ admits quantifier elimination.

If $\mathcal{A}(M)$ is mild then every definable $X \subset M^k$ can be written as $X = \bigcup_{i=1}^k (A_i \setminus B_i)$, where $A_i \subset B_i$ are K -analytic subsets of M^k and each A_i is irreducible. It follows that the topological closure of X is the union of the A_i 's and therefore K -analytic itself. It is not difficult now to conclude:

Theorem 4.1.2. *Let M be a mild K -manifold, $A \subset M^n$ a K -analytic irreducible subset, and $\pi: M^n \rightarrow M^k$ a projection. Then the topological closure $\text{cl}(\pi(A))$ of $\pi(A)$ is an irreducible K -analytic subset of M^k and $\text{cl}(\pi(A)) \setminus F \subset \pi(A)$ for some K -analytic subset $F \subsetneq \text{cl}(\pi(A))$.*

Example 4.1.3. According to [14, Theorem 8.3], every definably compact K -manifold (and in particular every compact complex manifold) is mild.

The example above together with the theorem below provide us with a wide class of mild K -manifolds.

Theorem 4.1.4. *Let M be a mild K -manifold and $A \subset M$ a K -analytic subset of M . Then*

- (i) $M \setminus A$ is a mild manifold.
- (ii) If A is irreducible then $\text{Reg}_K(A)$ is a mild K -manifold.

Proof. (i) Follows from Theorem 2.3.3, while (ii) follows from Theorem 2.3.4. \square

Example 4.1.5. As the theorem below shows, the unit disk $D \subseteq K$ is not a mild K -manifold, in any o-minimal structure over R .

Claim 4.1.6. *For a definable open nonempty subset $U \subseteq K$ the following conditions are equivalent.*

- (1) U is co-finite in K .
- (2) U is a mild K -manifold.
- (3) The structure $\mathcal{A}(U)$ is strongly minimal.

Proof. (1) \Rightarrow (2). If $U \subset K$ is co-finite then, by Theorems 2.3.2 and 2.3.3, the structure $\mathcal{A}(U)$ is the same as the structure induced on U by algebraic sets, hence it is strongly minimal.

(2) \Rightarrow (3). Since $\dim_K(U) = 1$ all proper K -analytic subsets of U are finite. By quantifier elimination, every $\mathcal{A}(U)$ -definable subset of U is either finite or co-finite. Since o-minimal structures have uniform finiteness property, the structure $\mathcal{A}(U)$ is strongly minimal.

(3) \Rightarrow (1). Assume now that U is not co-finite in K and let $X = K \setminus U$. Obviously, for any $c \in K$ the them map $x \mapsto c + x$ is a bijection between U and $c + U$ that

induces bijection between $\mathcal{A}(U)$ -definable sets and $\mathcal{A}(c+U)$ definable sets. Thus we may assume $0 \in U$. For every $a \in K$ the set $\{(x, y) \in U \times U : x = ay\}$ is a K -analytic subset of $U \times U$, hence the set $aU \cap U$ is definable in $\mathcal{A}(U)$. If we take a close enough to 0 so that $aX \cap U$ is infinite, we obtain that $aU \cap U$ is infinite and co-infinite. Hence $\mathcal{A}(U)$ is not strongly minimal. \square

We need the following technical result.

Lemma 4.1.7. *Let M be a mild K -manifold, A an irreducible K -analytic subset of M^n , $\pi: M^n \rightarrow M^k$ a projection such that $\pi(A)$ is infinite, and let B be the topological closure of $\pi(A)$ in M^k . Then there is a Zariski open dense subset U of $\text{Reg}_K(A)$ such that π has a constant rank on U and $\pi(U)$ is a Zariski open dense subset of $\text{Reg}_K(B)$.*

Proof. Let r be the maximal rank of f , restricted to the manifold $\text{Reg}_K(A)$. As was pointed out earlier, there exists a Zariski open dense subset $U_0 \subseteq \text{Reg}_K(A)$ such that π has constant rank r on U_0 . The set $\pi(U_0)$ is dense in B hence, by quantifier elimination, $\pi(U_0)$ contains a Zariski open dense subset W of B . We can now take $U = \pi^{-1}(W \cap \text{Reg}_K(B))$. \square

4.1.1. *Some open questions on mild manifolds.* Claim 4.1.6 suggests an immediate question:

Question. Is it true that a definable open $U \subset K^n$ is mild iff it is Zariski open in K^n ?

Question. Let M be a K -manifold. Assume that $\mathcal{A}(M)$ is strongly minimal, or even stable. Is M necessarily mild?

Question. Let M, N be mild K -manifolds. Is $M \times N$ necessarily mild?

Let $\mathcal{R}^* \succ \mathcal{R}$, R^* , K^* and \widetilde{M} and $\mathcal{A}(M^*)$ be as in Remark 4.0.5. Notice that if $\mathcal{A}(M)$ is mild then the structure \widetilde{M} eliminates quantifiers. However, one can still ask:

Question Let \mathcal{R}^* , M^* , $\mathcal{A}(M^*)$ and \widetilde{M} be as in Remark 4.0.5. Assume that $\mathcal{A}(M)$ is mild.

1. Is the structure $\mathcal{A}(M^*)$ mild as well?
2. Does $\mathcal{A}(M)$ remain mild in every o-minimal expansion of the structure \mathcal{R} ?

4.2. **Generic Types.** Let M be a mild K -manifold. If A is an irreducible K -analytic subset of M^n then, by quantifier elimination, in the structure $\mathcal{A}(M)$ the set $\{U \subset A : U \text{ is Zariski open and dense in } A\}$ is contained in a unique complete n -type over M . As usual we will call this type a *generic type of A* . Conversely, if p is a complete n -type over M and $A \in p$ is an irreducible K -analytic subset of minimal K -dimension then p is the generic type of A .

4.3. Strongly minimal mild K -manifolds.

Definition 4.3.1. A K -manifold M is called a *strongly minimal mild K -manifold* if it is mild and the structure $\mathcal{A}(M)$ is strongly minimal.

Since every o-minimal structure has uniform bounds on definable families of finite sets, it follows that a mild K -manifold M is strongly minimal if and only if every proper K -analytic subset of M is finite. In particular, every definably connected definably compact 1-dimensional K -manifold is strongly minimal and mild.

Theorem 4.3.2. *If M is a strongly minimal mild manifold then the structure $\mathcal{A}(M)$ is a Zariski geometry with K -analytic subsets taken as closed ones.*

Proof. Using the dimension theorem [14, Theorem 7.3], it is easy to see that strongly minimal mild manifolds with K -analytic subsets satisfy the axioms of Zariski geometries from [8]. \square

Our main goal in this section is to prove the following theorem.

Theorem 4.3.3. *Let M be a mild strongly minimal K -manifold. The structure $\mathcal{A}(M)$ is locally modular if and only if $\text{Mer}(M) = K$.*

We will need several claims.

Claim 4.3.4. *Let M be a strongly minimal mild K -manifold. Then for any $n \in \mathbb{N}^{>0}$ the generic type of M^n is the unique n -type in $\mathcal{A}(M)$ whose Morley rank is n .*

Proof. Easy, by induction on n . \square

Claim 4.3.5. *Let M be a strongly minimal mild K -manifold and $A \subset M^n$ an irreducible K -analytic subset. Then there is a projection $\pi: M^n \rightarrow M^k$ such that $\pi|_{\text{Reg}_K(A)}: \text{Reg}_K(A) \rightarrow M^k$ is a K -meromorphic finite covering.*

Proof. Let $p(\bar{x})$ be the generic type of A . Notice that the formula $x \in \text{Reg}_K(A)$ belongs to p . Let $\bar{a} = (a_1, \dots, a_n)$ be a realization of p in some elementary extension \widetilde{M} of $\mathcal{A}(M)$ (see discussion in Remark 4.0.5).

Let k be the Morley rank of p . After permuting coordinates if needed, we may assume that in the structure \widetilde{M} , the elements a_1, \dots, a_k are algebraically independent over M and \bar{a} is in the algebraic closure of a_1, \dots, a_k . Let $\pi: M^n \rightarrow M^k$ be the projection on-to the first k -coordinates.

By Claim 4.3.4, (a_1, \dots, a_k) realizes the generic type of M^k . Thus there is $m \in \mathbb{N}^{>0}$ such that the set

$$V_0 = \{x \in M^k: |\pi^{-1}(x) \cap \text{Reg}_K(A)| = m\}$$

is Zariski open and dense in M^k .

Let $U_0 = \pi^{-1}(V_0) \cap \text{Reg}_K(A)$. Obviously it is a Zariski open subset of A . By Theorem 4.1.7 there is a Zariski open dense $U \subset U_0$ such that $\pi|_U$ has constant rank and $V = \pi(U)$ is a Zariski open dense subset of M^k . It is easy to see that $\pi|_U: U \rightarrow V$ is a K -holomorphic m -covering, and therefore $\pi|_{\text{Reg}_K(A)}: \text{Reg}_K(A) \rightarrow M^k$ is a K -meromorphic finite covering. \square

Claim 4.3.6. *Let M be a strongly minimal mild K -manifold and $f: M \rightarrow K$ a non-constant K -meromorphic function. Then $f: M \rightarrow K$ is a K -meromorphic finite covering. In particular $\dim_K(M) = 1$.*

Proof. Let $U \subset M$ be a Zariski open set on which f is K -holomorphic. By the strong minimality of $\mathcal{A}(M)$, the set $M \setminus U$ is finite. Because f is nonconstant, the set $f(U)$ is infinite. It follows that the set $\{u \in U: df(u) = 0\}$ is a proper K -analytic subset of U which, by 2.3.4, is also definable in $\mathcal{A}(M)$, hence must be co-finite. Shrinking U , if needed, we may assume that the differential of f is nowhere zero on U .

Consider the equivalence relation E on U defined as $xEy \iff f(x) = f(y)$. Clearly, E is a K -analytic subset of $U \times U$ and, by Theorem 2.3.3, it is definable in $\mathcal{A}(M)$. By strong minimality, almost all E -classes have the same (finite) size. Shrinking U again, if needed, we may assume that all E -classes have the same size n . In particular, we already see that $\dim_K M = \dim_K K = 1$, and by our assumption on df , the map $f: U \rightarrow f(U)$ is a n -covering map. We are left to show that $f(U)$ is Zariski open and dense in K , i.e. is co-finite in K .

Since for any K -analytic subset $A \subseteq f(U)$ the closure of $f^{-1}(A)$ in M is K -analytic in M we obtain that the structure $\mathcal{A}(f(U))$ is isomorphic to a reduct of the structure on U/E induced by $\mathcal{A}(M)$. Since $\mathcal{A}(M)$ is strongly minimal, it implies that $\mathcal{A}(f(U))$ is strongly minimal as well, and, by the Claim 4.1.6, $f(U)$ is co-finite in K . \square

As a corollary we obtain one of the directions in Theorem 4.3.3.

Corollary 4.3.7. *Let M be a mild strongly minimal K -manifold. If $\text{Mer}(M) \neq K$ then the structure $\mathcal{A}(M)$ is not locally modular.*

Proof. If $\text{Mer}(M) \neq K$ then, by the previous claim, there is a K -meromorphic finite covering $f: M \rightarrow K$. It is easy to see that the field K is interpretable in $\mathcal{A}(M)$, hence $\mathcal{A}(M)$ is not locally modular. \square

4.3.1. Proof of Theorem 4.3.3. In this section we prove the second direction of Theorem 4.3.3. We fix a strongly minimal mild K -manifold M with $\text{Mer}(M) = K$ and will show that $\mathcal{A}(M)$ is locally modular.

Claim 4.3.8. (1) *For any $n \in \mathbb{N}^{>0}$ we have $\text{Mer}(M^n) = K$*
 (2) *For any irreducible K -analytic subset $A \subset M^n$, we have $\text{Mer}(\text{Reg}_K(A)) = K$.*

Proof. (1) is easy.

(2) By Theorem 4.3.5 there is a projection $\pi: M^n \rightarrow M^k$ such that $\pi: \text{Reg}_K(A) \rightarrow M^k$ is a K -meromorphic finite covering. Applying Theorem 3.1.4 we obtain that $\text{Mer}(\text{Reg}_K(A))$ is a finite extension of $\text{Mer}(M^k) = K$. Since K is algebraically closed, it does not have proper finite extensions, hence $\text{Mer}(\text{Reg}_K(A)) = K$. \square

Towards getting a contradiction we assume that $\mathcal{A}(M)$ is not locally modular.

Remark 4.3.9. The work [8] of Hrushovski and Zilber on Zariski structures implies that if M is not locally modular then it is a finite cover of an algebraic curve over an $\mathcal{A}(M)$ -definable field. However, because we need a more direct relationship to our underlying field K we take a different, more analytic, approach. The method we are using below is based on the techniques of Camapana and Fujiki in the setting of compact complex manifolds, as interpreted in [18] by Pillay and Ziegler and, in the o-minimal setting, in [14].

Let $\mathcal{R}^* \succ \mathcal{R}$ be an $|R|^+$ -saturated elementary extension of \mathcal{R} . We let $R^*, K^*, \mathcal{A}(M^*)$ and $\widetilde{M} \succ \mathcal{A}(M)$ be as in Remark 4.0.5. Notice that since every $m \in M$ is a K -analytic subset of M , we have that $M = \text{acl}(\emptyset)$ in \widetilde{M} .

Since $\mathcal{A}(M)$ is not locally modular and \widetilde{M} is sufficiently saturated, there are $a \in (M^*)^2$, $b_i \in (M^*)^l, i \in \omega$, and an $\mathcal{A}(M)$ -formula $\theta(x, y)$ such that in the structure \widetilde{M} :

- (1) $\text{tp}(ab_i) = \text{tp}(ab_j)$ for all $i, j \in \omega$;
- (2) $\text{tp}(a/b_i)$ is strongly minimal for each $i \in \omega$;
- (3) $\widetilde{M} \models \theta(a, b_i)$ for all $i \in \omega$;
- (4) $\theta(x, b_i) \wedge \theta(x, b_j)$ is finite for all $i \neq j \in \omega$.

Let $C \subset M^{2+l}$ be the irreducible K -analytic subset such that $\text{tp}(ab_0)$ is the generic type of C . By quantifier elimination we can assume that $\theta(x, y)$ defines a Zariski open dense subset W_0 of C . We will denote by π_2 and π_l the projections $\pi_2: M^{2+l} \rightarrow M^2$ and $\pi_l: M^{2+l} \rightarrow M^l$. Let A be the topological closure of $\pi_2(C)$ in M^2 and let B be the topological closure of $\pi_l(C)$ in M^l . By Theorem 4.1.2, A is an irreducible K -analytic subset of M^2 and B is an irreducible K -analytic subset of M^l .

Applying Theorem 4.1.7 we can find a Zariski open dense subset $W \subset W_0 \cap \text{Reg}_K(C)$ such that:

- (a) both π_2 and π_l have constant rank on W ;
- (b) the set $U = \pi_2(W)$ is a K -submanifold of M^2 and Zariski open in A ;
- (c) the $V = \pi_l(W)$ is a K -submanifold of M^l and Zariski open in B .

We will consider W as a K -submanifold of $U \times V$.

We now work in the structure \mathcal{R}^* . For $y \in V^*$ let

$$U_y^* = \{x \in U^* : (x, y) \in W^*\}.$$

Applying Theorem 9.1 from [16]² we obtain that there is a K^* -holomorphic vector bundle $\tau: L \rightarrow U^*$ and K^* -holomorphic map $\mu: W^* \rightarrow \mathbb{P}(L)$ such that the

²Theorem 9.1 and its proof in [16] were formulated in the context of the real field, however the proof was written with the nonstandard setting in mind and carries over to our setting

following diagram is commutative:

$$\begin{array}{ccc} W^* & \xrightarrow{\mu} & \mathbb{P}(L) \\ & \searrow \pi_2 & \swarrow \tau_1 \\ & U^* & \end{array}$$

and $\mu(x, y) = \mu(x, y')$ if and only if $U_y^* = U_{y'}^*$ near x .

Since the type $\text{tp}(ab_0)$ is the generic type of C we obtain that $(ab_i) \in W^*$ for all $i \in \omega$, and, since $U_{b_i}^* \cap U_{b_j}^*$ is finite for $i \neq j$, we have that $\mu(a, b_i) \neq \mu(a, b_j)$ for $i \neq j$.

Let $V_a^* = \{y \in V^* : (a, y) \in W^*\}$. Notice that all b_i are in V_a^* . After fixing a basis for $\tau^{-1}(a)$ we can identify $\tau_1^{-1}(a)$ with a projective space $\mathbb{P}_d(K^*)$. Let $f: V_a^* \rightarrow \mathbb{P}_d(K^*)$ be the map $y \mapsto \mu(a, y)$. Obviously f is K^* -holomorphic, and, since $f(b_i) \neq f(b_j)$ for $i \neq j$, we have that the range of f is infinite.

Since \mathcal{R}^* is an elementary extension of \mathcal{R} , we can find $\alpha \in V$ and K -holomorphic map g from $V_\alpha = \{y \in V : (\alpha, y) \in W\}$ into $\mathbb{P}_d(K)$ whose image is infinite. It is not hard now to derive a contradiction to Claim 4.3.8(2).

This finishes the proof of Theorem 4.3.3.

5. ON THE NON-STANDARD RIEMANN EXISTENCE THEOREM

Recall that the classical Riemann existence theorem states that every compact Riemann surface is bi-holomorphic with an algebraic curve. This theorem fails in the category of K -analytic manifolds. Namely, there is an example of a definably compact K -manifold of K -dimension one that is not K -biholomorphic with an algebraic curve (see [15, Corollary 5.6]).

In fact, Riemann existence theorem can be restated as having two parts.

Theorem 5.0.10 (Riemann existence theorem). *Let M be a compact connected Riemann surface.*

- (1) *There exists a non-constant holomorphic map $\pi: M \rightarrow \mathbb{P}_1(\mathbb{C})$.*
- (2) *For any such π there is a non-singular algebraic curve Γ over \mathbb{C} , a regular map $\tau: \Gamma \rightarrow \mathbb{P}_1(\mathbb{C})$ and a bi-holomorphic map $f: M \rightarrow \Gamma$ such that $\pi = \tau \circ f$.*

The second part of the above theorem can be restated slightly more generally.

Theorem 5.0.11. *Let B be a co-finite subset of \mathbb{C} , X a topological manifold, and $\pi: X \rightarrow B$ a finite covering. Then there exist an affine non-singular algebraic curve Γ over \mathbb{C} , a regular map $\tau: \Gamma \rightarrow B$ and a homeomorphism $f: X \rightarrow \Gamma$ such that $\pi = \tau \circ f$.*

Remark 5.0.12. The above theorem is well-known, but since we could not find it anywhere in the literature, we will outline its proof.

By adding finitely many points, X can be topologically embedded into a compact Riemann surface \hat{X} so that π extends to a holomorphic map $\hat{\pi}: \hat{X} \rightarrow$

$\mathbb{P}_1(\mathbb{C})$ (see [1, page 82] for details). By 5.0.10(2), \hat{X} is bi-holomorphic with an algebraic curve $\hat{\Gamma}$, hence X is homeomorphic to a co-finite subset Γ of $\hat{\Gamma}$.

Obviously Γ is not compact, hence Γ is a proper subset of $\hat{\Gamma}$. Using Riemann-Roch theorem we can find sufficiently many rational functions on $\hat{\Gamma}$ that are regular on Γ , and embed Γ into an affine space.

Our main goal below is to prove an analogue to Theorem 5.0.11 for definable finite coverings of co-finite subsets of K .

5.1. Definable covering maps. We refer to [3] for a definition of a definable space. **We always assume that a definable space is *definably regular*.** We use I to denote the closed interval $[0, 1]$ in R .

Definition 5.1.1. By a definable covering map we mean a definable map $\pi: X \rightarrow B$ between definable spaces that is also a finite covering. A definable n -covering map $\pi: X \rightarrow B$ is called *trivial* if X consists of n definable disjoint sets X_1, \dots, X_n and π maps each X_i homeomorphically onto B .

A *definable isomorphism* of two definable covering maps $\pi: X \rightarrow B$ and $\rho: Y \rightarrow B$ is a definable homeomorphism $f: X \rightarrow Y$ such that $\rho \circ f = \pi$.

Remark 5.1.2. In [5] a definable covering map is defined differently. We will show in Appendix A that in fact both definitions are equivalent. Also, since proofs of some basic properties of definable covering maps are almost identical to analogous statements from Algebraic Topology, we present them in Appendix.

Our goal is to prove the following theorem.

Theorem 5.1.3. *Let $B \subseteq K$ be a co-finite subset, X a definably regular definable space, and $\pi: X \rightarrow B$ a definable covering map. Then there exist a non-singular affine algebraic curve Γ over K , a regular map $\rho: \Gamma \rightarrow B$ and a definable homeomorphism $f: X \rightarrow \Gamma$ such that $\rho \circ f = \pi$.*

To prove the above theorem we will count, up to definable isomorphisms, the number of definable n -covering maps of a co-finite $B \subseteq K$ and then, using classical Riemann existence theorem, show that every such covering can be realized by an affine algebraic curve.

We will need the following claim whose proof is given in Appendix.

Claim 5.1.4. *Let $\pi: X \rightarrow B$ be a definable covering map. If B is definably contractible then π is trivial.*

5.1.1. Definable Paths. Recall that if X is a definable space then a *definable path* in X is a definable continuous function $\sigma: I \rightarrow X$. If $\sigma: I \rightarrow X$ is a definable path then the point $\sigma(0) \in X$ is called *the initial point of σ* , the point $\sigma(1)$ is called *the end point of σ* , and we also say that σ is a definable path from $\sigma(0)$ to $\sigma(1)$.

A *closed path or a loop* is a definable path whose initial point coincides with the end point. If a path σ is a loop with the initial point p then we also say that p is the base point of the loop σ .

Let $\pi: X \rightarrow B$ be a definable covering map and $\sigma: I \rightarrow B$ a definable path. As usual, a definable path $\tau: I \rightarrow X$ is called a *lift* of σ if $\sigma = \pi \circ \tau$.

Claim 5.1.5. *Let $\pi: X \rightarrow B$ be a definable covering map, $\sigma: I \rightarrow B$ a definable path and $\tau_1, \tau_2: I \rightarrow X$ two definable lifts of σ . If $\tau_1(t) = \tau_2(t)$ for some $t \in I$ then $\tau_1(t) = \tau_2(t)$ for all $t \in I$,*

Proof. The set $J = \{t \in I: \tau_1(t) = \tau_2(t)\}$ is both open and closed in I . Since I is definably connected, $J = I$ or empty. \square

The proof of the following claim is presented in Appendix.

Claim 5.1.6 (Path lifting). *Let $\pi: X \rightarrow B$ be a definable finite covering map and σ a definable path in B with the initial point b . Then for every $p \in \pi^{-1}(b)$ there is a unique definable lift τ of σ with the initial point p .*

Following [5] we will denote the end point $\tau(1)$ of the above path τ by $p \star \sigma$.

5.2. K-holomorphic finite cover. In this paper we are interested only in coverings of open subsets of K . Let B be an open definable subset of K and X a K -manifold. A *K-holomorphic covering map* is a K -holomorphic map $\pi: X \rightarrow B$ such that π is also a definable covering map.

Claim 5.2.1. *Let $B \subseteq K$ be a definable open set, X_1, X_2 be K -manifolds, and let $\pi_i: X_i \rightarrow B, i \in \{1, 2\}$, be K -holomorphic covering maps. If $f: X_1 \rightarrow X_2$ is a definable isomorphism of definable covering maps then f is K -holomorphic.*

Proof. Let σ be the isomorphism of the two coverings. Since every definable covering map is locally a bijection, we have locally $\sigma = \pi_1 \pi_2^{-1}$ so σ is K -holomorphic. \square

5.3. Counting definable covering maps of a punctured plane. We follow ideas from [9].

We fix integers n, k and k distinct points $s_1, \dots, s_k \in R^2$. Let $S = \{s_1, \dots, s_k\}$ and $B = R^2 \setminus S$.

We choose a base point $O \in B$ such that for any $s_i \in S$ the straight line through O and s_i does not contain any other point $s_j \in S$.

For every $i = 1, \dots, k$ let r_i be the ray in K with endpoint O that passes through s_i , and let r_i^+ denotes the part of r_i without the segment $[O, s_i)$, i.e.

$$r_i = \{t \cdot \overrightarrow{Os_i}: t \in R^{\geq 0}\} \text{ and } r_i^+ = \{t \cdot \overrightarrow{Os_i}: t \in R^{\geq 0}, t \geq 1\}.$$

Let $U = R^2 \setminus \bigcup_{i=1}^k r_i^+$. Obviously U is definably contractible and is contained in B .

By Claim 5.1.4, if $\pi: X \rightarrow B$ is a definable n -covering map then the restriction of π onto $\pi^{-1}(U)$ is trivial. Thus to describe π we only need to specify how components of $\pi^{-1}(U)$ are glued along $\pi^{-1}(r_i^+ \cap B)$. We fix a small positive $\varepsilon \in R$ such for each s_i the disk D_i of radius ε centered at s_i does not contain neither O nor any other $s_j \in S$.

For each $i = 1, \dots, k$ we pick a loop σ_i in B with base O that starts at O travels towards s_i along the ray r_i until it reaches the boundary of D_i , then it travels once counter-clock-wise along the boundary of D_i , and then comes back to O along r_i .

Let $\pi: X \rightarrow B$ be a definable n -covering map. For $i = 1, \dots, k$ let α_i be the permutation of $\pi^{-1}(O)$ defined as $\alpha_i: p \mapsto p * \sigma_i$. After an enumeration of $\pi^{-1}(O)$, the sequence $\alpha_1, \dots, \alpha_k$ will correspond to a tuple $(g_1, \dots, g_k) \in \text{Sym}(n)^k$. We will denote by $[g_1, \dots, g_k]$ the conjugacy class of (g_1, \dots, g_k) , namely

$$[g_1, \dots, g_k] = \{(hg_1h^{-1}, \dots, hg_kh^{-1}) : h \in \text{Sym}(n)\}$$

Obviously, the tuple (g_1, \dots, g_k) depends on an enumeration of $\pi^{-1}(O)$, but its conjugacy class does not.

We call the conjugacy class $[g_1, \dots, g_k]$ *the monodromy data of the covering* $\pi: X \rightarrow B$ (with the base point O).




Theorem 5.3.1. *For every tuple $(g_1, \dots, g_k) \in \text{Sym}(n)^k$, up-to a definable isomorphism, there is a unique definable n -covering $\pi: X \rightarrow B$ with the monodromy data $[g_1, \dots, g_k]$.*

Proof. For existence we will actually show that every monodromy data can be realized by a semi-algebraic covering.

We choose pairwise disjoint tubular neighborhoods V_i of r_i^+ in B . We denote by V_i^l and V_i^r the left and the right connected components of $V_i \cap U$.

Let $X_1 = U \times \{1, \dots, n\}$, $X_2 = \bigcup_{i=1}^k V_i \times \{1, \dots, n\}$, and $A = \bigcup_{i=1}^k (V_i^l \cup V_i^r) \times \{1, \dots, n\}$. We consider A as a subset of X_1 . Let $f: A \rightarrow X_2$ be the map defined as

$$f((p, j)) = \begin{cases} (p, j) & \text{if } p \in V_i^r \\ (p, g_i(j)) & \text{if } p \in V_i^l. \end{cases}$$



Let $X = X_1 \coprod_f X_2$ be the space obtained by attaching X_1 to X_2 via f (see [3, Chapter 10, Section 2.9] for a precise definition), and $\pi: X \rightarrow B$ be the map $(p, j) \mapsto p$. It is easy to see that π is a definable n -covering map with the monodromy data $[g_1, \dots, g_k]$.

For a uniqueness, let $\tau: Y \rightarrow B$ be a definable n -covering with the monodromy data $[g_1, \dots, g_k]$.

We can choose an enumeration of $\pi^{-1}(O)$ so that, with respect to this enumeration, $\alpha_i = g_i$ for $i = 1, \dots, k$. It is not hard to see that the covering $\tau: Y \rightarrow B$ is definably isomorphic to the semialgebraic covering $\pi: X \rightarrow B$ constructed above. \square

Corollary 5.3.2. *For any integers $k, n \in \mathbb{N}^{>0}$ there is a number $N(k, n)$ such that for any set $S \subseteq \mathbb{R}^2$ with $|S| = k$ up to a definable isomorphism there are exactly $N(k, n)$ definable n -covering maps $\pi: X \rightarrow \mathbb{R}^2 \setminus S$. This number $N(k, n)$ depends on n and k only and does not depend on neither the structure \mathcal{R} , nor the set S .*

Proof. It follows from the previous discussion that the number $N(k, n)$ is equal to the number of conjugacy classes in $\text{Sym}(n)^k$. \square

5.3.1. *The classical case.* In the case, when the field R is the field of the real number, it is not hard to see that two definable n -coverings of a co-finite subset of \mathbb{R}^2 are definably isomorphic if and only if they are isomorphic topologically.

Indeed, let $\pi_i: X_i \rightarrow B, i = 1, 2$, be definable n -coverings of a co-finite subset B of \mathbb{R}^2 . If $f: X_1 \rightarrow X_2$ is a homeomorphism, then, since in this case Claim 5.1.4 and Claim 5.1.5 hold without definability assumptions (e.g. see [7]), π_1 and π_2 have the same monodromy data, hence definably isomorphic.

5.4. **Proof of Theorem 5.1.3.** We need the following claim.

Claim 5.4.1. *Let $n \in \mathbb{N}$, $B \subseteq K$ a co-finite subset, Γ a non-singular affine algebraic curve over K , and π a regular function on Γ . Then $\pi: \Gamma \rightarrow B$ is a definable n -covering (with respect to the induced R -topology) if and only if π is surjective and the size of $\pi^{-1}(y)$ is exactly n for every $y \in B$.*

Proof. By Claim 3.1.1, we need only to check that π is a local homeomorphism. Since Γ is non. Let $\Gamma_1, \dots, \Gamma_l$ be the irreducible components of Γ , and let $\pi_i = \pi|_{\Gamma_i}$.

For $i = 1, \dots, l$ and $y \in B$ we have that $|\pi^{-1}(y)| \leq \deg(\pi_i)$ with equality for almost all $y \in B$. Thus the sum of degrees of π_i is n , and $|\pi_i^{-1}(y)| = \deg(\pi_i)$ for all $y \in B$.

Thus every π_i is an unramified finite map, its differential does not vanish anywhere on Γ_i , (see [19, Chapter II, Section 6.3, Theorem 5]), hence it is a local diffeomorphism \square

The above claim allows us to give a purely algebraic definition of an algebraic n -covering. Namely, let $B \subseteq K$ be a co-finite subset. A pair $\langle \Gamma, \pi \rangle$ is an *algebraic n -covering of B* if Γ is a non-singular affine algebraic curve over K , π is a regular function on Γ , B is the range of π , and $|\pi^{-1}(y)| = n$ for all $y \in B$. Two algebraic n -coverings $\langle \Gamma, \pi \rangle$ and $\langle \Gamma', \pi' \rangle$ are *isomorphic* if there is a regular map $f: \Gamma \rightarrow \Gamma'$ such that $\pi = \pi' \circ f$. If L is a subfield of K and both Γ and π are defined over L then we say that $\langle \Gamma, \pi \rangle$ is a covering over L .

Theorem 5.4.2. *Let $\langle \Gamma, \pi \rangle$ and $\langle \Gamma', \pi' \rangle$ be two algebraic n -coverings of a co-finite set $B \subset K$, $|K \setminus B| = k$. Then they are isomorphic as algebraic covers if and only if they are isomorphic as definable covering maps.*

In particular, there are at most $N(k, n)$ (see 5.3.2) algebraic n -coverings of B , independently of K and of B .

Proof. Clearly, every algebraic isomorphism of the two covers is also definable, so we need to prove the converse.

Assume that $\sigma: \Gamma \rightarrow \Gamma'$ is a definable isomorphism. Because $\Gamma \subset K^n$ and $\Gamma' \subset K^m$ are non-singular curves, they are K -manifolds. By Claim 5.2.1, the map σ is K -holomorphic, hence by our version of Chow's Theorem (Theorem 2.3.2), it is an algebraic map. The final clause follows from 5.3.2. \square

Thus, in order to prove Theorem 5.1.3 it is sufficient to prove the following purely algebraic result.

Theorem 5.4.3. *Let L be an algebraically closed field characteristic zero, $B \subseteq L$ a co-finite subset, $k = |L \setminus B|$. Then there are exactly $N(k, n)$ non-isomorphic algebraic n -coverings of B defined over the field L .*

Proof. We fix integers n and k .

For an algebraically closed characteristic zero field L and a co-finite subset $B \subseteq L$ with $|L \setminus B| = k$ we will denote by $N'(L, B)$ the number of non-isomorphic algebraic n -coverings of B defined over the field L . We need to prove that $N'(L, B) = N(k, n)$.

Remark 5.4.4. Using the equivalence of categories of algebraic finite coverings and ramified extensions of $L(X)$, this result can be derived from the main result of [4]. Here we present an alternative proof.

It follows from Theorem 5.4.2 that $N'(L, B) \leq N(k, n)$, and we only need to show that $N'(L, B) \geq N(k, n)$.

First notice that the above inequality holds in the case $L = \mathbb{C}$. Indeed, by Corollary 5.3.2 (applied to the o-minimal real field), and discussion in Section 5.3.1, there are $N(k, n)$ -many different topological n -coverings of $B \subseteq \mathbb{C}$. By Theorem 5.0.11 (i.e., by the classical Riemann existence Theorem), every such covering is topologically isomorphic to an n -algebraic covering of B . We therefore have $N(k, n)$ -many algebraic n -coverings of B that are non-isomorphic as algebraic curves (since they are not even topologically isomorphic).

For the purpose of getting a contradiction we assume that for some algebraically closed characteristic zero field L and $B \subseteq L$ with $|L \setminus B| = k$ such that $N'(L, B) < N(k, n)$. Let $N' = N'(L, B)$.

Let $\langle \Gamma_1, \pi_1 \rangle, \dots, \langle \Gamma_{N'}, \pi_{N'} \rangle$ be non-isomorphic algebraic coverings of B defined over L . Thus if $\langle \Gamma, \pi \rangle$ is an algebraic covering of B defined over L then it is isomorphic to one of the $\langle \Gamma_i, \pi_i \rangle$'s. Let $S = L \setminus B$. By restricting ourselves to a subfield of L over which S and Γ_i, π_i are all defined we may assume that L is a countable field.

Since the field \mathbb{C} is saturated, we can embed L into \mathbb{C} and assume that L is a countable subfield of \mathbb{C} . Let $\tilde{\Gamma}_i$ and $\tilde{\pi}_i$ be \mathbb{C} -points of Γ_i and π_i respectively; and let $\tilde{B} = \mathbb{C} \setminus S$

Since $N' < N(k, n)$, it follows from the case $L = \mathbb{C}$ that there exists an algebraic n -covering $\langle \mathcal{C}, \rho \rangle$ of \tilde{B} that is not isomorphic over \mathbb{C} to any of the $\langle \tilde{\Gamma}_i, \tilde{\pi}_i \rangle$'s. As we will show, this cover is isomorphic to a cover over L and this leads to a contradiction.

We choose polynomials $p_1(\bar{x}, \bar{u}), \dots, p_m(\bar{x}, \bar{u}), q(\bar{x}, \bar{u})$ over \mathbb{Z} , and $\bar{d} \in \mathbb{C}^l$ such that $\mathcal{C} = \{\bar{z} \in \mathbb{C}^s : p_1(\bar{z}, \bar{d}) = \dots = p_m(\bar{z}, \bar{d}) = 0\}$ and $\rho = q(\bar{x}, \bar{d})$.

For $\bar{a} \in \mathbb{C}^l$ let $\mathcal{C}_{\bar{a}} = \{\bar{z} \in \mathbb{C}^s : p_1(\bar{z}, \bar{a}) = \dots = p_m(\bar{z}, \bar{a}) = 0\}$ and $\rho_{\bar{a}} = q(\bar{x}, \bar{a})$. It is easy to see that the set

$$A = \{\bar{a} \in \mathbb{C}^l : \langle \mathcal{C}_{\bar{a}}, \rho_{\bar{a}} \rangle \text{ is an algebraic } n\text{-covering of } \tilde{B}\}$$

is definable (in the language of rings) over the field L .

Consider the equivalence relation \mathcal{E} on A given as

$$\mathcal{E}(\bar{a}, \bar{a}') \text{ iff the algebraic coverings } \langle \mathcal{C}_{\bar{a}}, \rho_{\bar{a}} \rangle \text{ and } \langle \mathcal{C}_{\bar{a}'}, \rho_{\bar{a}'} \rangle \text{ are isomorphic.}$$

It is not hard to see that $\mathcal{E}(\bar{x}, \bar{x}')$ is \forall -definable over L , i.e. it is a countable union of sets definable over L .

By 5.4.2, \mathcal{E} has at most $N(k, n)$ -many classes. Since \mathbb{C} is \aleph_1 -saturated, every \forall -definable equivalence relation on A with finitely many classes must be definable (every \mathcal{E} -class is \forall -definable with \forall -definable complement, hence definable). Since L is elementary substructure of \mathbb{C} , and \mathcal{E} is a definable (over L) equivalence relation with finitely many classes, every \mathcal{E} -class must have a representative in L . Thus there is $\bar{d}' \in L^l$ with $\mathcal{E}(\bar{d}, \bar{d}')$. The algebraic covering $\langle \mathcal{C}_{\bar{d}'}, \rho_{\bar{d}'} \rangle$ is defined over L and is not isomorphic to any of the $\langle \Gamma_i, \pi_i \rangle$'s. A contradiction. \square

This ends the proof of Theorem 5.1.3.

6. PROOF OF THE MAIN THEOREM

In this section we will prove the main theorem stated in the introduction.

Theorem 6.0.5. *Let M be a strongly minimal mild manifold. Then either the structure $\mathcal{A}(M)$ is locally modular or M is K -biholomorphic to a non-singular algebraic curve over K .*

Proof. Assume $\mathcal{A}(M)$ is not locally modular. Then, by Theorem 4.3.3, there is a non-constant K -meromorphic $\pi: M \rightarrow K$. By Claim 4.3.6, there is a finite subset $Z \subseteq M$ and a finite subset $S \subseteq K$ such that $\pi: M \setminus Z \rightarrow K \setminus S$ is a K -holomorphic finite covering. Using Theorem 5.1.3 we can find a non-singular affine curve Γ and a K -biholomorphism $g: M \setminus Z \rightarrow \Gamma$.

Since M is definably connected and Z is finite, $M \setminus Z$ is definably connected, hence Γ is irreducible.

Taking a projectivization of Γ and then, if needed, a normalization, we can find a non-singular projective curve \mathcal{C} and an injective regular map $h: \Gamma \rightarrow \mathcal{C}$ whose range is co-finite in \mathcal{C} . Composing g with h we obtain a K -holomorphic injective map $f: M \setminus Z \rightarrow \mathcal{C}$ whose range is co-finite in \mathcal{C} .

We first claim that f can be extended to a K -holomorphic map $F: M \rightarrow \mathcal{C}$.

Notice that $\dim_K M = 1$ and \mathcal{C} is definably compact (as a definable closed subset of a projective space). Let $G_f \subseteq M \times \mathcal{C}$ be the graph of f and A its topological closure. By [14, Corollary 9.1], A is a K -analytic subset of $M \times \mathcal{C}$. Since the o-minimal dimension of G_f is 2 we have that o-minimal dimension of A is also 2.

Since all proper k -analytic subsets of \mathcal{C} are finite, for every $m \in M$ the set $A_m = A \cap (\{m\} \times \mathcal{C})$ is either finite or co-finite. Because the o-minimal dimension of $A \setminus G_f$ is at most one, for every $z \in Z$, the set A_z must be finite.

Pick $z \in Z$. Let $U \subseteq M$ be a definable open subset of M definably K -biholomorphic to an open disk $D \subseteq K$ with $U \cap Z = \{z\}$. By [12, Theorem 2.37], it is sufficient to show that f can be extended continuously to z . Since \mathcal{C} is definably compact, $f(\gamma(t))$ has a limit in \mathcal{C} for every definable curve $\gamma(t)$ in M approaching z , hence A_z is not empty. Since $U \setminus \{z\}$ is definably connected, A_z is also definably connected (see [12, Fact 2.2]), hence it contains exactly one point. Using the definable compactness of \mathcal{C} it is not hard to see that A is the graph of a continuous function $F: M \rightarrow \mathcal{C}$ extending f .

We claim that F is injective. Indeed if $F(a) = F(b)$ for some $a \neq b \in M$, then, by Open Mapping Theorem [12, Corollary 2.34], for every $a' \in M$ close enough to a there is $b' \in M$ close enough to b such that $F(a') = F(b')$. This contradicts to the injectivity of f .

Let \mathcal{C}' be the range of F . Then \mathcal{C}' is a quasi-projective smooth curve K -biholomorphic to M . □

Example 6.0.6. As was shown in [15], there are one-dimensional definably compact K -groups (these are K -manifolds with K -holomorphic group operations) which are not isomorphic as K -manifolds to any algebraic curve. It follows from Theorem 6.0.5 that the every such K -group, with all its induced K -analytic structure, is locally modular.

Proof of Claim 5.1.6. Let $\pi: X \rightarrow B$ be a definable n -covering map, $\sigma: I \rightarrow B$ a definable path with the initial point b , and $p \in \pi^{-1}(b)$.

Uniqueness follows from Claim 5.1.5, and we only need to show existence of a lift $\tau: I \rightarrow X$ of σ with the initial point p .

Using definable choice we can find n definable functions $g_1, \dots, g_n: B \rightarrow X$ such that $\pi^{-1}(z) = \{g_1(z), \dots, g_n(z)\}$ for all $z \in B$. Using basic properties of o-minimal structures we can find $0 = t_0 < t_1 < \dots < t_k = 1$ such that for every $j = 1, \dots, n$ and $i = 1, \dots, k$ the function $h_j(t) = g_j(\sigma(t))$ is continuous on (t_{i-1}, t_i) . By induction on $i = 0, \dots, k$ we construct a definable continuous function $\tau_i: [0, t_i] \rightarrow X$ such that $\tau_i(0) = p$ and $\pi \circ \tau_i = \sigma$ on $[0, t_i]$. We set $\tau_0(0) = p$. Assume we have τ_i . Let V be a definable open neighbourhood of $\tau_i(t_i)$ such that π maps V homeomorphically onto an open neighborhood U of $\sigma(t_i)$. It is not hard to see that there is unique $j \in \{1, \dots, n\}$ such that $g_j(\sigma(t)) \in V$ for all $t \in (t_i, t_i + \varepsilon)$ for some sufficiently small $\varepsilon > 0$. Let

$$\tau'_{i+1} = \begin{cases} \tau_i(t) & \text{if } t \in [0, t_i) \\ g_j(\sigma(t)) & \text{if } t \in (t_i, t_{i+1}). \end{cases}$$

Obviously τ'_{i+1} is continuous on $[0, t_{i+1})$ and $\pi \circ \tau'_{i+1} = \sigma$ on $[0, t_{i+1})$. We now set $\tau_{i+1} = \tau'_{i+1}$ on $[0, t_{i+1})$ and $\tau_{i+1}(t_{i+1}) = q$, where q is a unique point in $\pi^{-1}(t_{i+1})$ with $\lim_{t \rightarrow t_{i+1}^-} \tau'_{i+1}(t) = q$. \square

Remark A.0.7. It is not hard to see that the above prove can be generalized to a definable family of definable paths. Namely if $\sigma_a, a \in A$, is a definable family of definable paths in B , and $f: A \rightarrow X$ is a definable function with $\pi(f(a)) = \sigma_a(0)$ then there is a definable family of paths $\tau_a: I \rightarrow X, a \in A$, such that each τ_a is a lift of σ_a with $\tau_a(0) = f(a)$.

Proof of Claim 5.1.4. Let $\pi: X \rightarrow B$ be a definable n -covering map with B definably contractible, i.e. there is $b \in B$ and a definable continuous $F: B \times I \rightarrow B$ such that

$$F(z, 1) = z \text{ and } F(z, 0) = b \text{ for all } z \in B.$$

We need to show that π is a trivial cover. Recall that a definable section of π is a definable continuous map $\rho: B \rightarrow X$ such that $\pi \circ \rho = \text{id}_B$. If ρ is a definable section of π then π restricted to $X \setminus \rho(B)$ is a definable $n - 1$ -covering map. Therefore, applying an induction on n , it is sufficient to show that there is a definable section of π .

We fix $p \in \pi^{-1}(b)$. For $z \in B$ let σ_z be the definable path $t \mapsto F(z, t)$. Thus we have a definable family $\sigma_z, z \in B$, of definable paths in B all with the same initial point b . By the previous remark, there is a definable family $\tau_z, z \in B$, of definable paths in X all with the initial point p such that $\pi(\tau_z(t)) = F(z, t)$ for all $z \in B, t \in I$. Let $\widehat{F}(z, t): B \times I \rightarrow X$ be the map $F(z, t) = \tau_z(t)$. We have

$$\pi \circ \widehat{F} = F, \quad \widehat{F}(z, 0) = p \text{ for all } z \in B,$$

and for each $z \in B$ the map $t \mapsto \widehat{F}(z, t)$ is continuous on I .

Let $\rho: B \rightarrow X$ be the map $z \mapsto \widehat{F}(z, 1)$. Clearly $\pi \circ \rho = \text{id}_B$, and it remains to show that ρ is continuous.

Let $z_0 \in B$ and $x_0 = \rho(z_0)$. We will show that ρ is continuous at z_0 .

Let $\sigma = \sigma_{z_0}$, and $\tau = \tau_{z_0}$. Thus σ is a path in B from b to z_0 , and τ is a path in X from p to x_0 .

For a definable open subset $U \subseteq B$ we say that π is trivial over U if $\pi^{-1}(U)$ consists of n definably connected components.

Since B is definably normal space, by [3, Chapter 10, Theorem 1.8], it is affine and we may assume that $B \subseteq R^m$. Let $d(\cdot, \cdot): R^m \times R^m \rightarrow R$ be the distance function on R^m .

For $z \in B$ and $\varepsilon \in R$ let $\mathcal{O}_\varepsilon(z) = \{z' \in B: d(z, z') < \varepsilon\}$.

For $t \in I$ let $h(t) = \sup\{\varepsilon \in R: \pi \text{ is trivial over } \mathcal{O}_\varepsilon(\sigma(t))\}$. Clearly, h is a definable function from I to $R^{>0} \cup \{+\infty\}$, and it is locally bounded from below, i.e for every $t \in I$ there is $\varepsilon > 0$ such that $h(t') > \varepsilon$ for all $t' \in I$ sufficiently close to t . Since I is definably compact, an easy application of curve selection yields that there is positive $\varepsilon_0 \in R$ such that $h(t) > \varepsilon_0$ for all $t \in I$.

Since F is continuous, I definably compact, and $F(z_0, 1) = z_0$, it is not hard to see that there is positive $\delta_0 \in R$ such that for all $z \in \mathcal{O}_{\delta_0}(z_0)$ we have $d(F(z, t), F(z_0, t)) < \varepsilon_0$ for all $t \in I$. Increasing δ_0 if needed, we may assume $\delta_0 \leq \varepsilon_0$.

Let V be the definably connected component of $\pi^{-1}(\mathcal{O}_{\delta_0}(z_0))$ containing x_0 . We have that π maps V homeomorphically onto $\mathcal{O}_{\delta_0}(z_0)$, and to show that ρ is continuous at z_0 , it is sufficient to show that $\rho(z) \in V$ for all $z \in \mathcal{O}_{\delta_0}(z_0)$.

Let $z' \in \mathcal{O}_{\delta_0}(z_0)$. It is not hard to see that the set

$$A = \{t \in I: \tau_{z'}(t) \text{ and } \tau(t) \text{ are in the same} \\ \text{definably connected component of } \pi^{-1}(\mathcal{O}_{\varepsilon_0}(\sigma(t)))\}$$

is both open and close. Since I is definably connected and $0 \in A$ we have $1 \in A$, hence $1 \in A$ and $\rho(z') = \tau_{z'}(1) \in V$. □

The following corollary shows that our definition of covering maps is equivalent to the definition in [5]

Corollary A.0.8. *Let $\pi: X \rightarrow B$ be a definable n -covering map. Then B can be covered by finitely many open definable sets $V_i, i = 1, \dots, k$ such that for each $\pi^{-1}(V_i)$ consists of n pair-wise disjoint definable sets U_i^1, \dots, U_i^n and π maps each U_i^j homeomorphically onto V_i .*

Proof. It is sufficient to show that B can be covered by finitely many open definable definably contractible sets V_i . We can take V_i to be star neighborhoods of vertexes in the barycentric subdivision of any triangulation of X . □

REFERENCES

- [1] Jean-Benoît Bost, *Introduction to compact Riemann surfaces, Jacobians, and abelian varieties*, From number theory to physics (Les Houches, 1989), Springer, Berlin, 1992, pp. 64–211.
- [2] Claude Chevalley, *Introduction to the theory of algebraic functions of one variable*, Mathematical Surveys, No. VI, American Mathematical Society, Providence, R.I., 1963.
- [3] Lou van den Dries, *Tame topology and o-minimal structures*, London Mathematical Society Lecture Note Series, vol. 248, Cambridge University Press, Cambridge, 1998.
- [4] Lou van den Dries and Paulo Ribenboim, *Application de la théorie des modèles aux groupes de Galois de corps de fonctions*, C. R. Acad. Sci. Paris Sér. A-B **288** (1979), no. 17, A789–A792 (French, with English summary).
- [5] Mário J. Edmundo and Margarita Otero, *Definably compact abelian groups*, J. Math. Log. **4** (2004), no. 2, 163–180.
- [6] Robin Hartshorne, *Algebraic geometry*, Springer-Verlag, New York, 1977. Graduate Texts in Mathematics, No. 52.
- [7] Allen Hatcher, *Algebraic topology*, Cambridge University Press, Cambridge, 2002.
- [8] Ehud Hrushovski and Boris Zilber, *Zariski geometries*, J. Amer. Math. Soc. **9** (1996), no. 1, 1–56.
- [9] A. Hurwitz, *Ueber Riemann'sche Flächen mit gegebenen Verzweigungspunkten*, Math. Ann. **39** (1891), no. 1, 1–60 (German).
- [10] Rahim Moosa, *A nonstandard Riemann existence theorem*, Trans. Amer. Math. Soc. **356** (2004), no. 5, 1781–1797 (electronic).
- [11] R. Moosa and S. Starchenko, *K-analytic versus ccm-analytic sets in nonstandard compact manifolds*, Fund. Math.
- [12] Ya'acov Peterzil and Sergei Starchenko, *Expansions of algebraically closed fields in o-minimal structures*, Selecta Math. (N.S.) **7** (2001), no. 3, 409–445.
- [13] ———, *Expansions of algebraically closed fields. II. Functions of several variables*, J. Math. Log., to appear.
- [14] ———, *Complex analytic geometry in a nonstandard setting*, Model Theory with applications to algebra and analysis, Vol. I, Cambridge University Press, Cambridge, 2008.
- [15] ———, *Uniform definability of the Weierstrass \wp functions and generalized tori of dimension one*, Selecta Math. (N.S.) **10** (2004), no. 4, 525–550.
- [16] ———, *Complex analytic geometry and analytic-geometric categories*, J. Reine Angew. Math., to appear.
- [17] Anand Pillay, *Some model theory of compact complex spaces*, geometry (Ghent, 1999), Contemp. Math., vol. 270, Amer. Math. Soc., Providence, RI, 2000, pp. 323–338.
- [18] Anand Pillay and Martin Ziegler, *Jet spaces of varieties over differential and difference fields*, Selecta Math. (N.S.) **9** (2003), no. 4, 579–599.

- [19] Igor R. Shafarevich, *Basic algebraic geometry. 1*, 2nd ed., Springer-Verlag, Berlin, 1994. Varieties in projective space; Translated from the 1988 Russian edition and with notes by Miles Reid.
- [20] B. Zil'ber, *Model theory and algebraic geometry*, Proceedings of the 10th Easter Conference on Model Theory, Seminarberichte, vol. 93, Humboldt Univ. Berlin, 1993, pp. 202–222.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAIFA, HAIFA, ISRAEL
E-mail address: `kobi@math.haifa.ac.il`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, NOTRE DAME, IN 46556
E-mail address: `starchenko.1@nd.edu`