

# A NOTE ON O-MINIMAL FLOWS AND THE AX-LINDEMANN-WEIERSTRASS THEOREM FOR ABELIAN VARIETIES OVER $\mathbb{C}$

YA'ACOV PETERZIL AND SERGEI STARCHENKO

ABSTRACT. In this short note we present an elementary proof of Theorem 1.2 from [9], and also the Ax-Lindemann-Weierstrass theorem for abelian and semi-abelian varieties. The proof uses ideas of Pila, Ullmo, Yafaev, Zannier (see e.g. [7]) and is based on basic properties of sets definable in o-minimal structures. It does not use the Pila-Wilkie counting theorem.

## 1. INTRODUCTION

In their article [7], Pila and Zannier proposed a new method to tackle problems in Arithmetic geometry, a method which makes use from model theory, and in particular the theory of o-minimal structures. Their goal was to produce a new proof for the Manin-Mumford conjecture and it went roughly as follows: Consider the transcendental uniformizing map  $\pi : \mathbb{C}^n \rightarrow A$  for an  $n$ -dimensional abelian variety  $A$ . Given an algebraic variety  $V \subseteq A$ , with “many” torsion points, consider its pre-image  $\tilde{V} = \pi^{-1}(V)$ . The analytic periodic set  $\tilde{V}$ , when restricted to a fundamental domain  $F \subseteq \mathbb{C}^n$ , is definable in the o-minimal structure  $\mathbb{R}_{an}$ . At the heart of the proposed method was a theorem by Pila and Wilkie, [6], used to conclude that  $\tilde{V}$  contains an algebraic variety  $X$ . In the last step of the proof one shows that  $X$  is contained in a coset of a  $\mathbb{C}$ -linear subspace  $L$  of  $\mathbb{C}^n$ , with  $L \subseteq \tilde{V}$ . Finally, the Zariski closure of  $\pi(L)$  is a coset of an abelian subvariety of  $V$ , which is the goal of the theorem. Because of various equivalent formulations this last step of the argument became known as the “Ax-Lindemann-Weierstrass” statement for abelian varieties, which we call here ALW.

Following the seminal paper of Pila, [4] on the Andre-Oort Conjecture for  $\mathbb{C}^n$  it became clear that the Pila-Zannier method was very

---

*Date:* November 14, 2016.

The authors thank the US-Israel Binational Science Foundation for its support.

The second author was partially supported by the NSF Research Grant DMS-150067.

effective in attacking other problems in arithmetic geometry. Each such problem was broken-up into various parts and the ALW was isolated as a separate statement. Somewhat surprisingly, despite the fact that ALW does not seem to have a clear arithmetic content, Pila found an ingenious way to apply the Pila-Wilkie theorem again in order to prove it in the setting of the Andre-Oort conjecture (this is sometimes called “the hyperbolic ALW”). The method of Pila was applied extensively since then to settle several variants of ALW ([3], [5], [2]).

Our goal in this note is to show that at least in the setting of semi-abelian varieties, where the uniformizing map is actually a group homomorphism, the use of Pila-Wilkie is unnecessary and the proof of ALW and most of its variants becomes quite elementary. We believe that this simpler approach can clarify the picture substantially and eventually yield new results as well.

**1.1. Geometric restatements of ALW for semi-abelian varieties.** The following theorem follows from a more general theorem of Ax (see [1, Theorem3]) and often is called the full Ax-Lindemann-Weierstrass Theorem. The original proof of Ax used algebraic differential methods.

**Theorem 1.1** (Full ALW). *Let  $G$  be a connected semi-abelian variety over  $\mathbb{C}$ ,  $\mathbf{T}_G$  the Lie algebra of  $G$  and  $\exp_G: \mathbf{T}_G \rightarrow G$  the exponential map. Let  $W \subseteq \mathbf{T}_G$  be an irreducible algebraic variety and  $\xi_1, \dots, \xi_k \in \mathbb{C}(W)$ . If  $\xi_1, \dots, \xi_k$  are  $\mathbb{Q}$ -linearly independent over  $\mathbb{C}$  then  $\exp_G(\xi_1), \dots, \exp_G(\xi_k)$  are algebraically independent over  $\mathbb{C}(W)$ .*

If in above theorem we change the conclusion to “algebraically independent over  $\mathbb{C}$ ” then we get a weaker statement that often is called Ax-Lindemann-Weierstrass theorem (ALW theorem for short).

It is not hard to see that both full ALW and ALW theorems can be interpreted geometrically (see e.g. [8] for more details).

**Theorem 1.2** (ALW, Geometric Version). *Let  $G$  be a connected semi-abelian variety over  $\mathbb{C}$ ,  $\mathbf{T}_G$  the Lie algebra of  $G$  and  $\exp_G: \mathbf{T}_G \rightarrow G$  the exponential map.*

*Let  $X \subseteq \mathbf{T}_G$  be an irreducible algebraic variety and  $Z \subseteq G$  the Zariski closure of  $\exp_G(X)$ . Then  $Z$  is a translate of an algebraic subgroup of  $G$ .*

We can also restate full ALW.

**Theorem 1.3** (Full ALW, Geometric Version). *Let  $G$  be a connected semi-abelian variety over  $\mathbb{C}$ ,  $\mathbf{T}_G$  the Lie algebra of  $G$ ,  $\exp_G: \mathbf{T}_G \rightarrow$*

$G$  the exponential map, and  $\pi: \mathbf{T}_G \rightarrow \mathbf{T}_G \times G$  be the map  $\pi(z) = (z, \exp_G(z))$ .

Let  $X \subseteq \mathbf{T}_G$  be an irreducible algebraic variety and let  $Z \subseteq \mathbf{T}_G \times G$  be the Zariski closure of  $\pi(X)$ . Then  $Z = X \times B$ , where  $B$  is a translate of an algebraic subgroup of  $G$ .

## 2. PRELIMINARIES

We work in an o-minimal expansion  $\mathcal{R}$  of the real field  $\mathbb{R}$ , and by definable we always mean  $\mathcal{R}$ -definable (with parameters). The only property of o-minimal structures that we need is that every definable discrete subset is finite.

If  $V$  is a finite dimensional vector space over  $\mathbb{R}$  and  $X$  a subset of  $V$  then, as usual, we say that  $X$  is *definable* if it becomes definable after fixing a basis for  $V$  and identifying  $V$  with  $\mathbb{R}^n$ . Clearly this notion does not depend on a choice of basis.

Let  $\pi: V \rightarrow G$  be a group homomorphism, where  $V$  is a finite dimensional vector space over  $\mathbb{R}$  and  $G$  a connected commutative algebraic group over  $\mathbb{C}$ . We denote the group operation of  $G$  by  $\cdot$ .

Let  $\Lambda = \pi^{-1}(e)$ . We say that a subset  $F \subseteq V$  is a *large domain* for  $\pi$  if  $F$  is a connected open subset of  $V$  with  $V = F + \Lambda$ . If in addition the restriction of  $\pi$  to  $F$  is definable then we say that  $F$  is a *definable large domain* for  $\pi$ .

*Remark 2.1.* In the above setting if  $\pi$  is real analytic and  $\Lambda$  is a lattice in  $V$  then  $V/\Lambda$  is compact and there is a relatively compact large domain for  $\pi$  definable in  $\mathbb{R}_{\text{an}}$ .

## 3. KEY OBSERVATIONS

In this section we fix a finite dimensional  $\mathbb{C}$ -vector space  $V$ , a connected commutative algebraic group  $G$  over  $\mathbb{C}$  and  $\pi: V \rightarrow G$  a complex analytic group homomorphism. We assume that  $\Lambda = \pi^{-1}(e)$  is a discrete subgroup of  $V$  and that  $\pi$  has a definable large domain  $F$ .

Let  $X$  be a definable connected real analytic submanifold of  $V$  and let  $Z$  be the Zariski closure of  $\pi(X)$  in  $G$ .

Let  $\tilde{Z} = \pi^{-1}(Z)$  and  $\tilde{Z}_F = \tilde{Z} \cap F$ . The set  $\tilde{Z}$  is a complex analytic  $\Lambda$ -invariant subset of  $V$  and  $\tilde{Z}_F$  is a definable subset of  $F$ .

Let

$$(3.1) \quad \Sigma_F(X) = \{v \in V: v + X \cap F \neq \emptyset \text{ and } v + X \cap F \subseteq \tilde{Z}_F\}.$$

Clearly  $\Sigma_F(X)$  is a definable subset of  $V$ .

The following is an elementary observation.

**Observation 3.1.** (1) If  $\lambda \in \Lambda$  and  $\lambda + F \cap X \neq \emptyset$  then  $-\lambda \in \Sigma_F(X)$ .

In particular  $X \subseteq F - (\Sigma_F(X) \cap \Lambda)$ .

(2) If  $v$  is in  $\Sigma_F(X)$  then  $v + X \subseteq \tilde{Z}$ .

As a consequence we have the following claim.

**Claim 3.2.**  $\pi(\Sigma_F(X)) \subseteq \text{Stab}_G(Z) = \{g \in G: g \cdot Z = Z\}$ .

*Proof.* If  $v$  is in  $\Sigma_F(X)$  then by Observation 3.1(2) we have  $X \subseteq \tilde{Z} - v$ , and hence  $\pi(X) \subseteq \pi(v)^{-1} \cdot \pi(\tilde{Z}) = \pi(v)^{-1} \cdot Z$ . Since  $Z$  is the Zariski closure of  $\pi(X)$  and  $\pi(v)^{-1} \cdot Z$  is a subvariety of  $G$  we have  $Z \subseteq \pi(v)^{-1} \cdot Z$ , hence  $\pi(v)$  is in the stabilizer of  $Z$ .  $\square$

*Remark 3.3.* Both Observation 3.1 and Claim 3.2 hold for a complex irreducible algebraic subvariety  $X$  of  $V$ . It can be done either by a direct argument or replacing  $X$  with the set  $X_{\text{reg}}$  of smooth points on  $X$  and using the fact that  $X_{\text{reg}}$  is a connected complex submanifold of  $V$  that is dense in  $X$ .

We deduce a slight generalization of Theorem 1.2 from [9].

**Proposition 3.4.** Let  $\pi: V \rightarrow G$  be a complex analytic group homomorphism from a finite dimensional  $\mathbb{C}$ -vector space  $V$  to a connected commutative algebraic group  $G$  over  $\mathbb{C}$ . Let  $\Lambda = \pi^{-1}(e)$ . Assume  $\pi$  has a large definable domain  $F$ .

Let  $X \subseteq V$  be a definable connected real analytic submanifold (or an irreducible complex algebraic subvariety) and  $Z \subseteq G$  the Zariski closure of  $\pi(X)$  in  $G$ .

If  $X$  is not covered by finitely many  $\Lambda$ -translate of  $F$  then  $\text{Stab}_G(Z)$  is infinite.

*Proof.* If  $X$  is not covered by finitely many  $\Lambda$ -translate of  $F$ , then by Observation 3.1(1) the set  $\Sigma_F(X)$  is infinite. Since it is also definable,  $\pi(\Sigma_F(X))$  must be also infinite (otherwise  $\Sigma_F(X)$  would be an infinite definable discrete subset contradicting o-minimality).  $\square$

The following proposition is a key in our proof of ALW.

**Proposition 3.5.** Let  $G$  be a connected commutative algebraic group over  $\mathbb{C}$ ,  $\mathbf{T}_G$  the Lie algebra of  $G$ , and  $\exp_G: \mathbf{T}_G \rightarrow G$  the exponential map. Assume  $\exp_G$  has a definable large domain  $F$ .

Let  $X \subseteq \mathbf{T}_G$  be a definable real analytic submanifold (or an irreducible algebraic subvariety), and  $\mathbf{T}_B < \mathbf{T}_G$  the Lie algebra of the stabilizer  $B$  of the Zariski closure of  $\exp_G(X)$  in  $G$ .

Then there is a finite set  $S \subset \mathbf{T}_G$  such that

$$X \subseteq \mathbf{T}_B + S + F.$$

*Proof.* Let  $\Lambda = \exp_G^{-1}(e)$ . It is a discrete subgroup of  $\mathbf{T}_G$ .

Let  $Z \subseteq G$  be the Zariski closure of  $\exp_G(X)$  and  $B$  be the stabilizer of  $Z$  in  $G$ .

We define  $\Sigma_F(X)$  as in (4.1).

Let  $B^0$  be the connected component of  $B$ . It is an algebraic subgroup of  $G$ , has a finite index in  $B$  and with  $\exp_G(\mathbf{T}_B) = B^0$ , where  $\mathbf{T}_B < \mathbf{T}_G$  is the Lie algebra of  $B$ .

We choose  $b_1, \dots, b_n \in B$  with  $B = \bigcup_{i=1}^n b_i \cdot B^0$ , and also choose  $h_1, \dots, h_n \in \mathbf{T}_G$  with  $\exp_G(h_i) = b_i$ . We have

$$\exp_G\left(\bigcup_{i=1}^n (h_i + \mathbf{T}_B)\right) = B,$$

hence by Claim 3.2  $\exp_G(\Sigma_F(X)) \subseteq \exp_G\left(\bigcup_{i=1}^n (h_i + \mathbf{T}_B)\right)$  and

$$\Sigma_F(X) \subseteq \mathbf{T}_B + \left(\bigcup_{i=1}^n (h_i + \Lambda)\right).$$

Since  $\Lambda$  is a discrete subgroup of  $\mathbf{T}_G$ , the set  $\bigcup_{i=1}^n (h_i + \Lambda)$  is a discrete subset of  $\mathbf{T}_G$ . By o-minimality, since  $\Sigma_F(X)$  is definable we obtain that there is a finite set  $S \subseteq \bigcup_{i=1}^n (h_i + \Lambda)$  with  $\Sigma_F(X) \subseteq \mathbf{T}_B + S$ . The proposition now follows from Observation 3.1(1).  $\square$

*Remark 3.6.* The above proposition immediately implies ALW Theorem for abelian varieties. Indeed let  $G$  be an abelian variety,  $\exp_G: \mathbf{T}_G \rightarrow G$  the exponential map,  $X \subseteq \mathbf{T}_G$  an irreducible algebraic subvariety,  $B < G$  the stabilizer of the Zariski closure of  $\exp_G(X)$  and  $\mathbf{T}_B < \mathbf{T}_G$  the Lie algebra of  $B$ .

Since  $G$  is compact, there is a relatively compact fundamental domain  $F$  for  $\exp_G$  definable in the o-minimal structure  $\mathbb{R}_{\text{an}}$ .

Using Proposition 3.5, we have that  $X \subseteq \mathbf{T}_B + S + F$ , for some finite  $S \subseteq \mathbf{T}_G$ . Since  $F$  is relatively compact we obtain that  $X \subseteq \mathbf{T}_B + K$  for some compact  $K \subseteq \mathbf{T}_G$ .

Let  $L$  be a  $\mathbb{C}$ -linear subspace of  $\mathbf{T}_G$  complementary to  $\mathbf{T}_B$ . The projection of  $X$  to  $L$  along  $\mathbf{T}_B$  is bounded. Since  $X$  is an irreducible variety, it has to be a point. It follows then that  $X \subseteq \mathbf{T}_B + h$  for some  $h \in \mathbf{T}_G$  and  $\exp_G(X) \subseteq \exp_G(h) \cdot B$ .

#### 4. FULL ALW FOR SEMI-ABELIAN VARIETIES

In this section we prove a general statement that implies full ALW Theorem and hence also ALW Theorem for semi-abelian varieties.

**Proposition 4.1.** *Let  $G$  be a connected semi-abelian variety over  $\mathbb{C}$ ,  $\mathbf{T}_G$  the Lie algebra of  $G$ ,  $\exp_G: \mathbf{T}_G \rightarrow G$  the exponential map,  $V$  a vector group over  $\mathbb{C}$  and  $\pi: V \oplus \mathbf{T}_G \rightarrow V \times G$  the map  $\pi = \text{id}_V \times \exp_G$ .*

*Let  $Y \subseteq V \oplus \mathbf{T}_G$  be an irreducible algebraic variety and  $Z \subseteq \mathbf{T}_G \times G$  the Zariski closure of  $\pi(Y)$ . Then  $Z = Z_V \times Z_G$ , where  $Z_V$  is a subvariety of  $V$  and  $Z_G$  a translate of an algebraic subgroup of  $G$ .*

*Remark 4.2.* Since  $Z$  is the Zariski closure of  $Y$ , it is easy to see that if  $Z = Z_V \times Z_G$  then  $Z_V$  must be the Zariski closure of  $\text{pr}_V(Y)$  and  $Z_G$  must be the Zariski closure of  $\exp_G(\text{pr}_{\mathbf{T}_G}(Y))$ , where  $\text{pr}_V$  and  $\text{pr}_{\mathbf{T}_G}$  are the projections from  $V \oplus \mathbf{T}_G$  to  $V$  and  $\mathbf{T}_G$  respectively.

Before proving the proposition let's remark how it implies both versions of ALW. To get ALW we take  $V$  to be the trivial vector group  $0$ . To get full ALW we take  $V = \mathbf{T}_G$  and  $Y \subseteq \mathbf{T}_G \oplus \mathbf{T}_G$  the image of  $X$  under the diagonal map, i.e.  $Y = \{(u, u) \in \mathbf{T}_G \oplus \mathbf{T}_G: u \in X\}$ .

We now proceed with the proof of Proposition 4.1.

*Proof.* Let  $H = V \times G$ . It is a commutative algebraic group with the Lie algebra  $\mathbf{T}_H = V \oplus \mathbf{T}_G$  and with the exponential map  $\exp_H = \pi$ . Hence  $Z$  is the Zariski closure of  $\exp_H(Y)$ .

We denote the group operation of  $H$  by  $\cdot$ , and view  $V$  and  $G$  as subgroups of  $H$ . Very often for subsets  $S_1 \subseteq V$  and  $S_2 \subseteq G$  we write  $S_1 \times S_2$  instead of  $S_1 \cdot S_2$  to indicate that in this case  $S_1 \cdot S_2$  can be also viewed as the Cartesian product of  $S_1$  and  $S_2$ .

Notice that since  $\exp_H$  restricted to  $V$  is the identity map we have  $\exp_H^{-1}(e) = \exp_G^{-1}(e)$ .

Let  $\text{Stab}_H(Z)$  be the stabilizer of  $Z$  in  $H$ . It is an algebraic subgroup of  $V \times G$ . Since  $V$  is a vector group and  $G$  is a semi-abelian variety,  $\text{Stab}_H(Z)$  splits as  $\text{Stab}_H(Z) = V_0 \times B$ , where  $V_0 < V$  and  $B < G$  are algebraic subgroups.

We first show that  $Z \subseteq V \times (p \cdot B)$  for some  $p \in G$ .

**Lemma 4.3.** *We have  $Y - h \subseteq V + \mathbf{T}_B$  for some  $h \in \mathbf{T}_H$ , where  $\mathbf{T}_B < \mathbf{T}_G$  is the Lie algebra of  $B$ .*

*Proof of Lemma.* Since  $G$  is a connected semi-abelian variety it admits a short exact sequence

$$e \rightarrow G_0 \rightarrow G \rightarrow A \rightarrow e,$$

where  $A$  is an abelian variety and  $G_0$  is an algebraic torus, i.e. an algebraic group isomorphic to  $(\mathbb{C}^*, \cdot)^k$ .

We do a standard decomposition of  $\mathbf{T}_G$ .

Let  $d$  be the dimension of  $G$  and  $k$  the dimension of  $G_0$ . Let  $\Lambda = \exp_G^{-1}(e)$ . It is a discrete subgroup of  $\mathbf{T}_G$  whose  $\mathbb{C}$ -span is  $\mathbf{T}_G$ . Also  $\Lambda$  is a free abelian group of rank  $2d - k$ .

Let  $\mathbf{T}_0 < \mathbf{T}_G$  be the Lie algebra of  $G_0$ . It is a  $\mathbb{C}$ -linear subspace of  $\mathbf{T}_G$  of dimension  $k$ . Let  $\Lambda_0 = \Lambda \cap \mathbf{T}_0$ . It is easy to see that  $\Lambda_0$  is a pure subgroup of  $\Lambda$  (i.e. for  $\lambda \in \Lambda$  and  $n \in \mathbb{N}$ ,  $n\lambda \in \Lambda_0$  implies  $\lambda \in \Lambda_0$ ), hence it has a complementary subgroup  $\Lambda_a$  in  $\Lambda$ , i.e. a subgroup  $\Lambda_a$  of  $\Lambda$  with  $\Lambda = \Lambda_0 \oplus \Lambda_a$ . Let  $L_a < \mathbf{T}_G$  be the  $\mathbb{R}$ -span of  $\Lambda_a$ .

We have that  $\mathbf{T}_G = \mathbf{T}_0 \oplus L_a$ , and  $\Lambda_a$  is a lattice in  $L_a$ .

The restriction of  $\exp_G$  to  $\mathbf{T}_0$  is a complex Lie group homomorphism from  $\mathbf{T}_0$  onto  $G_0$  whose kernel is  $\Lambda_0$ . Choosing an appropriate basis for  $\mathbf{T}_0$  and after identifying  $G_0$  with  $(\mathbb{C}^*, \cdot)^k$ , we may assume that  $\mathbf{T}_0 = \mathbb{C}^k$  and the restriction of  $\exp_G$  to  $\mathbf{T}_0$  has form  $(z_1, \dots, z_k) \mapsto (e^{2\pi iz_1}, \dots, e^{2\pi iz_k})$ . In particular  $\Lambda_0 = \mathbb{Z}^k$  and the restriction of  $\exp_G$  to  $i\mathbb{R}^k$  is definable in  $\mathbb{R}_{\text{exp}}$ .

From now on we identify  $\mathbf{T}_0$  with  $\mathbb{C}^k$  and use decompositions

$$\mathbf{T}_G = \mathbb{C}^k \oplus L_a = \mathbb{R}^k \oplus i\mathbb{R}^k \oplus L_a \text{ and } \mathbf{T}_H = V \oplus \mathbb{R}^k \oplus i\mathbb{R}^k \oplus L_a.$$

Since both  $L_a/\Lambda_a$  and  $\mathbb{R}^k/\mathbb{Z}^k$  are compact we can choose relatively compact large domains  $F_a \subseteq L_a$  and  $F_0 \subseteq \mathbb{R}^k$  for  $\exp_G \upharpoonright L_a$  and  $\exp_G \upharpoonright \mathbb{R}^k$  respectively, definable in  $\mathbb{R}_{\text{an}}$

It is easy to see that  $F_0 + i\mathbb{R}^k + F_a$  is a large domain for  $\exp_G$  and  $F = V + F_0 + i\mathbb{R}^k + F_a$  is a large domain for  $\exp_H$ , both definable in  $\mathbb{R}_{\text{an,exp}}$ .

Let  $\mathbf{T}_B < \mathbf{T}_H$  be the Lie algebra of  $B$ . Since  $\exp_H^{-1}(e) = \exp_G^{-1}(e) = \Lambda$ , we apply Proposition 3.5 to  $Y$  and  $\exp_H$  and get a finite  $S \subset \mathbf{T}_H$  with  $Y \subseteq \mathbf{T}_B + S + F$ . Thus we have

$$Y \subseteq \mathbf{T}_B + S + F = V + \mathbf{T}_B + S + F_0 + i\mathbb{R}^k + F_a.$$

Since the closures of  $F_0$  and  $F_a$  are compact, we can find a compact subset  $K \subseteq \mathbf{T}_H$  with  $S + F_0 + F_a \subseteq K$ , and hence

$$(4.1) \quad Y \subseteq V + \mathbf{T}_B + i\mathbb{R}^k + K.$$

Let  $M = V + \mathbf{T}_B + i\mathbb{R}^k$ . It is an  $\mathbb{R}$ -linear subspace of  $\mathbf{T}_H$ . We first claim that  $Y \subseteq M + h$  for some  $h \in \mathbf{T}_H$ . Indeed, using elementary linear algebra it is sufficient to show that for any  $\mathbb{R}$ -linear map  $\xi: \mathbf{T}_H \rightarrow \mathbb{R}$  vanishing on  $M$  the image of  $Y$  under  $\xi$  is a point. Let  $\xi: \mathbf{T}_H \rightarrow \mathbb{R}$  be an  $\mathbb{R}$ -linear map vanishing on  $M$ . From (4.1) we obtain that  $\xi(Y)$  is bounded. Therefore, since  $Y$  is an irreducible algebraic variety and the map  $\bar{\xi}: \mathbf{T}_H \rightarrow \mathbb{C}$  given by  $\bar{\xi}: z \mapsto \xi(z) - i\xi(z)$  is a  $\mathbb{C}$ -linear map, the set  $\bar{\xi}(Y)$  must be a point. Thus we have  $Y \subseteq M + h$  for some  $h \in \mathbf{T}_H$ .

We will use the following fact that it is not difficult to prove.

**Fact 4.4.** *Let  $Y' \subseteq \mathbf{T}_H$  be an irreducible complex analytic subset. If  $W \subseteq \mathbf{T}_H$  is the  $\mathbb{R}$ -span of  $Y'$  (i.e. the smallest  $\mathbb{R}$ -linear subspace containing  $Y'$ ) then  $W$  is a  $\mathbb{C}$ -linear subspace of  $\mathbf{T}_H$ .*

*In particular if  $Y' \subseteq U$  for some  $\mathbb{R}$ -linear subspace  $U$  of  $\mathbf{T}_H$  then  $Y' \subseteq iU$ .*

Applying the above fact to  $Y' = Y - h$  we obtain

$$(4.2) \quad Y - h \subseteq M \cap iM = (V + \mathbf{T}_B + i\mathbb{R}^k) \cap (V + \mathbf{T}_B + \mathbb{R}^k).$$

Thus to finish the proof of Lemma, it remains to show that

$$(4.3) \quad (V + \mathbf{T}_B + i\mathbb{R}^k) \cap (V + \mathbf{T}_B + \mathbb{R}^k) = V + \mathbf{T}_B.$$

Since  $B$  is a semi-abelian subvariety of  $G$ , the intersection  $B_1 = B \cap G_0$  is an algebraic torus with the Lie algebra  $\mathbf{T}_{B_1} = \mathbf{T}_B \cap \mathbb{C}^k$ . Since  $B_1$  is an algebraic subtorus of  $G_0$ ,  $\mathbf{T}_{B_1}$  has a  $\mathbb{C}$ -basis in  $\Lambda \cap \mathbb{C}^k = \mathbb{Z}^k \subset \mathbb{R}^k$ .

It follows then that  $\mathbf{T}_{B_1}$  has form  $E \oplus iE$  for some  $\mathbb{R}$ -linear subspace  $E \subseteq \mathbb{R}^k$ , and hence

$$\mathbf{T}_B \cap (\mathbb{R}^k + i\mathbb{R}^k) = E \oplus iE.$$

We are now ready to show (4.3). Let  $\alpha \in (V + \mathbf{T}_B + i\mathbb{R}^k) \cap (V + \mathbf{T}_B + \mathbb{R}^k)$ . Then

$$\alpha = v_1 + u_1 + w_1 = v_2 + u_2 + iw_2$$

for some  $v_1, v_2 \in V, u_1, u_2 \in \mathbf{T}_B, w_1, w_2 \in \mathbb{R}^k$ . Since  $\mathbf{T}_H = V \oplus \mathbf{T}_G$ , we get  $v_1 = v_2$ , and  $(u_1 - u_2) = -w_1 + iw_2$ .

Thus  $-w_1 + iw_2 \in \mathbf{T}_B \cap (\mathbb{R}^k + i\mathbb{R}^k) = E \oplus iE$ , so  $w_1, w_2 \in E$ , and hence  $w_1, w_2 \in \mathbf{T}_B$ . It implies that  $\alpha \in V + \mathbf{T}_B$ , that shows (4.3). It finishes the proof of Lemma.  $\square$

We choose  $p \in G$  with  $V \cdot \exp_H(h) = V \cdot p$  and obtain

$$\exp_H(Y) \subseteq V \times (p \cdot B) \text{ for some } p \in G.$$

hence

$$(4.4) \quad Z \subseteq V \times (p \cdot B).$$

Let  $Z_V = \{v \in V : v \times p \in Z\}$ . It is an algebraic subvariety of  $V$  and we claim that  $Z = Z_V \times (p \cdot B)$ .

If  $v \in Z_V$  then  $v \cdot p \in Z$ , and since  $B$  lies in the stabilizer of  $Z$  we have  $v \times (p \cdot B) \subseteq Z$ . Hence  $Z_V \times (p \cdot B) \subseteq Z$ .

Let  $v \in V, g \in G$  with  $v \cdot g \in Z$ . Since  $B$  lies in the stabilizer of  $Z$  we have  $v \times (g \cdot B) \subseteq Z$ . By (4.4),  $v \times (g \cdot B) \subseteq V \times (p \cdot B)$ , hence  $g \cdot B = p \cdot B$ ,  $v \cdot p \in Z$ ,  $v \in Z_V$  and  $v \cdot g \in Z_V \times (p \cdot B)$ . It shows that  $Z \subseteq Z_V \times (p \cdot B)$ .  $\square$

## REFERENCES

- [1] AX, J. Some topics in differential algebraic geometry. I. Analytic subgroups of algebraic groups. *American Journal of Mathematics* 94 (1972), 1195–1204.
- [2] KLINGLER, B., ULLMO, E., AND YAFAEV, A. The hyperbolic Ax-Lindemann-Weierstraß conjecture. *Publications Mathématiques. Institut de Hautes Études Scientifiques* 123 (2016), 333–360.
- [3] ORR, M. Introduction to abelian varieties and the Ax-Lindemann-Weierstrass theorem. In *O-minimality and diophantine geometry*. Cambridge Univ. Press, Cambridge, 2015, pp. 100–128.
- [4] PILA, J. O-minimality and the André-Oort conjecture for  $\mathbb{C}^n$ . *Annals of Mathematics. Second Series* 173, 3 (2011), 1779–1840.
- [5] PILA, J., AND TSIMERMAN, J. Ax-Lindemann for  $\mathcal{A}_g$ . *Annals of Mathematics. Second Series* 179, 2 (2014), 659–681.
- [6] PILA, J., AND WILKIE, A. J. The rational points of a definable set. *Duke Mathematical Journal* 133, 3 (2006), 591–616.
- [7] PILA, J., AND ZANNIER, U. Rational points in periodic analytic sets and the Manin-Mumford conjecture. *Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl.* 19, 2 (2008), 149–162.
- [8] TSIMERMAN, J. Ax-Schanuel and o-minimality. In *O-minimality and diophantine geometry*. Cambridge Univ. Press, Cambridge, 2015, pp. 216–221.
- [9] ULLMO, E., AND YAFAEV, A. O-minimal flows on abelian varieties. *arXiv.org* (Oct. 2016).

UNIVERSITY OF HAIFA

*E-mail address:* kobi@math.haifa.ac.il

UNIVERSITY OF NOTRE DAME

*E-mail address:* sstarcke@nd.edu