

# A THEORY OF PAIRS FOR NON-VALUATIONAL STRUCTURES

ELITZUR BAR-YEHUDA, ASSAF HASSON<sup>†</sup>, AND YA'ACOV PETERZIL

ABSTRACT. Given a weakly o-minimal structure  $\mathcal{M}$  and its o-minimal completion  $\bar{\mathcal{M}}$ , we first associate to  $\bar{\mathcal{M}}$  a canonical language and then prove that  $Th(\mathcal{M})$  determines  $Th(\bar{\mathcal{M}})$ . We then investigate the theory of the pair  $(\bar{\mathcal{M}}, \mathcal{M})$  in the spirit of the theory of dense pairs of o-minimal structures, and prove, among other results, that it is near model complete, and every definable open subset of  $\bar{M}^n$  is already definable in  $\bar{\mathcal{M}}$ .

We give an example of a weakly o-minimal structure which interprets  $\bar{\mathcal{M}}$  and show that it is not elementarily equivalent to any reduct of an o-minimal trace.

## 1. INTRODUCTION

An expansion  $\mathcal{M}$  of an ordered group is *weakly o-minimal non-valuational* (below we use “non-valuational” for short) if it is weakly o-minimal (every definable subset of  $M$  is a finite union of convex sets) and does not admit any definable non-trivial convex sub-groups. Non-valuational structures were introduced in [6] and more systematically studied in [10] and [11]. In those works Wencel showed that to a non-valuational structure  $\mathcal{M}$  one can associate an o-minimal structure  $\bar{\mathcal{M}}$ , whose universe is  $\bar{M}$  – the definable Dedekind completion of  $\mathcal{M}$  – and with the additional property that the structure which  $\bar{\mathcal{M}}$  induces on (the natural embedding of)  $M$  (in  $\bar{\mathcal{M}}$ ) is precisely the structure  $\mathcal{M}$ . Wencel called the structure  $\bar{\mathcal{M}}$  *the canonical o-minimal completion of  $\mathcal{M}$* . In [5] Keren shows that  $\bar{\mathcal{M}}$  has the same definable sets as the structure  $\mathcal{M}^*$ , whose atomic sets are all sets of the form  $\text{cl}_{\bar{M}}(S) \subseteq \bar{M}^n$  for  $\mathcal{M}$ -definable  $S \subseteq M^n$ , (see Proposition 2.7 below). Both Wencel and Keren’s constructions have the problem that the signatures of the resulting structures depend on the structure  $\mathcal{M}$ , rather than on its signature.

In the present paper we address this problem by considering, for  $A \subseteq M$ , structures of the form  $\mathcal{M}_A^*$  whose atomic sets are all sets of the form  $\text{cl}_{\bar{M}}(S)$  for  $S$  an  $\mathcal{M}$ -definable set over  $A$ . The starting point of the present work, and the main result of Section 2 is:

**Theorem 1.** *Let  $\mathcal{M}$  be a non-valuational structure. Then  $\mathcal{M}_\emptyset^*$  and  $\mathcal{M}^*$  have the same definable sets. Moreover, if  $\mathcal{M} \equiv \mathcal{N}$  then  $\mathcal{M}_\emptyset^* \equiv \mathcal{N}_\emptyset^*$ .*

This result shows that to a non-valuational theory  $T$  we can associate an o-minimal theory  $T^*$  which can be viewed as an invariant of  $T$ . Consequently, any of the o-minimal properties of  $T^*$  can reflect on the weakly o-minimal  $T$  and vice versa. This plays a crucial role in the proof of Theorem 3 below.

Section 5 is dedicated to the study of the theory of the pair  $\mathcal{M}^P = (\mathcal{M}_\emptyset^*, \mathcal{M})$  for  $\mathcal{M}$  non-valuational, in the spirit of van den Dries’ study of o-minimal dense pairs (see [9]). Our main result is the following:

**Theorem 2.** *Let  $\mathcal{M}$  be non-valuational.*

- (1) *If  $\mathcal{M} \equiv \mathcal{N}$  then  $\mathcal{M}^P \equiv \mathcal{N}^P$ .*

*We let  $T^P = Th(\mathcal{M}^P)$  and assume  $\tilde{\mathcal{N}} = (\mathcal{N}', \mathcal{N}) \models T^P$ .*

- (2) *If  $Y \subseteq (N')^n$  is  $\emptyset$ -definable in  $\tilde{\mathcal{N}}$  then it can be written as a boolean combination of sets defined by formulas of the form*

$$(1) \quad \exists x_1 \cdots \exists x_k \left( \bigwedge_{i=1}^k x_i \in P \ \& \ \varphi(x_1, \dots, x_k, y) \right),$$

*and  $\varphi(x, y)$  is a formula of the o-minimal structure  $\mathcal{M}'$ .*

- (3) *If  $X \subseteq N^k$  is definable in  $\tilde{\mathcal{N}}$  over  $A \subseteq N$  then  $X$  is already definable in the weakly o-minimal  $\mathcal{N}$ .*

---

*Date:* November 28, 2017.

<sup>†</sup> Supported by ISF grant No. 181/16.

(4) If  $U \subseteq (N')^k$  is a definable open set in  $\tilde{\mathcal{N}}$  then  $U$  is already definable in the o-minimal structure  $\mathcal{N}'$ . In particular,  $\tilde{\mathcal{N}}$  has an o-minimal open core.

The above results show that pairs  $(\mathcal{M}', \mathcal{M})$  as above fit into the setting of recent works by Eleftherious, Gunaydin and Hieronymi (see for example [3]) on expansions of o-minimal structures by dense predicates.

Non-valuational structures arise naturally in the study of dense pairs of o-minimal structures. Namely, if  $\mathcal{M} \prec \mathcal{N}$  are o-minimal expansions of ordered groups and  $M$  is dense in  $N$  then the structure induced on  $M$  from  $\mathcal{N}$  is non-valuational (weak o-minimality follows from [1] and non-valuationality is easy, see e.g., [4]). Since every ordered group which is a reduct of a non-valuational structure, or even elementarily equivalent to one, is also such, a question arises whether every non-valuational structure arises in this manner.

First, some terminology. A non-valuational structure  $\mathcal{M}$  is called an *o-minimal trace* if there is a dense pair  $\mathcal{M}_0 \prec \mathcal{N}$  such that  $\langle M_0, < \rangle = \langle M, < \rangle$  (i.e., the structures  $\mathcal{M}_0$  and  $\mathcal{M}$  have the same underlying ordered set) and the induced structure on  $M$  in the pair  $(\mathcal{N}, \mathcal{M}_0)$  has the same definable sets as  $\mathcal{M}$  (see [4] for details). In [4] we showed that an ordered reduct of a non-valuational o-minimal trace need not be an o-minimal trace itself, and that the class of reducts of o-minimal traces is not closed under elementary equivalence. In the present paper we show that even after closing the class of o-minimal traces under reducts and elementary equivalence we still do not cover all non-valuational structures:

**Theorem 3.** *Let  $\mathbb{Q}^{\sqrt{2}}$  be the expansion of  $(\mathbb{Q}, +)$  by the predicate  $y < \sqrt{2}x$ . Then  $\mathbb{Q}^{\sqrt{2}}$  is non-valuational and not elementarily equivalent to a reduct of an o-minimal trace.*

Along the way we reveal a new dividing line between two types of non-valuational structures:

- *Tight* structures (of which  $\mathbb{Q}^{\sqrt{2}}$  is a typical example), in which  $\mathcal{M}^*$  is interpretable in  $\mathcal{M}$ . These are *small* (in the sense of [9]), and in that respect differ significantly from o-minimal traces.
- Non-tight structures, whose theory resembles to a much greater extent that of o-minimal traces.

*This project was initiated by the M.Sc thesis of the first author at Ben Gurion University, under the supervision of the other authors. We thank Pantelis Eleftheriou for his helpful comments.*

## 2. PRELIMINARIES

We fix a non-valuational structure  $\mathcal{M}$  and its definable completion  $\bar{M}$ . Recall that the elements of  $\bar{M}$  are all (unique) realizations of definable cuts in  $\mathcal{M}$ . These will be identified here with the definable open subsets of  $M$  that are bounded above and downward closed. The set  $\bar{M}$  is equipped with ordering by inclusion. The structure  $\langle M, < \rangle$  is naturally embedded into  $\bar{M}$  via the map  $a \mapsto (-\infty, a)$ , and from now on we will view  $M$  as a subset of  $\bar{M}$ . The topology on  $\bar{M}$  and  $\bar{M}^n$  are the order and the product topology, respectively. We let  $\text{cl}_{\bar{M}}(-)$ ,  $\partial_{\bar{M}}(-)$  denote the corresponding topological operations in  $\bar{M}^n$ . Unless otherwise stated, all definability below refers to the structure  $\mathcal{M}$ .

Recall that a partial function  $f : M^n \rightarrow \bar{M}$  is said to be *definable* if the set  $\{(x, y) \in M^{n+1} : y < f(x)\}$  is definable. Equivalently, the family of cuts  $\{y \in \bar{M} : y < f(x)\}$ , for  $x \in M^n$ , is a definable family (and can be identified with a sort in  $\mathcal{M}$ ).

We start by collecting several useful facts concerning the relationship of  $\mathcal{M}$  and various structures on  $\bar{M}$ . We first recall the definition of a strong cell  $C \subseteq M^n$  from [10]<sup>1</sup> The definition will be inductive in  $n$  and for the induction step we will also associate inductively to each strong cell  $C \subseteq M^n$  its so-called *iterative convex hull*  $\bar{C}$ ,  $C \subseteq \bar{C} \subseteq \bar{M}^n$ . Having defined  $C$  and  $\bar{C}$  below, we say that an  $\mathcal{M}$ -definable function  $f : C \rightarrow \bar{M}$  is *strongly continuous* if it extends continuously to  $\bar{f} : \bar{C} \rightarrow \bar{M}$ , and in addition either  $f(C) \subseteq M$  or  $f(C) \subseteq \bar{M} \setminus M$ . We are now ready to state the definition:

**Definition 2.1.** A set  $C \subseteq M$  is a *strong cell* if it is either a point, in which case  $\bar{C} = C$ , or an open convex set, in which case  $\bar{C}$  is defined as the convex hull of  $C$  in  $\bar{M}$ .

Inductively, If  $C \subseteq M^n$  is a strong cell (with the associated  $\bar{C} \subseteq \bar{M}^n$ ) and  $f, g : C \rightarrow \bar{M}$  are strongly continuous with  $\bar{f}(x) < \bar{g}(x)$  for all  $x \in \bar{C}$  (note the strong assumption here!) then  $\Gamma_f(C)$  – the graph of  $f$  on  $C$

<sup>1</sup>We are using Wencel's definition, in a slightly different formulation than in [6].

– and  $(f, g)_C := \{(x, y) \in M^{n+1} : f(x) < y < g(y)\}$  are strong cells. In the first case the iterative convex hull is defined to be the graph of the extension  $\bar{f} : \bar{C} \rightarrow \bar{M}$ , and in the second case it is defined to be

$$\{(x, y) \in \bar{M}^{n+1} : x \in \bar{C} \ \& \ \bar{f}(x) < y < \bar{g}(x)\}.$$

**Remark 2.2.** (1) It is easy to verify that for each strong cell  $C \subseteq M^n$  there exists a homeomorphic projection  $\pi_C : C \rightarrow D \subseteq M^k$  onto  $k$  of the coordinates,  $k \leq n$ , whose image is an open strong cell in  $M^k$ . In this case  $\dim C := k$ . The coordinate functions of  $\pi_C^{-1}$  are strongly continuous on  $D$ .

(2) Notice that each strong cell  $C$  is a subset of  $M^n$  that is definable in  $\mathcal{M}$ , and furthermore the various functions  $f$  and  $g$  in the inductive definition of  $C$  are definable in  $\mathcal{M}$ , even though they might take values in  $\bar{M} \setminus M$ . However, in general  $\bar{C} \subseteq \bar{M}^n$  is not definable in  $\mathcal{M}$  in any obvious sense because it might not be contained in finitely many sorts in  $\mathcal{M}$ .

We can now describe Wencel's canonical completion  $\bar{\mathcal{M}}$ , but we refine his definition so we have a better control of parameters.

**Definition 2.3.** Given  $A \subseteq M$ , we let  $\bar{\mathcal{M}}_A$  be the expansion of  $\bar{M}$  by all iterative convex hulls  $\bar{C} \subseteq \bar{M}^n$ , so that  $C \subseteq M^n$  is a strong cell defined over  $A$ .

It is easy to see that the order relation  $<$  is an atomic relation in  $\bar{\mathcal{M}}_A$ . Since  $\langle M, <, + \rangle$  is divisible, [6], and  $M$  is dense in  $\bar{M}$ , the group operation extends uniquely to  $\bar{M}$ , so it is strongly continuous, and its graph  $C_+$  is a strong cell whose iterative convex hull is the graph of a group operation on  $\bar{M}$  that we still denote by  $+$ .

We now collect some of the main results from [11]

**Fact 2.4.** Let  $\mathcal{M}$  be a weakly  $o$ -minimal non-valuational structure.

- (1) Every  $A$ -definable set has a decomposition into finitely many strong cells, each defined over  $A$ .
- (2) The structure  $\bar{\mathcal{M}}_M$  is  $o$ -minimal.
- (3) If  $X \subseteq \bar{M}^n$  is definable in  $\bar{\mathcal{M}}$  then  $X \cap M^n$  is definable in  $\mathcal{M}$ .

In [5], the language of  $\bar{\mathcal{M}}_A$  was replaced by another one, which we find more convenient to work with.

**Definition 2.5.** Given  $A \subseteq M$ , and an  $A$ -definable set  $X \subseteq M^n$  in  $\mathcal{M}$ , we associate to  $X$  a predicate symbol  $\hat{X}$ . We interpret  $\hat{X}$  in  $\bar{M}^n$  as the topological closure of  $X$  in  $\bar{M}^n$ , denoted by  $\text{cl}_{\bar{M}}(X)$ , and let  $\mathcal{M}_A^*$  be the expansion of  $\bar{M}$  by all  $\hat{X}$ , for  $X \subseteq M^n$  definable over  $A$ .

It was proved in [5] that the structures  $\bar{\mathcal{M}}_M$  and  $\mathcal{M}_M^*$  have the same definable sets. We re-prove here a more precise version. We first prove:

**Lemma 2.6.** If  $C \subseteq M^n$  is a strong cell then  $\text{cl}_{\bar{M}}(C) = \text{cl}_{\bar{M}}(\bar{C})$ .

*Proof.* Since  $C \subseteq \bar{C}$  it suffices to show that  $\bar{C} \subseteq \hat{C}$  for every strong cell  $C$ . We use induction on  $n$ .

If  $C \subseteq M$  the claim is obvious. Now, suppose that  $\bar{C} \subseteq \hat{C}$  for some strong cell  $C$  and let  $f_1, f_2 < f_3$  be strongly continuous such that the range of  $f_1$  is in  $M$ . We let  $C_1 = \Gamma(f_1)_C$  and  $C_2 = (f_2, f_3)_C$  be the associated strong cells, and will show that  $\bar{C}_1 \subseteq \hat{C}_1$  and  $\bar{C}_2 \subseteq \hat{C}_2$ .

Let  $(c, m) \in \bar{C} \times \bar{M}$ . If  $(c, m) \in \bar{C}_1$  then  $\bar{f}_1(c) = m$ . But then since  $c \in \bar{C}$  and  $\Gamma(f_1)_C$  is dense in  $\Gamma(\bar{f}_1)_{\bar{C}}$  (because  $C$  is dense in  $\bar{C}$ ) then  $(c, m)$  is a limit point of  $f_1$  and therefore  $(c, m) \in \hat{C}_1$ . If  $(c, m) \in \bar{C}_2$  then  $\bar{f}_2(c) < m < \bar{f}_3(c)$ , and again since  $c \in \bar{C}$  and  $(f_2, f_3)_C$  is dense in  $(\bar{f}_2, \bar{f}_3)_{\bar{C}}$  then  $(c, m)$  is a limit point of  $(f_2, f_3)_C$  and therefore  $(c, m) \in \hat{C}_2$ .  $\square$

We can now prove:

**Proposition 2.7.** For every  $A \subseteq M$ , the ( $o$ -minimal) structures  $\mathcal{M}_A^*$  and  $\bar{\mathcal{M}}_A$  have the same  $\emptyset$ -definable sets (so in particular the same definable sets).

*Proof.* We first show that every atomic set in  $\mathcal{M}_A^*$  is  $\emptyset$ -definable in  $\bar{\mathcal{M}}_A$ . So we take an  $A$ -definable  $X \subseteq M^k$ , and consider its closure  $\hat{X} \subseteq \bar{M}^k$ . By Fact 2.4,  $X$  can be written as the union  $\bigcup_{i=1}^k C_i$  of strong cells that are definable over  $A$  in  $\mathcal{M}$ . By Lemma 2.6, each  $\bar{C}_i$  is dense in  $\text{cl}_{\bar{M}}(C_i)$ . It follows that  $\text{cl}_{\bar{M}}(X) = \bigcup_{i=1}^k \text{cl}_{\bar{M}}(\bar{C}_i)$ . Since each  $\bar{C}_i$  is  $\emptyset$ -definable in  $\bar{\mathcal{M}}_A$ , and the closure operation is itself definable, it follows that  $\text{cl}_{\bar{M}}(X)$  is  $\emptyset$ -definable in  $\bar{\mathcal{M}}_A$ .

For the other inclusion, we need to see that for every strong cell  $C \subseteq M^n$  that is definable over  $A$ , the set  $\bar{C}$  is  $\emptyset$ -definable in  $\mathcal{M}_A^*$ . This is done by induction on  $n$ .

For 0-cells in  $\mathcal{M}$  this is clear. If  $C \subseteq M$  is a 1-cell then  $\bar{C}$  is an open interval  $(a, b)$  in  $\bar{M}$ . The interval  $[a, b]$  is  $\emptyset$ -definable in  $\mathcal{M}_A^*$ , hence so is  $\bar{C}$ . So we now assume that we have proved the result for all strong cells in  $M^n$  and we prove it for strong-cells in  $M^{n+1}$ . Let  $C \subseteq M^n$  be a strong cell defined over  $A$ . Let  $f : C \rightarrow M$  be a strongly continuous function definable in  $\mathcal{M}$  over  $A$ , and let  $Y$  be  $\Gamma_f$ , the graph of  $f$ . Then  $\bar{Y} := \{(x, \bar{f}(x)) : x \in \bar{C}\}$ . We have to show that  $\bar{Y}$  is  $\emptyset$ -definable in  $\mathcal{M}_A^*$ . As  $\bar{f}$  is continuous we get that  $\bar{Y} = \text{cl}_{\bar{M}}(\Gamma_f) \cap (\bar{C} \times \bar{M})$ , which is  $\emptyset$ -definable in  $\mathcal{M}_A^*$  by the inductive hypothesis.

Now let  $f, g : C \rightarrow \bar{M}$  be  $A$ -definable strongly continuous functions in  $\mathcal{M}$ , with  $f < g$  (unlike the above, we cannot assume here that they take values in  $M$ ). We have to show that the iterative convex hull of  $Y := (f, g)_C$  is  $\emptyset$ -definable in  $\mathcal{M}_A^*$ . By definition,

$$\bar{Y} = \{(x, y) : x \in \bar{C}, \bar{f}(x) < y < \bar{g}(x)\}.$$

Since, by induction  $\bar{C}$  is  $\emptyset$ -definable in  $\mathcal{M}_A^*$ , it will suffice to show that  $\bar{f}$  (and similarly  $\bar{g}$ ) is  $\emptyset$ -definable in  $\mathcal{M}_A^*$ . If  $f$  is the constant function  $-\infty$ , then there is nothing to prove. So we assume this is not the case. By definition, the set

$$F := \{(x, y) : x \in C, y < f(x)\}$$

is  $A$ -definable in  $\mathcal{M}$ . For every  $c \in \bar{C}$  let

$$s(c) := \sup\{y \in \bar{M} : (c, y) \in \text{cl}_{\bar{M}}(F)\}.$$

Since  $f$  is strongly continuous,  $s(c)$  is well defined, and by definition it coincides with  $f$  on  $C$ . Since  $C$  is dense in  $\bar{C}$  and  $\bar{f}$  is the unique continuous extension of  $f$  to  $\bar{C}$ , necessarily  $s = \bar{f}$ , and as  $s$  is  $\emptyset$ -definable in  $\mathcal{M}_A^*$ , we are done.  $\square$

From now on we can use interchangeably the structures  $\mathcal{M}_A^*$  and  $\bar{\mathcal{M}}_A$ . Notice however, that the language of  $\bar{\mathcal{M}}_A$  depends on the specific structure  $\mathcal{M}$ , thus for different  $\mathcal{M}$  and  $\mathcal{N}$ , even if elementarily equivalent, the structures  $\bar{\mathcal{M}}_M$  and  $\bar{\mathcal{N}}_N$  are of different signature. One of the initial goals of this work was to obtain a uniform signature by showing that the definable sets in  $\bar{\mathcal{M}}_\emptyset$  and  $\bar{\mathcal{M}}_M$  are the same. We need the following observations

- Proposition 2.8.** (1) *Every  $\emptyset$ -definable set in  $\mathcal{M}$  can be written as a boolean combination of  $\emptyset$ -definable sets each of which is the closure of an open  $\emptyset$ -definable set. In particular, this is true if  $\mathcal{M}$  is o-minimal.*  
(2) *The o-minimal structure  $\mathcal{M}_M^*$  eliminates quantifiers. Moreover, it is sufficient to take as atomic relations all  $\text{cl}_{\bar{M}}(X)$  with  $X \subseteq M^n$  an open definable set.*

*Proof.* (1) We first prove the result for an arbitrary definable open set  $X \subseteq M^n$ . Note that  $X = \text{cl}(X) \setminus \partial(X)$  (here  $\partial(X)$  is the boundary of  $X$ ), and then that

$$\partial(X) = \text{cl}(X) \cap \text{cl}(M^n \setminus \text{cl}(X)).$$

The set on the right is of the desired form, so we are done.

For an arbitrary definable  $X \subseteq M^n$ , we apply strong cell decomposition, so we may assume that  $X$  is a cell. Hence,  $X$  is either a point or the graph of a definable map  $f$  from an open cell  $C \subseteq M^{n-k}$  into  $M^k$  (the  $n - k$  coordinates need not be the first ones), and each of the coordinate functions of  $f$  are strongly continuous.

Thus it is sufficient to show that the graph of each strongly continuous  $f_i : C \rightarrow M$  is definable in the desired form. By the continuity of  $f_i$ , such a graph can be written as the complement in  $C \times M$  of the open set:

$$\{(x, y) \in C \times M : y > f_i(x)\} \cup \{(x, y) \in C \times M : y < f_i(x)\}.$$

Since each of the open sets can be defined in the required form, so is the graph of  $f_i$ , and hence so is  $X$ .  $\square$

For (2), we first apply (1) to the o-minimal structure  $\mathcal{M}_M^*$  and reduce the problem to definable sets  $\hat{X} \subseteq \bar{M}^n$ , which are the closure of an open definable set  $U \subseteq M^n$ . Since  $M^n$  is dense in  $\bar{M}^n$ ,  $\text{cl}_{\bar{M}}(U) = \text{cl}_{\bar{M}}(U \cap M^n)$ . By fact 2.4, the set  $U \cap M^n$  is definable in  $\mathcal{M}$  (possibly over parameters). We now apply (1).  $\square$

In the text the first part of the above proposition will be applied, mostly, when  $\mathcal{M}$  is, in fact, o-minimal.

**Lemma 2.9.** *Let  $C \subseteq M^{k+n}$  be a strong cell,  $a \in \pi(C)$ , where  $\pi$  is the projection onto the first  $k$ -coordinate. Let  $C_a = \{x \in M^n : (a, x) \in C\}$ . Then*

- (1)  $C_a$  is a strong cell.
- (2)  $(\bar{C})_a = \bar{C}_a$ .

*Proof.* It is sufficient to prove the result for  $k = 1$  (and then proceed by induction). This is straightforward from the definition of a strong cell.  $\square$

**Theorem 2.10.** *For every  $A \subseteq M$ , the structures  $\bar{\mathcal{M}}_M$  and  $\bar{\mathcal{M}}_A$  have the same definable sets.*

*Proof.* Absorbing  $A$  to the language we, at this stage, assume that  $A = \emptyset$ . We first claim that for every  $n \in \mathbb{N}$ , we have

$$(2) \quad \{Y \cap M^n : Y \subseteq \bar{M}^n \text{ definable in } \bar{\mathcal{M}}_\emptyset\} = \{Y \cap M^n : Y \subseteq \bar{M}^n \text{ definable in } \bar{\mathcal{M}}_M\}.$$

Since  $\bar{\mathcal{M}}_\emptyset$  is a reduct of  $\bar{\mathcal{M}}_M$  it is sufficient to prove the right-to-left inclusion. We first show: For every  $\emptyset$ -definable  $X \subseteq M^n$ , there exists a  $\emptyset$ -definable  $Y \subseteq \bar{M}^n$  in  $\bar{\mathcal{M}}_\emptyset$  such that  $Y \cap M^n = X$ . Indeed,  $X$  has a decomposition into  $\emptyset$ -definable strong cells (see Fact 2.4), and for each  $\emptyset$ -definable strong cell  $C_i$  we have  $\bar{C}_i \cap M^n = C_i$ , so  $Y = \bigcup_i \bar{C}_i$  is the desired set.

Now, let  $Z \subseteq M^n$  be definable in  $\bar{\mathcal{M}}_M$ . By Fact 2.4,  $Z \cap M^n$  is definable in  $\mathcal{M}$ , possibly over parameters. Hence, it is of the form  $X_a$ , for some  $X \subseteq M^{n+k}$  which is  $\emptyset$ -definable in  $\mathcal{M}$  and  $a \in M^k$ . By what we just shown, there is  $Y \subseteq \bar{M}^k$  which is  $\emptyset$ -definable in  $\bar{\mathcal{M}}_\emptyset$ , such that  $X = Y \cap M^{n+k}$ . Hence,

$$Z = (Y \cap M^{n+k})_a = Y_a \cap M^n,$$

and  $Y_a$  is definable in  $\bar{\mathcal{M}}_\emptyset$ . This ends the proof of (2).

We now make the following general observation:

**Lemma 2.11.** *Let  $\langle N, < \rangle$  be a densely ordered set, with  $M \subseteq N$  a dense subset. Assume that  $\mathcal{N}_1, \mathcal{N}_2$  are two o-minimal expansions of  $\langle N, < \rangle$  with the property that for every  $n \in \mathbb{N}$ , we have*

$$(3) \quad \{Y \cap M^n : Y \subseteq \bar{M}^n \text{ definable in } \mathcal{N}_1\} = \{Y \cap M^n : Y \subseteq \bar{M}^n \text{ definable in } \mathcal{N}_2\}.$$

*Then  $\mathcal{N}_1$  and  $\mathcal{N}_2$  have the same definable sets.*

*Proof.* It easily follows from the assumptions that we have

$$(4) \quad \{Y \cap M^n : Y \subseteq \bar{M}^n \text{ open definable in } \mathcal{N}_1\} = \{Y \cap M^n : Y \subseteq \bar{M}^n \text{ open definable in } \mathcal{N}_2\}.$$

By Proposition 2.8 (1), it is enough to know that for every open  $U \subseteq N^n$ , the set  $\text{cl}(U)$  is definable in  $\mathcal{N}_1$  if and only if it is definable in  $\mathcal{N}_2$ . However, since  $M$  is dense in  $N$ , it is enough to consider sets of the form  $\text{cl}(U \cap M^n)$ . By (4), both collections of sets of the form  $U \cap M^n$ , where  $U$  is definable in either  $\mathcal{N}_1$  or in  $\mathcal{N}_2$ , are the same..  $\square$

In order to prove Theorem 2.10, we apply Lemma 2.11 to the structures  $\bar{\mathcal{M}}_\emptyset$  and  $\bar{\mathcal{M}}_M$  using (2).  $\square$

### 3. THE STRUCTURE $\mathcal{M}_A$ AND ELEMENTARY EXTENSIONS

Again, we let  $\mathcal{M}$  be a fixed non-valuational structure. From now on we shall work with  $\mathcal{M}_A^*$  rather than  $\bar{\mathcal{M}}_A$ .

**3.1. The canonical completion and elementary extensions.** Let  $\mathcal{N}$  be an elementary extension of  $\mathcal{M}$ . Every definable cut  $C$  in  $\mathcal{M}$  has a natural realization  $C(\mathcal{N})$  in  $\mathcal{N}$  and so  $\bar{M}$  can be embedded into  $\bar{N}$ . Under this embedding, if  $n \in \bar{N}$  is the supremum of a cut in  $N$  which is definable over some  $A \subseteq M$  then  $n$  is already in  $\bar{M}$ . We have:

$$\begin{array}{ccc} \mathcal{N} & \xrightarrow{\iota} & \bar{\mathcal{N}} \\ & \prec & \uparrow \\ \mathcal{M} & \xrightarrow{\iota} & \bar{\mathcal{M}} \end{array}$$

Where  $\iota$  is the natural embedding of  $(M, <)$  in  $(\bar{M}, <)$ . We now fix an arbitrary  $A \subseteq M$  and consider the structures  $\mathcal{M}_A^*$  and  $\mathcal{N}_A^*$ . Both structures are in the language  $\mathcal{L}_A^*$ , and we claim that  $\mathcal{M}_A^*$  is a substructure of  $\mathcal{N}_A^*$ : Indeed, first note that for a fixed  $x \in \bar{M}^n$ , and  $\epsilon > 0$  in  $M$ , the set  $B(x, \epsilon) \cap M^n = \{y \in M^n : |x - y| < \epsilon\}$  is

definable in  $\mathcal{M}$  and moreover, it is uniformly definable as  $\epsilon$  varies in  $M_{>0}$  ( $x$  still fixed). It easily follows that for  $x \in \bar{M}^n$ , being in the closure of a definable  $X \subseteq M^n$  is a first order property. Namely, for  $x \in \bar{M}^n$ ,

$$x \in \text{cl}_{\bar{M}}(X(M)) \Leftrightarrow x \in \text{cl}_{\bar{N}}(X(N)).$$

Said differently,  $\hat{X}(N) \cap \bar{M}^n = \hat{X}(M)$ , so  $\mathcal{M}_A^*$  is a substructure of  $\mathcal{N}_A^*$ .

Our goal is to show that  $\mathcal{M}_A^*$  is in fact an elementary substructure of  $\mathcal{N}_A^*$ . We do that in several steps.

**Lemma 3.1.** *Assume that  $A \subseteq M$  and that  $\mathcal{M}$  is  $|A|^+$ -saturated. If  $Y \subseteq \bar{M}^n$  is  $\emptyset$ -definable in  $\mathcal{M}_A^*$  then  $Y \cap M^n$  is  $A$ -definable in  $\mathcal{M}$ .*

*Proof.* By fact 2.4,  $Y \cap M^n$  is definable in  $\mathcal{M}$ . By the saturation assumption it is enough to show that any automorphism of  $\mathcal{M}$  which fixes  $A$  point-wise leaves  $Y \cap M^n$  invariant. Let  $\alpha : M \rightarrow M$  be such an automorphism. We claim that  $\alpha$  has a (unique) extension to a bijection  $\bar{\alpha} : \bar{M} \rightarrow \bar{M}$  which is an automorphism of  $\mathcal{M}_A^*$ . Because  $\alpha$  is an automorphism of  $\mathcal{M}$  it sends definable cuts to definable cuts so extends naturally to  $\bar{\alpha} : \bar{M} \rightarrow \bar{M}$ . The map  $\bar{\alpha}$  is an order preserving bijection so in particular continuous on  $\bar{M}$ . To see that  $\bar{\alpha}$  is an automorphism of  $\mathcal{M}_A^*$ , let  $X \subseteq M^n$  be  $A$ -definable and consider its closure  $\hat{X}$ . Since  $\alpha(X) = X$ , continuity implies that  $\bar{\alpha}(\hat{X}) = \hat{X}$ , thus  $\bar{\alpha}$  is an automorphism of  $\mathcal{M}_A^*$ .

Since  $Y$  was  $\emptyset$ -definable in  $\mathcal{M}_A^*$  it is left invariant under  $\bar{\alpha}$ , and because  $\bar{\alpha}(M) = M$ , we have

$$\alpha(Y \cap M^n) = \bar{\alpha}(Y \cap M^n) = \bar{\alpha}(Y) \cap \bar{\alpha}(M^n) = Y \cap M^n.$$

□

**Lemma 3.2.** *For  $A \subseteq M$  arbitrary, if  $\mathcal{M} \prec \mathcal{N}$  then  $\mathcal{M}_A^* \prec \mathcal{N}_A^*$ .*

*Proof.* First note that we may assume that  $\mathcal{N}$  is sufficiently saturated. Indeed, we may consider  $\mathcal{N}' \succ \mathcal{N}$  which is saturated enough. The above would then imply that  $\mathcal{M}_A^* \prec (\mathcal{N}'_A)^*$  and  $\mathcal{N}_A^* \prec (\mathcal{N}'_A)^*$ , from which it follows that  $\mathcal{M}_A^* \prec \mathcal{N}_A^*$ .

By The Tarski-Vaught Criterion, it is enough to prove, for every nonempty  $Y \subseteq \bar{N}$  which is definable in  $\mathcal{N}_A^*$  over  $\bar{M}$ , that  $Y \cap \bar{M} \neq \emptyset$ .

Since  $\mathcal{N}_A^*$  is an o-minimal expansion of a group,  $Y$  contains some element  $b \in \text{dcl}_{\mathcal{N}_A^*}(\bar{M})$ . So, there exists a finite tuple  $a = (a_1, \dots, a_r)$  from  $\bar{M}$ , such that  $b \in \text{dcl}_{\mathcal{N}_A^*}(a)$ . Each  $a_i$  realizes a cut in  $M$ , definable in  $\mathcal{M}$  over some finitely many parameters. Thus there is a finite  $F \subseteq M$  such that each  $a_i$  realizes a cut definable over  $F$ . If we now let  $A' = A \cup F \subseteq M \subseteq N$  then clearly every element in  $A'$  is  $\emptyset$ -definable in  $\mathcal{N}_{A'}^*$ , hence  $b$  is in  $\text{dcl}_{\mathcal{N}_{A'}^*}(\emptyset)$  so the set  $(-\infty, b)$  is  $\emptyset$ -definable in  $\mathcal{N}_{A'}^*$ . Since  $\mathcal{N}$  is sufficiently saturated it follows from Lemma 3.1 that  $(-\infty, b) \cap N$  is  $A'$ -definable in  $\mathcal{N}$ .

Since  $\mathcal{M} \prec \mathcal{N}$  and  $A' \subseteq M$  it follows, as we already noted above, that  $b \in \bar{M}$ , so  $Y \cap \bar{M} \neq \emptyset$ . Thus  $\mathcal{M}_A^* \prec \mathcal{N}_A^*$ . □

**Note:** It only makes sense to compare  $\mathcal{M}_A^*$  and  $\mathcal{N}_A^*$  for  $A \subseteq M$ , since otherwise the two structures do not have a common language.

Finally, we can now prove:

**Theorem 3.3.** *For  $A \subseteq M$  (with no saturation assumption), assume that  $X \subseteq \bar{M}^n$  is  $\emptyset$ -definable in the structure  $\mathcal{M}_A^*$ . Then  $X \cap M^n$  is  $A$ -definable in  $\mathcal{M}$ . In particular, if  $f : \bar{M}^n \rightarrow \bar{M}$  is  $\emptyset$ -definable in  $\mathcal{M}_A^*$  then  $f \upharpoonright M^n : M^n \rightarrow \bar{M}$  is  $A$ -definable in  $\mathcal{M}$ .*

*Proof.* We consider an elementary extension  $\mathcal{N}$  of  $\mathcal{M}$  that is  $|A|^+$ -saturated. By Lemma 3.2, we have  $\mathcal{M}_A^* \prec \mathcal{N}_A^*$  and by Lemma 3.1, the set  $Y = X(\bar{N}) \cap N^n$  is definable in  $\mathcal{N}$  over  $A$ . Since  $\mathcal{M} \prec \mathcal{N}$  we can conclude that  $Y \cap M^n = X(\bar{N}) \cap M^n$  is also definable over  $A$  in  $\mathcal{M}$ . It is left to see that this last set equals  $X \cap M^n$ . Because  $\mathcal{M}_A^* \prec \mathcal{N}_A^*$  we have  $X(\bar{N}) \cap \bar{M}^n = X$ , and therefore

$$Y \cap M^n = X(\bar{N}) \cap M^n = X \cap M^n.$$

For the second clause, just note that the set  $\{x \in M^n : x < f(x)\}$  is the intersection of a  $\emptyset$ -definable subset of  $\bar{M}^n$  with  $M^n$ . □

We now return to Proposition 2.8 and Theorem 2.10 and prove finer results:

**Proposition 3.4.** *For any  $A \subseteq M$ ,*

- (1) *The structure  $\mathcal{M}_A^*$  eliminates quantifiers. In fact every  $\emptyset$ -definable set is a boolean combination of sets of the form  $\text{cl}_{\bar{M}}(X)$  for  $X \subseteq M^n$  open and definable in  $\mathcal{M}$  over  $A$ .*
- (2) *If  $X \subseteq \bar{M}^n$  is  $\emptyset$ -definable in  $\mathcal{M}_A^*$  then it is  $A$ -definable in  $\mathcal{M}_\emptyset^*$ .*

*Proof.* (1) We may repeat the short argument in the proof of 2.8 with the additional data given by Theorem 3.3, that whenever  $X \subseteq \bar{M}^n$  is  $\emptyset$ -definable in  $\mathcal{M}_A^*$ , the set  $X \cap M^n$  is  $A$ -definable in  $\mathcal{M}$ . For (2), assume that  $Z$  is  $\emptyset$ -definable in  $\mathcal{M}_A^*$ . By (1),  $Z$  is a boolean combination of atomic sets (with no extra parameters), so it is sufficient to prove that each atomic such set  $Z$  is  $A$ -definable in  $\mathcal{M}_\emptyset^*$ . By the first paragraph of the proof of Proposition 2.7  $Z = \bigcup_{i=1}^k \text{cl}_{\bar{M}}(\bar{C}_i)$  for some  $A$ -definable strong cells  $C_i \subseteq M^n$ . So  $C_i = (D_i)_a$  for some  $\emptyset$ -definable set  $D_i$  and  $a \subseteq A$ . By strong cell decomposition, each  $D_i$  is itself a finite union of  $\emptyset$ -definable strong cells, so we may write each  $C_i$  as a union of the form  $\bigcup_j (D_{i,j})_a$ , where each  $D_{i,j}$  is a  $\emptyset$ -definable strong cell.

By Lemma 2.9 we know that  $(D_{i,j})_a = (\overline{D_{i,j}})_a$  for every  $j$ . The right-hand side of this equation is  $A$ -definable in  $\mathcal{M}_\emptyset^*$ , and hence so is its  $\bar{M}$ -closure. Therefore the closure of each  $C_i$  is a finite union of sets that are  $A$ -definable in  $\mathcal{M}_\emptyset^*$ . The conclusion follows.  $\square$

Since any two elementarily equivalent structures have a common elementary extension, we can also conclude from Lemma 3.2:

**Corollary 3.5.** *If  $\mathcal{M}$  is non-valuational and  $\mathcal{N} \equiv \mathcal{M}$  then  $\bar{\mathcal{M}}_\emptyset \equiv \bar{\mathcal{N}}_\emptyset$  (both are  $\bar{\mathcal{L}}_\emptyset$ -structures), and  $\mathcal{M}_\emptyset^* \equiv \mathcal{N}_\emptyset^*$  (as  $\mathcal{L}_\emptyset^*$ -structures)*

Finally, we shall be using the following technical lemma:

**Lemma 3.6.** *For every  $A \subseteq M$ ,  $\text{dcl}_{\mathcal{M}_\emptyset^*}(A) \cap M = \text{dcl}_{\mathcal{M}}(A)$ .*

*Proof.* Assume that  $a \in \text{dcl}_{\mathcal{M}_\emptyset^*}(A) \cap M$ . Then it follows that  $a \in \text{dcl}_{\mathcal{M}_A^*}(\emptyset)$  (since each element of  $A$  is  $\emptyset$ -definable in  $\mathcal{M}_A^*$ ). Hence, the interval  $(-\infty, a) \subseteq \bar{M}$  is  $\emptyset$ -definable in  $\mathcal{M}_A^*$ , so by Theorem 3.3, the intersection of  $(-\infty, a)$  with  $M$  is  $A$ -definable in  $\mathcal{M}$ . Because  $a \in M$ , we have  $a \in \text{dcl}_{\mathcal{M}}(A)$ .

For the converse, assume that  $a \in \text{dcl}_{\mathcal{M}}(A)$  (so in particular in  $M$ ). Thus, the interval  $(-\infty, a)$  is definable in  $\mathcal{M}$  over  $A$  and its iterative convex hull, the interval  $(\infty, a) \subseteq \bar{M}$ , is  $\emptyset$ -definable in  $\mathcal{M}_A$ . By Proposition 2.8(2), this interval is  $A$ -definable in  $\mathcal{M}_\emptyset^*$  so  $a \in \text{dcl}_{\mathcal{M}_\emptyset^*}(A) \cap M$ .  $\square$

#### 4. TIGHT WEAKLY O-MINIMAL STRUCTURES

As was pointed out before, the set  $\bar{M}$  can be viewed as a union of sorts in  $\mathcal{M}$ , where each sort corresponds to a  $\emptyset$ -definable family of cuts in  $\mathcal{M}$ . In general, there might be infinitely many such sorts, but in some cases there are only finitely many such sorts.

##### 4.1. Definition and basic properties.

**Definition 4.1.** A non-valuational structure  $\mathcal{M}$  is *tight* if there are finitely many  $\emptyset$ -definable families of cuts in  $\mathcal{M}$  such that every definable cut belongs to one of them.

Clearly, if  $\mathcal{M}$  is an o-minimal structure then it is (trivially) non-valuational and tight, since the family of definable cuts is just all intervals of the form  $(-\infty, a)$ , as  $a$  varies in  $M$ .

It immediately follows that if  $\mathcal{M} \equiv \mathcal{N}$  then  $\mathcal{M}$  is tight if and only if  $\mathcal{N}$  is tight. Thus, we may use the term “tight” for  $T$  as well.

**Proposition 4.2.** *The structure  $\mathcal{M}$  is tight if and only if there are finitely many  $\emptyset$ -definable functions  $f_i : M^{n_i} \rightarrow \bar{M}$ ,  $i = 1, \dots, k$ , such that  $\bar{M} \subseteq \bigcup_{i=1}^k \text{Im}(f_i)$ .*

*In particular,  $\mathcal{M}$  is tight then the structure  $\mathcal{M}^*$  is interpretable in  $\mathcal{M}$  without parameters.*

*Proof.* The first clause is easy to verify. For the second clause, note first that the universe of  $\bar{M}$  is a quotient of some  $M^n$  by a definable set, and furthermore the embedding of  $M$  in this quotient (i.e. the family of cuts  $\{C_x : x \in M\}$ , where  $C_x = \{y < x\}$ ) is definable in  $\mathcal{M}$ . It is easy to see that the ordering on  $\bar{M}$  is definable in  $\mathcal{M}$  and hence  $\text{cl}_{\bar{M}}(X)$  is definable in  $\mathcal{M}$  for every  $\mathcal{M}$ -definable  $X \subseteq M^n$ .  $\square$

**Remark 4.3.** The above proof shows, in fact, that the pair  $(\mathcal{M}^*, \mathcal{M})$  is bi-interpretable with  $\mathcal{M}$ , i.e., not only is  $\mathcal{M}^*$  interpretable in  $\mathcal{M}$ , but so is the natural embedding of  $M$  in  $\bar{M}$ .

**4.2. An example of a tight structure.** We shall now see that there are examples of tight structures which are not o-minimal.

Let  $\mathbb{Q}_{vs} = \langle \mathbb{Q}, <, +, 1, \{\lambda_q\}_{q \in \mathbb{Q}} \rangle$  denote the group of rational numbers, viewed as an ordered vector space over itself, with a function symbol for every rational scalar. Let  $\mathbb{Q}^{\sqrt{2}}$  be the expansion of  $\mathbb{Q}_{vs}$  by the relation

$$P_{\sqrt{2}} = \{(x, y) \in \mathbb{Q}^2 : y < \sqrt{2}x\}.$$

We denote the language by  $\mathcal{L}_{\sqrt{2}}$ . (In [4, Section 3] a similar expansion of  $\mathbb{Q}_{vs}$  by the predicate  $P_\pi$  was investigated.)

The idea is to eventually identify  $P_{\sqrt{2}}$  with a map  $x \mapsto \sqrt{2}x$  from the structure  $\mathbb{Q}^{\sqrt{2}}$  into its canonical completion. Our goal is to show that  $\text{Th}(\mathbb{Q}^{\sqrt{2}})$  is axiomatised by the following theory  $T$ :

- (1) The ordered  $\mathbb{Q}$ -vector space axioms.
- (2) An axiom expressing the fact that  $P_{\sqrt{2}}$  is “linear”:

$$(\forall x_1, y_1, x_2, y_2) ((x_1, y_1) \in P_{\sqrt{2}} \wedge (x_2, y_2) \in P_{\sqrt{2}} \rightarrow (x_1 + x_2, y_1 + y_2) \in P_{\sqrt{2}}).$$

- (3) (Ensuring that we define the positive  $\sqrt{2}$ ):  $(\exists x, y)((x, y) \in P_{\sqrt{2}} \wedge x > 0 \wedge y > 0)$
- (4) For all  $r \in \mathbb{Q}$ , such that  $r < \sqrt{2}$ , we have  $\forall x (x > 0 \rightarrow (x, rx) \in P_{\sqrt{2}})$ , and for all  $r \in \mathbb{Q}$  such that  $r > \sqrt{2}$ , we have  $\forall x (x > 0 \rightarrow (x, rx) \notin P_{\sqrt{2}})$ .
- (5) For all  $x \neq 0$ , the set

$$\{y : (x, y) \in P_{\sqrt{2}}\}$$

is closed downwards, and has no supremum. Furthermore,

$$\text{Inf} \{y_2 - y_1 : (x, y_1) \in P_{\sqrt{2}} \& (x, y_2) \notin P_{\sqrt{2}}\} = 0.$$

- (6) An axiom expressing the fact that the composition of  $x \mapsto \sqrt{2}x$  with itself yields the map  $x \mapsto 2x$ :

$$\forall (x, y > 0) ((\exists z > 0) P_{\sqrt{2}}(x, z) \wedge P_{\sqrt{2}}(z, y)) \iff y < 2x).$$

- (7) The quantifier-free theory of  $\mathbb{Q}^{\sqrt{2}}$ .

Clearly,  $\mathbb{Q}^{\sqrt{2}}$  is a model of  $T$ .

For simplicity we write  $F = \mathbb{Q}(\sqrt{2})$ . Before we prove quantifier elimination we note that if  $\mathcal{M}$  is a model of  $T$  then we may consider the associated  $F$ -vector space  $V = F \otimes_{\mathbb{Q}} M$ . If we identify  $M$  with the  $\mathbb{Q}$ -subspace  $1 \otimes M$ , then each element of  $V$  can be written uniquely as  $x + \sqrt{2}y$  for  $x, y \in M$ . We can now endow  $V$  with an ordering by declaring  $x + \sqrt{2}y > 0$  when  $(y, -x) \in P_{\sqrt{2}}$ . Indeed, the above axioms imply that this is a linear ordering of the vector space  $V$ , compatible with the ordering of  $F$ .

The definition of the ordering and Axiom (3) allows us to conclude:

- Claim 4.4.** (1) For  $x, y \in M$ , we have  $(x, y) \in P_{\sqrt{2}} \iff$  if and only if  $y < \sqrt{2}x$  in  $V$ .  
(2)  $M$  is dense in  $V$ .

We can now endow  $V$  with an  $\mathcal{L}_{\sqrt{2}}$ -structure, by interpreting  $P_{\sqrt{2}}$  as we did over  $\mathbb{Q}$ . Clause (1) above then implies that  $\mathcal{M}$  is a substructure of  $V$  as an  $\mathcal{L}_{\sqrt{2}}$ -structure.

The following lemma is similar to [4, Proposition 3.3]:

**Lemma 4.5.** *The theory  $T$  is complete and has quantifier elimination.*

*Proof.* Let  $\mathcal{Q}_1, \mathcal{Q}_2 \models T$  be  $\kappa$ -saturated models of the same cardinality. In order to prove quantifier elimination it suffices to prove (see for example [7, Corollary 3.1.6]):

If  $A$  is a substructure of  $\mathcal{Q}_1$  and  $\mathcal{Q}_2$  of cardinality smaller than  $\kappa$ , then for every  $a_1 \in \mathcal{Q}_1$  there is  $a_2 \in \mathcal{Q}_2$  such that  $a_1$  and  $a_2$  have the same quantifier-free type over  $A$ .

As above, consider the ordered  $F$ -vector spaces  $\mathcal{G}_i := F \otimes_{\mathbb{Q}} \mathcal{Q}_i$ . Since  $\mathcal{Q}_i$  is dense in  $G_i$ , and  $\mathcal{G}_i$  is o-minimal, the saturation of  $\mathcal{Q}_i$  implies that  $\mathcal{G}_i$  is also  $\kappa$ -saturated. Let  $B_i$  be the  $F$ -span of  $A$  inside  $\mathcal{G}_i$ . Then  $B_1$  and  $B_2$  are isomorphic-over- $A$  ordered vector spaces (both isomorphic to  $A + \sqrt{2}A$ , with the same ordering). Thus we may write  $B = B_1 = B_2$

Let  $p(x) := \text{tp}_{\mathcal{G}_1}(a_1/B)$ . We may assume that  $a_1 \notin A$  and hence  $a_1 \notin B$  (note that  $B \cap \mathcal{Q}_1 = A$ ). By the completeness of the theory of ordered  $F$ -vector spaces and saturation, we can find  $a_2 \in \mathcal{G}_2$  such that  $a_2 \models p(x)$ . In fact, because  $\mathcal{G}_2$  is  $\kappa$ -saturated and  $p$  is non-algebraic there is more than one such  $a_2$ , so since  $\mathcal{Q}_2$  is dense in  $G_2$ , we can find such an  $a_2$  inside  $\mathcal{Q}_2$ .



Finally, since each  $\mathcal{Q}_i$  is a substructure of  $\mathcal{G}_i$ , and  $a_1, a_2 \models p$ , it follows that the quantifier-free types of  $a_1$  and  $a_2$  over  $A$ , in the structures  $\mathcal{Q}_1, \mathcal{Q}_2$ , respectively, are the same. This completes the proof of quantifier elimination.

To see that  $T$  is complete we just notice that every model of  $T$  contains the structure  $\mathbb{Q}^{\sqrt{2}}$ , which is itself a model of  $T$ .  $\square$

**Corollary 4.6.** *The theory  $T$  is a tight weakly o-minimal non-valuational theory and  $T^*$  is the theory of ordered  $\mathbb{Q}^{\sqrt{2}}$ -ordered vector spaces (in the language  $\mathcal{L}_{\sqrt{2}}$ ).*

*Proof.* The atomic subsets of  $\mathcal{Q}$ , the universe of any  $\mathcal{Q} \models T$ , are rays with or without endpoints. By quantifier elimination the definable subsets of  $\mathcal{Q}$  are in the boolean algebra generated by those, proving the weak o-minimality. The same argument also shows that the only definable cuts are non-valuational, because so are the atomic cuts.

By the proof of lemma 4.5 and the preceding discussion, each model  $\mathcal{Q}$  of  $T$  is a dense substructure of the o-minimal structure  $V = F \otimes_{\mathbb{Q}} \mathcal{Q}$ . It is easy to verify that the intersection with  $\mathcal{Q}$  of every ray  $(-\infty, a)$  in  $V$  is definable in  $\mathcal{Q}$ , and hence every element of  $V$  realizes a definable cut in  $\mathcal{Q}$ . Conversely, by quantifier elimination, the definable cuts in any model  $\mathcal{Q}$  of  $T$  are of the form  $x_1 + r\sqrt{2}x_2$  for  $r \in \mathbb{Q}$  and  $x_1, x_2 \in \mathcal{Q}$ , so they are realized in  $V$ . It follows that  $V$  is the canonical completion of  $\mathcal{Q}$ , and its theory, in the language  $\mathcal{L}_{\sqrt{2}}$ , is that of an ordered  $F$ -vector space.

To see that  $T$  is tight we note that each definable cut in  $\mathcal{Q}$  can also be written as  $x_1 + \sqrt{2}x_2$ , for  $x_1, x_2 \in \mathcal{Q}$ , and that this is a definable family in  $T$ .  $\square$

Note that the above construction worked because of the algebraicity of  $\sqrt{2}$ . If we consider  $\mathbb{Q}^t$ , the expansion of  $\mathbb{Q}_{v_s}$  by  $x \mapsto tx$  where  $t$  realizes a cut defining a real transcendental number we would not obtain a tight structure. See the example  $\mathbb{Q}_{v_s}^\pi$  in [4].

We now prove:

**Theorem 4.7.** *The structure  $\mathbb{Q}^{\sqrt{2}}$  is not elementarily equivalent to a reduct of an o-minimal trace.*

*Proof.* Assume towards a contradiction that there is a dense pair  $(\mathcal{R}, \mathcal{Q})$  of o-minimal expansions of groups such that  $\mathbb{Q}^{\sqrt{2}}$  is elementarily equivalent to a reduct of the trace which this pair induces on a structure  $\mathcal{Q}$ . By that we mean that there is some expansion  $\hat{\mathcal{Q}}$  of  $\langle \mathcal{Q}, <, + \rangle$  satisfying  $T$ , such that every definable set in  $\hat{\mathcal{Q}}$  is definable in the dense pair  $(\mathcal{R}, \mathcal{Q})$ . While some of these sets are already definable in the o-minimal structure  $\mathcal{Q}$  others may be the intersection with  $\mathcal{Q}^n$  of subsets of  $R^n$  that are definable in  $\mathcal{R}$  over parameters which are not in  $\mathcal{Q}$ . The order relation  $<$  and the group operation  $+$  are assumed to be definable in  $\mathcal{Q}$ .

Let us consider the predicate  $P_{\sqrt{2}}(\hat{\mathcal{Q}})$ . It is a definable set in  $(\mathcal{R}, \mathcal{Q})$ , hence by [9, Theorem2], there is a definable  $Y_{\sqrt{2}} \subseteq R^2$  in the o-minimal structure  $\mathcal{R}$  such that  $Y_{\sqrt{2}} \cap \mathcal{Q} = P_{\sqrt{2}}(\hat{\mathcal{Q}})$ . Because  $\mathcal{Q}$  is dense in  $R$  (the universe of the o-minimal structure  $\mathcal{R}$ ), it easily follows that for every  $x \in \mathcal{Q}$ , there is  $y(x) \in R$  such that

$$y(x) = \sup\{y \in \mathcal{Q} : (x, y) \in Y_{\sqrt{2}}\}.$$

By taking the closure of the graph of  $y(x)$  we obtain an  $\mathcal{R}$ -definable function, which we will denote by  $\lambda_{\sqrt{2}} : R \rightarrow R$ , which gives  $y(x)$  for every  $x \in \mathcal{Q}$ . It is not hard to see that  $\lambda_{\sqrt{2}}$  is a definable automorphism of  $\langle R, + \rangle$  satisfying  $\lambda_{\sqrt{2}} \circ \lambda_{\sqrt{2}}(x) = 2x$ .

We now consider two cases. If the function  $\lambda_{\sqrt{2}}$  is  $\emptyset$ -definable in  $\mathcal{R}$  then it comes from a definable function in the o-minimal structure  $\mathcal{Q}$ , and in particular, for every  $x \in \mathcal{Q}$ , the set  $\{y \in \mathcal{Q} : (x, y) \in P_{\sqrt{2}}\}$  has a supremum in  $\mathcal{Q}$ . This contradicts the axioms of  $T$ .

On the other hand, if  $\lambda_{\sqrt{2}}$  is not  $\emptyset$ -definable then by [8], one can define in the o-minimal structure  $\mathcal{R}$  a multiplication function  $\cdot$  on  $R^2$ , making  $\langle R, <, +, \cdot \rangle$  a real closed field, call it  $K$ . A-priori the multiplication function might not be  $\emptyset$ -definable but in that case there is a  $\emptyset$ -definable family of such multiplications all of which expand  $\langle R, + \rangle$  to a real closed field. By definable choice we may find one such multiplication function that is  $\emptyset$ -definable.

Since  $\lambda_{\sqrt{2}}$  is an  $\mathcal{R}$ -definable automorphism of the additive group of  $K$  it must be of the form  $x \mapsto c \cdot x$  for some scalar  $c \in K$ . Because  $\lambda_{\sqrt{2}} \circ \lambda_{\sqrt{2}}(x) = 2x$ , and because  $\lambda_{\sqrt{2}}$  takes positive values on  $x > 0$ , the scalar  $c$  is necessarily  $\sqrt{2}$  (in the sense of  $K$ ). In particular,  $\lambda_{\sqrt{2}}$  is  $\emptyset$ -definable in  $\mathcal{R}$ , yielding a contradiction as before.  $\square$

5. THE THEORY OF  $(\mathcal{M}^*, \mathcal{M})$ 

From now on, given a complete non-valuational theory  $T$  we will denote by  $T^*$  the theory of the associated o-minimal completion, in the language  $\mathcal{L}_\emptyset^*$  (by Corollary 3.5, the theory  $T$  indeed determines  $T^*$ ). We write  $\bar{\mathcal{M}}$  and  $\mathcal{M}^*$ , for the structure  $\bar{\mathcal{M}}_\emptyset$ , and  $\mathcal{M}_\emptyset^*$ , respectively.

While  $\mathcal{M}$  and  $\mathcal{M}_\emptyset^*$  initially have different signatures it will be convenient to treat them in the same language. We thus modify the language of  $\mathcal{M}$ .

**Lemma 5.1.** *Let  $\mathcal{M}$  be a weakly o-minimal non-valuational structure. Let  $\mathcal{M}_0$  be the reduct of  $\mathcal{M}$  generated by all  $\emptyset$ -definable closed sets. Then every  $\emptyset$ -definable set in  $\mathcal{M}$  is  $\emptyset$ -definable in  $\mathcal{M}_0$ . In particular,  $\mathcal{M}$  and  $\mathcal{M}_0$  have the same  $\emptyset$ -definable sets.*

*Proof.* This follows from the proof of Proposition 2.8.  $\square$

So from now on we will assume that  $\mathcal{M}$  is given in the signature consisting of a function symbol for  $+$ , the ordering  $<$ , and a predicate for each  $\emptyset$ -definable closed set in  $M^n$ . We let  $\mathcal{L}$  be the associated language, so we may use the same language for  $\mathcal{M}^*$ . By Proposition 4.5, the structure  $\mathcal{M}^*$  eliminates quantifiers.

We let  $\mathcal{L}^P = \mathcal{L} \cup \{P\}$ , where  $P$  is a unary predicate. We consider the  $\mathcal{L}^P$ -structure

$$\mathcal{M}^P = \langle \mathcal{M}^*, \mathcal{M} \rangle,$$

where the interpretation of  $P$  is  $M$ . As we will see, the theory of  $\mathcal{M}^P$  depends only on  $T$ . We propose the following axiomatization for this theory:

Let  $T^d$  be the  $\mathcal{L}^P$ -language axiomatized as follows (we write  $(\mathcal{M}', \mathcal{M})$  for models of  $T^d$ ),

- (1)  $\mathcal{M} \models T$ ,  $\mathcal{M}' \models T^*$ .
- (2)  $M$  dense in  $M'$ .
- (3) Every definable cut in  $\mathcal{M}$  has a supremum in  $M'$ .
- (4) (when  $T$  is tight) Every element of  $\mathcal{M}'$  realizes a definable cut in  $\mathcal{M}$ .

Our goal is to prove:

**Theorem 5.2.** *The theory  $T^d$  is complete.*

**5.1. The tight case.** Assume that  $T$  is tight. As we saw in Proposition 4.2, the structure  $\mathcal{M}^*$  is interpretable in  $\mathcal{M}$  without parameters. Using axiom (4) above we immediately conclude:

**Lemma 5.3.** *Assume that  $T$  is tight.*

- (1) *If  $(\mathcal{M}', \mathcal{M}) \models T^d$  then necessarily  $\mathcal{M}' = \mathcal{M}^*$ .*
- (2) *For all  $\mathcal{M}, \mathcal{N} \models T$ , we have  $(\mathcal{N}^*, \mathcal{N}) \equiv (\mathcal{M}^*, \mathcal{M})$ .*

**5.2. The general case.**

**Theorem 5.4.** *If  $\mathcal{M}^d = (\mathcal{M}', \mathcal{M})$  and  $\mathcal{N}^d = (\mathcal{N}', \mathcal{N})$  are models of  $T^d$ , then  $\mathcal{M}^d \equiv \mathcal{N}^d$ .*

*Proof.* We may assume that  $T$  is non-tight. We may assume that  $\mathcal{M}^d$  and  $\mathcal{N}^d$  are  $\kappa$ -saturated for sufficiently large  $\kappa$ .

Notice that every  $\mathcal{M}$ -definable cut is realized in  $\mathcal{M}'$  exactly once, hence there is a natural embedding of  $\mathcal{M}^*$  into  $\mathcal{M}'$ , and the same holds for  $\mathcal{N}'$  and  $\mathcal{N}$ . However, by saturation, unless  $\mathcal{M}$  is tight it is not the case that  $\mathcal{M}'$  equals  $\mathcal{M}^*$ , since it realizes cuts which are not definable as well. Our goal is to show that there are  $(B, A) \prec (\mathcal{M}', \mathcal{M})$  and  $(D, C) \prec (\mathcal{N}', \mathcal{N})$  which are isomorphic.

Notice first that both  $M$  and  $M' \setminus M$  are dense in  $M'$ , for  $i = 1, 2$ . Indeed, this follows from the fact that  $T$  is non-valuational, so if  $c \in \bar{M} \setminus M$  is any element then  $c + M \subseteq \bar{M}$  is dense in  $\bar{M}$ , so also in  $M'$ .

Since  $\mathcal{M}^* \models T^*$  and  $\mathcal{M}^*$  eliminates quantifiers, the pair  $(\mathcal{M}', \mathcal{M}^*)$  is an elementary dense pair of o-minimal structures, so we shall apply to it the theory of dense pairs as in [9].

We first need:

**Lemma 5.5.** *Let  $(\mathcal{M}', \mathcal{M}) \models T^d$ . Let  $M_0 \prec \mathcal{M}$ . Then  $\text{dcl}_{\mathcal{M}_0^*}(M_0) = \bar{M}_0$ . Moreover,  $\text{dcl}_{\mathcal{M}'}(M_0) = \bar{M}_0$ .*

*Proof.* It will suffice to prove the first part of the lemma as the second part follows from the fact that  $\mathcal{M}_0^* \prec \mathcal{M}'$ .

First we show the right-to-left inclusion. For that we need:

**Claim 5.6.** Assume that  $f : M_0^n \rightarrow \bar{M}_0$  is a  $\emptyset$ -definable function in  $\mathcal{M}_0$ . Then there are in  $\mathcal{M}_0$  finitely many  $\emptyset$ -definable strong cells of the form  $C_1, \dots, C_k \subseteq M_0^n$ , with  $M_0^n \subseteq \bigcup_i C_i$ , and in  $\mathcal{M}_0^*$  there are finitely many  $\emptyset$ -definable functions  $\bar{f}_i : \bar{C}_i \rightarrow \bar{M}_0$ , such that for all  $x \in C_i$ ,  $\bar{f}_i(x) = f(x)$ .

*Proof.* We decompose  $M_0^n$  into  $\emptyset$ -definable strong cells,  $C_1, \dots, C_k$ , on each of which  $f$  is strongly continuous. For each  $i$ , the graph of  $\bar{f} \upharpoonright \bar{C}_i$  is the iterative convex hull of  $\Gamma(f \upharpoonright C_i)$ , so it is  $\emptyset$ -definable in  $\mathcal{M}_0^*$ .  $\square$

Assume now that  $b \in \bar{M}_0$ , then by definition of the completion, the cut  $Y = \{x \in M_0 : x < b\}$  is definable in  $\mathcal{M}_0$ , over a tuple of parameters  $a$ . We may assume that  $Y = Y_a$  for a  $\emptyset$ -definable family of sets  $\{Y_t : t \in T\}$  and  $\emptyset$ -definable set  $T \subseteq M_0^m$ , and that we have  $b = \sup Y_a$ . It follows that there is in  $\mathcal{M}_0$  a  $\emptyset$ -definable function  $f : T \rightarrow \bar{M}_0$ , such that  $f(a) = b$ .

By the above claim, we have  $T = \bigcup C_i$  a union of  $\emptyset$ -definable strong cells in  $\mathcal{M}$ , and there are  $f_i : \bar{C}_i \rightarrow \bar{M}_0$  all  $\emptyset$ -definable in  $\mathcal{M}_0^*$ , such that

$$(5) \quad \bigwedge_{i=1}^k \forall x \in C_i \bar{f}_i(x) = f(x).$$

In particular, there is  $i \in \{1, \dots, k\}$  such that  $a \in C_i$  and  $b = \bar{f}_i(a)$  is in  $\text{dcl}_{\mathcal{M}_0^*}(M_0)$ . Thus,  $\bar{M}_0 \subseteq \text{dcl}_{\mathcal{M}_0^*}(M_0)$ .

For the converse, we assume that  $g(a) = b$  for some  $\emptyset$ -definable function  $g$  in  $\mathcal{M}_0^*$  and  $a \in M_0^m$ . We want to show that  $b \in \bar{M}_0$ , namely that  $b$  is the supremum of a definable cut in the structure  $\mathcal{M}_0$ .

The function  $g$  is  $\emptyset$ -definable in the o-minimal structure  $\mathcal{M}_0^*$ , so by Theorem 3.3, the set

$$Y = \{(x, y) \in M_0^{n+1} : y < g(x)\}$$

is  $\emptyset$ -definable in  $\mathcal{M}_0$  and we have

$$(6) \quad \forall x \in M_0^n g(x) = \sup(Y_x).$$

It follows that  $b = g(a) = \sup Y_a$ , with  $Y \subseteq M_0^{n+1}$  a  $\emptyset$ -definable set in  $\mathcal{M}_0$ . Hence,  $b \in \bar{M}_0$ .  $\square$

We will also need:

**Claim 5.7.** For  $A \subseteq M$  and  $a \in M$ , the  $\mathcal{M}$ -type of  $a$  over  $A$  is determined by the cut of  $a$  in  $\text{dcl}_{\mathcal{M}'}(A)$ .

*Proof.* Assume that  $a$  and  $b$  in  $M$  realize the same cut over  $\text{dcl}_{\mathcal{M}'}(A)$ . To see that  $a$  and  $b$  realize the same  $\mathcal{M}$ -type over  $A$ , it is sufficient, by the weak o-minimality of  $\mathcal{M}$ , to show, for every cut  $C \subseteq M$  definable in  $\mathcal{M}$  over  $A$ , that  $a \in C$  iff  $b \in C$ . Using our assumptions, it is enough to prove that the supremum of  $C$  exists in  $M'$  and belongs to  $\text{dcl}_{\mathcal{M}'}(A)$ .

If  $C$  has a supremum  $s$  in  $M$  then  $s \in \text{dcl}_{\mathcal{M}}(A) \cap M$ , and therefore (Lemma 3.6)  $s \in \text{dcl}_{\mathcal{M}^*}(A)$ . Since  $\mathcal{M}^*$  is an elementary substructure of  $\mathcal{M}'$  we have  $s \in \text{dcl}_{\mathcal{M}'}(A)$ .

If  $C$  has no supremum in  $M$  then, by definition, its supremum is realized in  $\bar{M}$ . As  $C$  is definable in  $\mathcal{M}$  over  $A$ , its closure in  $\bar{M}$  is  $\emptyset$ -definable in  $\mathcal{M}_A^*$ , so by 4.5(2) it is definable in  $\mathcal{M}_0^*$  over  $A$ . But then  $\sup C \in \text{dcl}_{\mathcal{M}^*}(A) = \text{dcl}_{\mathcal{M}'}(A)$ . This finishes the proof.  $\square$

The rest of the proof follows closely the arguments from [9]. In order to proceed we borrow the following terminology:

**Definition 5.8.** For  $B \subseteq M'$  and  $A = B \cap M$ , we say that  $(B, A)$  is *free* if  $\dim_{\mathcal{M}'}(B'/A) = \dim_{\mathcal{M}'}(B'/M)$  for every finite  $B' \subseteq B$ . Namely, every subset of  $B$  which is  $\mathcal{M}'$ -independent over  $A$  remains independent over  $M$ . We make the same definitions for subsets of  $N'$  and  $N$ .

We consider all  $(B, A) \subseteq (M', M)$  (and similarly  $(D, C)$  in  $(N', N)$ ) which satisfy:

- (i)  $B \cap M = A$ .
- (ii)  $\text{dcl}_{\mathcal{M}'}(B) = B$ .
- (iii)  $(B, A)$  is free.

We now begin the construction of the intended isomorphism. By saturation, there is  $\mathcal{M}_0 \prec \mathcal{M}$ , of cardinality smaller than  $\kappa$  that is isomorphic to some  $\mathcal{N}_0 \prec \mathcal{N}$ .

If we let  $A_0 := M_0$ ,  $B_0 := \bar{A}_0$  and  $C_0 := \bar{N}_0$ ,  $D_0 := \bar{C}_0$ . Then (i) holds. By Lemma 5.5  $\dim_{\mathcal{M}'}(B_0/A_0) = 0$ , so  $(B_0, A_0)$  is (trivially) free. Also, by this lemma,  $B_0$  is definably closed in  $\mathcal{M}'$ , so  $(B_0, A_0)$  satisfy (i),(ii),(iii). Similarly,  $(D_0, C_0)$  satisfies (i),(ii),(iii).

Our goal is to use back-and-forth and Tarski-Vaught in order to build isomorphic elementary substructures of  $(\mathcal{M}', \mathcal{M})$  and  $(\mathcal{N}', \mathcal{N})$ . Towards that goal we need to prove the following result:

**Lemma 5.9.** *Assume that  $(B, A) \subseteq (\mathcal{M}', \mathcal{M})$  and  $(D, C) \subseteq (\mathcal{N}', \mathcal{N})$  satisfy (i),(ii),(iii), and isomorphic (namely, there is an  $\mathcal{L}$ -isomorphism  $\alpha : B \rightarrow D$  sending  $A$  onto  $C$ ), with  $|A| < \kappa$ . Then, for every  $b \in M'$ , there are  $B' \subseteq M'$ ,  $A' \subseteq M$  with  $b \in B'$ , and there are  $D' \subseteq N'$ ,  $C' \subseteq N$ , such that  $(B', A')$ ,  $(D', C')$  satisfy (i),(ii),(iii), and there is an isomorphism  $\alpha' : (B', A') \rightarrow (D', C')$  extending  $\alpha$ .*

(We also have the analogous result for  $(D, C)$  and  $d \in N'$ .)

*Proof.* We divide the argument into several cases:

**Case I.**  $b \in M$ .

First, we find  $d \in N$  such that  $\alpha(\text{tp}_{\mathcal{M}'}(b/B)) = \text{tp}_{\mathcal{N}'}(d/D)$  (so by Lemma 5.7, also  $\alpha(\text{tp}_{\mathcal{M}}(b/A)) = \text{tp}_{\mathcal{N}}(d/C)$ ). Indeed, this is possible because  $N$  is dense in  $N'$  and  $\mathcal{N}'$  is  $\kappa$ -saturated. The function  $\alpha$  then extends naturally to an isomorphism  $\alpha'$  of the o-minimal structures  $B' := \text{dcl}_{\mathcal{M}'}(Bb)$  and  $D' := \text{dcl}_{\mathcal{N}'}(Dd)$ . We let  $A' = B' \cap M$  and  $C' = D' \cap N$ . In order to see that  $\alpha'$  is an isomorphism of  $(B', A')$  and  $(D', C')$  it is left to verify is that for every  $a \in B'$ ,

$$(7) \quad a \in M \Leftrightarrow \alpha'(a) \in N.$$

So, we take  $a \in \text{dcl}_{\mathcal{M}'}(Bb)$  and prove (7).

Assume first that  $a \in \text{dcl}_{\mathcal{M}'}(Ab)$ . By Lemma 5.5,  $a \in \bar{M}$ , so we have  $a \in \text{dcl}_{\mathcal{M}^*}(Bb)$ . Hence, there exists a  $\emptyset$ -definable function  $F$  of  $(n+1)$ -variables in  $\mathcal{M}^*$ , and  $e \in (\bar{M})^n$ , with  $F(b, e) = a$ . The function  $F$  is definable in  $\mathcal{M}^*$ , and, by 3.3, its restriction to  $M^{n+1}$  is  $\emptyset$ -definable in  $\mathcal{M}$  (as a function into  $\bar{M}$ ). Thus, we can definably in  $\mathcal{M}$  partition its domain into  $\emptyset$ -definable strong cells on each of which  $F$  takes either values in  $M$  or in  $\bar{M} \setminus M$ . This partition is part of the weakly o-minimal theory  $T$ , and thus holds in both  $\mathcal{M}$  and  $\mathcal{N}$ . Since  $\alpha(\text{tp}_{\mathcal{M}}(b/A)) = \text{tp}_{\mathcal{N}}(d/C)$  it follows that  $a = F(b, e) \in M$  if and only if  $\alpha'(a) = F(d, \alpha(e)) \in N$ .

Assume now that  $a \in \text{dcl}_{\mathcal{M}'}(Bb) \setminus \text{dcl}_{\mathcal{M}'}(Ab)$  (so  $\alpha'(a) \in \text{dcl}_{\mathcal{N}'}(Dd) \setminus \text{dcl}_{\mathcal{N}'}(Cd)$ ). We claim that  $a \notin M$  and  $\alpha'(a) \notin N$ .

Indeed, assume towards a contradiction that  $a \in M$ , and let  $Y \subseteq B$  be a minimal finite set which is  $\text{dcl}_{\mathcal{M}'}(Ab)$ -independent over  $Ab$  such that  $a \in \text{dcl}_{\mathcal{M}'}(YAb)$ . Because  $a \notin \text{dcl}_{\mathcal{M}'}(Ab)$  the set  $Y$  is nonempty so fix  $y_0 \in Y$ . We have  $a \in \text{dcl}_{\mathcal{M}'}(Y'y_0Ab)$ , with  $Y' = Y \setminus \{y_0\}$ , so by exchange (and minimality of  $Y'$ ),  $y_0 \in \text{dcl}_{\mathcal{M}'}(Y'Ab)$ . Because  $a, b \in M$  and  $A \subseteq M$ , it follows that  $Y$  is not independent over  $M$ , even though it is independent over  $A$ . This contradicts the fact that  $(B, A)$  was free, so  $a \notin M$ . The same argument shows that  $\alpha'(a) \notin N$ .

Thus, we showed that  $\alpha' : (B', A') \rightarrow (D', C')$  is an isomorphism. It is clear, that the pairs satisfy (i) and (ii), so we are left to see that they are free. So, we take  $Y \subseteq B'$  independent over  $A'$  and claim that it remains independent over  $M$ . Indeed, because  $b \in A'$  (since  $b \in M$ ), it must be the case that  $Y \subseteq B$ , and the result follows immediately from the freeness of  $(B, A)$  (because  $A \subseteq A'$ ). This ends Case I.

**Case II.**  $b \in \text{dcl}_{\mathcal{M}'}(BM)$ .

In this case, there is  $\bar{m} = (m_1, \dots, m_k) \in M^k$  such that  $b \in \text{dcl}_{\mathcal{M}'}(B\bar{m})$ . We first apply Case I to each  $m_i$ , and thus may assume that  $\bar{m} \subseteq B$ , and in particular may assume that  $b$  is already in  $B$ .

**Case III.**  $b \notin \text{dcl}_{\mathcal{M}'}(BM)$

Notice first that in this case  $\mathcal{M}$  (and hence also  $\mathcal{N}$ ) is not tight (since in the tight case  $M' = \bar{M} = \text{dcl}_{\mathcal{M}'}(M)$ ). We let  $B' = \text{dcl}_{\mathcal{M}'}(Bb)$  and  $A' = B' \cap M$ . Our goal is to show that  $(B', A')$  satisfies (i),(ii),(iii), so we need to show that it is free.

We first claim that  $A' = A$ . Indeed, if  $a \in \text{dcl}_{\mathcal{M}'}(Bb) \cap M$  then either  $a \in \text{dcl}_{\mathcal{M}'}(B)$ , so  $a \in A$ , or if not then by exchange,  $b \in \text{dcl}_{\mathcal{M}'}(Ba)$ , contradicting the assumption on  $b$ .

Assume now that  $Y \subseteq B'$  is independent over  $A' = A$ . If  $Y \subseteq B$  then  $Y$  is independent over  $M$ , and otherwise, we may assume that it is of the form  $Y'b$  with  $Y' \subseteq B$ . By freeness of  $(B, A)$  we have  $Y'$  independent over  $M$  and by assumption on  $b$  we may conclude that  $Y'b$  independent over  $M$ . Thus,  $(B', A')$  is indeed free.

Next, we claim that we may find in  $\mathcal{N}'$  an element  $d$  such that  $\alpha(\text{tp}_{\mathcal{M}'}(b/B)) = \text{tp}_{\mathcal{N}'}(d/D)$  and in addition  $d \notin \text{dcl}_{\mathcal{N}'}(DN)$ . It is here that we use the fact that  $\mathcal{N}$  is non-tight. We prove:

**Lemma 5.10.** *Let  $D \subseteq N'$  be of cardinality smaller than  $\kappa$ . Then for every  $\mathcal{M}'$ -type  $p(x)$  over  $B$ , there is a realization of  $\alpha(p)$  which is not in  $\text{dcl}_{\mathcal{N}'}(DN)$ .*

*Proof.* By the saturation of  $(N', N)$  it is sufficient to prove that  $X \not\subseteq \text{dcl}_{\mathcal{N}'}(DN)$  for every infinite set  $X \subseteq N'$  that is definable in  $\mathcal{N}'$  over  $D$ . For that it is clearly sufficient to show that  $X \not\subseteq \text{dcl}_{\mathcal{N}'}(D\bar{N})$ . By applying the theory of dense pairs to the pair of o-minimal structures  $(\mathcal{N}', \bar{\mathcal{N}})$ , we may conclude from [9, Lemma 4.1], that no interval in  $\mathcal{N}'$  is in the image of  $\bar{N}^n$  under an  $\mathcal{N}'$ -definable map. This is easily seen to imply the result we want.  $\square$

This ends the proof of Lemma 5.9.  $\square$

Going back to our proof of completeness of  $T^d$ , we find  $d \in N'$  with  $\alpha(\text{tp}_{\mathcal{M}'}(b/B)) = \text{tp}_{\mathcal{N}'}(d/D)$  and with  $d \notin \text{dcl}_{\mathcal{N}'}(DN)$ . We let  $D' = \text{dcl}_{\mathcal{N}'}(Dd)$  and  $C' = D' \cap N'$  (which equals  $C$ ), so as before  $(D', C')$  is free. It is left to see that the natural extension of  $\alpha$  to  $\alpha' : B' \rightarrow D'$  preserves  $M \cap B'$ . However,  $B' \cap M = A'$  so by applying what we already know to both  $\alpha$  and  $\alpha^{-1}$  we conclude that  $x \in M' \Leftrightarrow \alpha'(x) \in N'$ . This ends the proof of Theorem 5.4.  $\square$

Notice that the proof above showed that any isomorphism of weakly o-minimal structures  $M_1 \prec M$  and  $M_2 \prec N$  can be extended to an isomorphism of elementary substructures  $(B, A) \prec (M', M)$  and  $(D, C) \prec (N', N)$ . Lemma 5.9 also implies:

**Lemma 5.11.** *Assume that  $(\mathcal{M}', \mathcal{M}), (\mathcal{N}', \mathcal{N}) \models T^d$  and  $(B, A) \subseteq (\mathcal{M}', \mathcal{M}), (D, C) \subseteq (\mathcal{N}', \mathcal{N})$  satisfy (i),(ii),(iii). If  $\alpha : B \rightarrow D$  is an  $\mathcal{L}$ -isomorphism sending  $A$  to  $C$  and  $\alpha(b) = d$  for some  $b \in B^n$  then*

$$\alpha(\text{tp}_{(\mathcal{M}', \mathcal{M})}(b/\emptyset)) = \text{tp}_{(\mathcal{N}', \mathcal{N})}(d/\emptyset).$$

We can now prove analogues of several theorems from [9]. The proofs are very similar to the original ones.

**Theorem 5.12.** *Let  $\mathcal{M}^d = (\mathcal{M}', \mathcal{M})$  be a model of  $T^d$ .*

(1) *In  $\mathcal{M}^d$ , every  $\emptyset$ -definable subset of  $(M')^n$  is a boolean combination of sets defined by formulas of the form*

$$(8) \quad \exists x_1 \cdots \exists x_k \left( \bigwedge_{i=1}^k x_i \in P \ \& \ \varphi(x_1, \dots, x_k, y) \right),$$

where  $|y| = n$  and  $\varphi(x, y)$  is an  $\mathcal{L}$  formula.

- (2) *Let  $B \subseteq M'$  be such that  $(B, B \cap M)$  is free. Then every subset of  $M^k$  that is definable in  $\mathcal{M}^d$  over  $B \subseteq M'$  is of the form  $Y \cap M^k$  for some  $Y \subseteq (M')^k$  that is definable in  $\mathcal{M}'$  over  $B$ .*
- (3) *Every subset of  $M^k$  that is definable in  $\mathcal{M}^d$  over  $A_0 \subseteq M$  is definable in the structure  $\mathcal{M}$  over  $A_0$ .*
- (4) *Every subset of  $M^n$  that is definable in  $(\mathcal{M}^*, \mathcal{M})$  (here  $\mathcal{M}^*$  is the completion of  $\mathcal{M}$ ) is definable in the structure  $\mathcal{M}$ .*

*Proof.* Without loss of generality,  $(\mathcal{M}', \mathcal{M})$  is sufficiently saturated.

(1) By standard model theoretic considerations it is enough to prove the following: For any  $b, d \in (M')^k$ , assume that  $b$  satisfies a formula of the form (8) if and only if  $d$  does. Then  $b$  and  $d$  have the same type in  $\mathcal{M}^*$  over  $\emptyset$ .

Let  $r = \dim_{\mathcal{M}'}(b/M)$ . We can find  $a \subseteq M$  finite such that  $\dim_{\mathcal{M}'}(b/a) = r$ . It follows that if we let  $B = \text{dcl}_{\mathcal{M}'}(ab)$  and  $A = \text{dcl}_{\mathcal{M}'}(a)$  then  $(B, A)$  is free and  $A = B \cap M$ .

We consider the  $\mathcal{L}$ -type of  $(b, a)$  over  $\emptyset$ . Because  $b$  and  $d$  realize the same formulas of the form (8), and because of saturation we can find  $c \in M$  such that  $\text{tp}_{\mathcal{M}'}(b, a/\emptyset) = \text{tp}_{\mathcal{M}'}(d, c/\emptyset)$ . The pair  $(D, C)$ , with  $D = \text{dcl}_{\mathcal{M}'}(cd)$  and  $C = \text{dcl}_{\mathcal{M}'}(c)$  is free with  $C = D \cap M$ . Just like in the proof of Lemma 5.9, the natural  $\mathcal{L}$ -isomorphism of  $B$  and  $D$  (sending  $(b, a)$  to  $(d, c)$ ) sends  $A$  to  $C$ .

By Lemma 5.11, the  $\mathcal{L}^P$ -types of  $b$  and  $d$  in  $(\mathcal{M}', \mathcal{M})$  are the same. Thus we proved (1).

(2) By standard model theoretic arguments it is sufficient to prove: If  $b_1, b_2 \in M^k$  satisfy the same  $\mathcal{M}'$ -type over  $B$  then they satisfy the same  $\mathcal{L}^P$ -type over  $B$ . For that, let  $A = B \cap M$ . It is sufficient to show that there are  $(B_1, A_1), (B_2, A_2) \prec (\mathcal{M}', \mathcal{M})$ , with  $(B, A) \subseteq (B_i, A_i)$  and  $b_i \in B_i$  for  $i = 1, 2$ , and there is an  $\mathcal{L}$ -isomorphism between  $(B_1, A_1)$  and  $(B_2, A_2)$ , which fixes  $B$  point-wise, and sending  $b_1$  to  $b_2$ .

We are now in the setting of Case I of the proof of Lemma 5.9, with our  $b_1, b_2$  replacing  $b, d$  there. Thus, we may first find two free pairs  $(B'_1, A'_1)$  and  $(B'_2, A'_2)$  with  $B \subseteq B'_i$  and  $b_i \in B'_i$ ,  $i = 1, 2$ , and an isomorphism  $\alpha : (B'_1, A'_1) \rightarrow (B'_2, A'_2)$  extending the identity map, with  $\alpha(b_1) = b_2$ . We now proceed exactly as in the proof of Theorem 5.4 and obtain the desired  $(B_1, A_1), (B_2, A_2) \prec (\mathcal{M}', \mathcal{M})$ . Thus,  $b_1$  and  $b_2$  realize the same  $\mathcal{L}^P$  type over  $B$  and we may conclude (1).

For (3), let  $X \subseteq M^k$  be definable in  $(\mathcal{M}', \mathcal{M})$  over  $A_0 \subseteq M$ . Notice that the mere definability of  $X$  in  $\mathcal{M}$  follows immediately from (2) but we want to show that  $X$  is definable over the same  $A_0$ . For that, it is sufficient to prove that any  $a_1, a_2 \in M$  which realize the same  $\mathcal{M}$ -type over  $A_0$  realize the same  $\mathcal{L}^P$ -type over  $A_0$ .

To do that, we first find a small model  $\mathcal{M}_1 \prec \mathcal{M}$  containing  $A_0, a_1, a_2$ , and an automorphism  $\alpha$  of  $\mathcal{M}_1$  over  $A_0$ , sending  $a_1$  to  $a_2$ . As we commented previously, we may extend  $\alpha$  to an isomorphism of two structures  $(B, A), (D, C) \prec (\mathcal{M}', \mathcal{M})$ . This is clearly sufficient.

To see (4), we note that every element of  $\mathcal{M}^*$  is in  $\text{dcl}_{\mathcal{M}^*}(N)$  and hence every definable subset of  $\bar{M}^k$  in  $\mathcal{M}^*$  can be defined over  $M$ . We now apply (3).  $\square$

Note that (3) above fails if we omit the requirement that  $M_0 \subseteq M$ , since in the non-tight case, in general,  $\mathcal{M}'$  will realize cuts which are not definable in  $\mathcal{M}$  and thus their intersection with  $M$  is not definable in  $\mathcal{M}$ .

We also point out:

**Lemma 5.13.** *If  $\mathcal{M}^d = (\mathcal{M}', \mathcal{M}) \models T^d$  then it is definably complete.*

*Proof.* If  $X \subseteq M'$  is definable in  $\mathcal{M}^d$  and bounded below then the intersection of its convex hull with  $M$  is definable in  $\mathcal{M}^d$ , and thus has the form  $Y \cap M$  for some  $Y \subseteq M'$  which is definable in  $\mathcal{M}'$ . Without loss of generality,  $Y$  is also convex and thus  $\text{Inf}Y = \text{Inf}X$ . This suffices, by o-minimality of  $\mathcal{M}'$ .  $\square$

We can now conclude, using Boxall and Hieronymi, [2]:

**Theorem 5.14.** *Let  $\mathcal{M}^d = (\mathcal{M}', \mathcal{M}) \models T^d$ . If  $U \subseteq (M')^n$  is open and definable in  $\mathcal{M}^d$  then it is definable in  $\mathcal{M}'$ . More precisely, if an open  $U$  is defined in  $\mathcal{M}^d$  over  $B \subseteq M'$  such that  $(B, B \cap M)$  is free, then  $U$  is definable in  $\mathcal{M}'$  over  $B$ . In particular,  $\mathcal{M}^P$  has an o-minimal open core.*

*Proof.* This is an immediate corollary of [2, Corollary 3.2] and what we proved so far. We extract from their argument a direct proof, which is underlined by the following simple corollary of cell decomposition.

**Fact 5.15.** *If  $Y \subseteq (M')^n$  is definable in  $\mathcal{M}'$  and  $\dim Y < n$  then  $Y \cap M^n$  has empty interior in  $M^n$ .*

We now first claim that  $\text{cl}_{M'}(U)$  is definable in  $\mathcal{M}'$  over  $B$ . Indeed, by Theorem 5.12 (2), there is  $Y \subseteq (M')^n$  definable in  $\mathcal{M}'$  over  $B$  such that  $Y \cap M^n = U \cap M^n$ . By the above observation,  $\dim Y = n$ .

Since  $M^n$  is dense in  $(M')^n$ , the set  $\text{Int}(Y) \cap M^n$  is dense in the open set  $\text{Int}(Y)$ . We claim that it is also dense in  $U$ . Indeed, we know that  $Y \cap M^n = U \cap M^n$  is open in  $M^n$  and dense in  $U$ , and by o-minimality  $\dim_{\mathcal{M}'}(Y \setminus \text{Int}(Y)) < n$ . It thus follows from Fact 5.15, that  $\text{Int}(Y) \cap M^n$  is dense in  $U$ .

So,

$$\text{cl}_{M'}(U) = \text{cl}_{M'}(\text{Int}(Y) \cap M^n) = \text{cl}_{M'}(\text{Int}(Y)).$$

Because  $Y$  was definable in  $\mathcal{M}'$  over  $B$ ,  $\text{cl}_{M'}(U)$  is definable in  $\mathcal{M}'$  over  $B$ .

We thus showed that the closure of every  $\mathcal{M}^d$ -definable open set over  $B \subseteq M'$  is definable in  $\mathcal{M}'$  over  $B$ . It follows that every  $\mathcal{M}^d$ -definable continuous function  $f : (M')^n \rightarrow M$  is definable in  $\mathcal{M}'$ , over the same parameters. Indeed, the closure of the open set  $\{(x, y) \in (M')^{n+1} : y < f(x)\}$  is exactly  $\{(x, y) \in (M')^{n+1} : y \leq f(x)\}$ , from which the definability of  $f$  follows.

Finally, we show that every closed  $F \subseteq (M')^n$  set which is  $\mathcal{M}^d$ -definable over  $B \subseteq M'$  is definable in  $\mathcal{M}'$  over  $B$ . For every  $x \in M^n$  we let  $f(x) = d(x, F) = \text{Inf}\{d(x, y) : y \in F\}$ . By Lemma 5.13, this is a well defined function in  $\mathcal{M}^d$  (over  $B$ ), and since  $F$  is closed, the function  $f$  is continuous and  $F$  is its zero set. Because  $f$  is definable in  $\mathcal{M}'$  over  $B$ , so is the set  $F$ .

Since every definable set in  $\mathcal{M}'$  can be defined over some  $B \subseteq M'$  with  $(B, B \cap M)$  free, the theorem follows.  $\square$

## REFERENCES

- [1] Yerzhan Baisalov and Bruno Poizat. Paires de structures o-minimales. *J. Symbolic Logic*, 63(2):570–578, 1998.
- [2] Gareth Boxall and Philipp Hieronymi. Expansions which introduce no new open sets. *J. Symbolic Logic*, 77(1):111–121, 2012.
- [3] A. Eleftheriou, P. E. Gunaydin and P. Hieronymi. Structure theorems in tame expansions of o-minimal structures by a dense set. *ArXiv e-prints*, 2017.
- [4] P. E. Eleftheriou, A. Hasson, and G. Keren. On definable Skolem functions in weakly o-minimal non-valuational structures. *ArXiv e-prints*, August 2015.
- [5] Gil Keren. Definable compactness in weakly o-minimal structures. Master’s thesis, Ben Gurion University of the Negev, 2014.
- [6] Dugald Macpherson, David Marker, and Charles Steinhorn. Weakly o-minimal structures and real closed fields. *Trans. Amer. Math. Soc.*, 352(12):5435–5483 (electronic), 2000.
- [7] David Marker. *Model theory*, volume 217 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2002. An introduction.
- [8] Chris Miller and Sergei Starchenko. A growth dichotomy for o-minimal expansions of ordered groups. *Trans. Amer. Math. Soc.*, 350(9):3505–3521, 1998.
- [9] Lou van den Dries. Dense pairs of o-minimal structures. *Fund. Math.*, 157(1):61–78, 1998.
- [10] Roman Wencel. Weakly o-minimal nonvaluational structures. *Ann. Pure Appl. Logic*, 154(3):139–162, 2008.
- [11] Roman Wencel. On the strong cell decomposition property for weakly o-minimal structures. *MLQ Math. Log. Q.*, 59(6):452–470, 2013.

*E-mail address:* [elitzur.by@gmail.com](mailto:elitzur.by@gmail.com)

DEPARTMENT OF MATHEMATICS, BEN GURION UNIVERSITY OF THE NEGEV, BE’ER SEHVA, ISRAEL

*E-mail address:* [hassonas@math.bgu.ac.il](mailto:hassonas@math.bgu.ac.il)

*URL:* <http://www.math.bgu.ac.il/~hasson/>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAIFA, HAIFA, ISRAEL

*E-mail address:* [kobi@math.haifa.ac.il](mailto:kobi@math.haifa.ac.il)

*URL:* <http://math.haifa.ac.il/kobi/>