

Dominant tournament families *

Raphael Yuster †

Abstract

For a tournament H with h vertices, its typical density is $h!2^{-\binom{h}{2}}/aut(H)$, i.e. this is the expected density of H in a random tournament. A family \mathcal{F} of h -vertex tournaments is *dominant* if for all sufficiently large n , there exists an n -vertex tournament G such that the density of each element of \mathcal{F} in G is larger than its typical density by a constant factor. Characterizing all dominant families is challenging already for small h . Here we characterize several large dominant families for every h . In particular, we prove the following for all h sufficiently large: (i) For all tournaments H^* with at least $5 \log h$ vertices, the family of all h -vertex tournaments that contain H^* as a subgraph is dominant. (ii) The family of all h -vertex tournaments whose minimum feedback arc set size is at most $\frac{1}{2} \binom{h}{2} - h^{3/2} \sqrt{\ln h}$ is dominant. For small h , we construct a dominant family of 6 (i.e. 50% of the) tournaments on 5 vertices and dominant families of size larger than 40% for $h = 6, 7, 8, 9$. For all h , we provide an explicit construction of a dominant family which is conjectured to obtain an absolute constant fraction of the tournaments on h vertices. Some additional intriguing open problems are presented.

AMS subject classifications: 05C20, 05C35

Keywords: tournament; density

1 Introduction

All graphs in this paper are finite and simple. Our main objects of study are *tournaments*, namely orientations of the complete graph. The *density* of a tournament H with h vertices in a larger tournament G is the probability $d_H(G)$ that a randomly chosen set of h vertices of G induces a tournament that is isomorphic to H (i.e. an H -copy in G). Stated otherwise, if $c_H(G)$ denotes the number of H -copies in an n -vertex tournament G , then $d_H(G) = c_H(G)/\binom{n}{h}$.

There are several papers that consider possible densities of a given tournament in larger tournaments [4, 5, 6, 8, 7, 12, 13]. Broadly speaking, there are a few designated regimes of interest. The maximum density of H , denoted by $d_{max}(H)$ is the limsup of the sequence whose n 'th element is the maximum possible value of $d_H(G)$ ranging over n -vertex tournaments G . The maximum density

*This research was supported by the Israel Science Foundation (grant No. 1082/16).

†Department of Mathematics, University of Haifa, Haifa 3498838, Israel. Email: raphael.yuster@gmail.com

is sometimes called the *inducibility* of H [13]. Clearly $d_{max}(H) = 1$ if and only if $H = T_h$ is the transitive tournament on h vertices. Determining $d_{max}(H)$ for some H may be quite challenging; for some small H , flag algebra techniques are useful [7, 8, 13, 15]. One can similarly consider the minimum density of H denoted by $d_{min}(H)$, but of course $d_{min}(H) = 0$ unless $H = T_h$. For the latter, $d_{min}(T_h)$ is the liminf of the sequence whose n 'th element is the minimum possible value of $d_{T_h}(G)$ ranging over n -vertex tournaments G . The *typical density*, denoted by $d(H)$ is the expected density of H in a random tournament. By a random tournament we mean, as usual, the probability space of n -vertex tournaments where the direction of each edge is chosen independently and uniformly at random. Observe that $d(H)$ is independent of n and is easy to compute. The probability of a labeled random h -vertex tournament to be isomorphic to a labeled copy of H is $2^{-\binom{h}{2}}$. Hence, $d(H) = h!2^{-\binom{h}{2}}/aut(H)$ where $aut(H)$ is the size of the automorphism group of H . In particular, $d(T_h) = h!2^{-\binom{h}{2}}$. The typical density plays an important role in the study of quasi-random tournaments [4, 6, 7, 12].

By their definitions, we have that $d_{min}(H) \leq d(H) \leq d_{max}(H)$ for every H . There are a few tournaments where one of the inequalities is an equality. For the transitive tournament T_h it is well-known that $d_{min}(T_h) = d(T_h)$ (see Exercise 10.44(b) of [14]). There are a few sporadic cases where $d_{max}(H) = d(H)$. This is easily shown to hold for $H = C_3$, the directed triangle, but it is also known to hold for the tournament on 5 vertices H_5^8 of Figure 1 as proved by Coreglano et al. [7] (there called T_5^8). It is known that all tournaments on four vertices have $d_{max}(H) > d(H)$ as well as all tournaments on at least 7 vertices [4].

Let \mathcal{T}_h denote the set of all tournaments on h vertices. So on the one hand, for a given $H \in \mathcal{T}_h$ (except for the few sporadic cases where $d_{max}(H) = d(H)$ discussed above), one can construct arbitrarily large tournaments G in which $d_H(G)$ is significantly larger than the typical density $d(H)$, but certainly no such G can be universal for all elements of \mathcal{T}_h since clearly for any G we have

$$1 = \sum_{H \in \mathcal{T}_h} d(H) = \sum_{H \in \mathcal{T}_h} d_H(G).$$

So, the natural question that emerges is, to what extent can a significant subset $\mathcal{F} \subset \mathcal{T}_h$ have the property that there are arbitrarily large tournaments G that are universal for all elements of \mathcal{F} .

Definition 1.1. *A set $\mathcal{F} \subset \mathcal{T}_h$ is dominant if there exists $\beta > 0$ such that for all sufficiently large n , there exists an n -vertex tournament G for which $d_H(G) \geq (1 + \beta)d(H)$ for all $H \in \mathcal{F}$.*

Trivially, all singletons (except for the sporadic cases discussed above where $d_{max}(H) = d(H)$) are dominant, but we are of course interested with the existence of large dominant \mathcal{F} . Clearly, if one can characterize all maximal dominant \mathcal{F} then this would characterize all dominant \mathcal{F} , but at present this seems like a problem beyond our reach (we do not even have an exact formula for the number of elements of \mathcal{T}_h). A more realistic goal is to determine large \mathcal{F} that can be explicitly

characterized in the sense that the members of \mathcal{F} are exactly the ones that satisfy some natural property (namely, given a tournament H , one can deterministically check whether H satisfies the property). This is indeed what we do in this paper for a few very natural properties.

In Section 2, we prove that for all sufficiently large h , the family of tournaments whose minimum feedback arc set size is at most $\frac{1}{2}\binom{h}{2} - h^{3/2}\sqrt{\log h}$ ¹ is a dominant family. We note that this result cannot be improved by much as it is well-known that the minimum feedback arc set size of *every* h -vertex tournament is at most $\frac{1}{2}\binom{h}{2} - \Theta(h^{3/2})$ [16]. Our main tool in the proof is the notion of the bias polynomial (a notion defined in Section 2). We also prove that the subset of all h -vertex tournaments whose bias polynomial has a local minimum at 0, is dominant. We show that for some small h , this subset is of significant size. For example, for each $h = 6, 7, 8, 9$ more than 40% of the tournaments on h vertices are of this type, and half of the tournaments on 5 vertices are of this type. We conjecture that for all h , the fraction of such tournaments out of all h -vertex tournaments is at least a positive constant independent of h .

In Section 3, we prove that for all sufficiently large h , if H^* is a tournament with at least $5 \log h$ vertices, then the family of all elements of \mathcal{T}_h that contain an H^* -copy, is dominant. Again, this result cannot be improved by much as it is well-known [18] that every element of \mathcal{T}_h contains $T_{\lceil \log h \rceil}$.

In section 4, we discuss a few open problems and conjectures related to dominant families. Solving some of these problems may be challenging.

2 The bias polynomial and dominant families

2.1 The bias polynomial

We define a probability space on labeled n -vertex tournaments that generalizes the standard uniform probability space (the random tournament model). Consider tournaments with labeled vertices $[n] = \{1, \dots, n\}$ and let $p \in [0, 1]$. If $i < j$ then make (i, j) an edge with probability p (so (j, i) is an edge with probability $1 - p$) where all $\binom{n}{2}$ choices are independent. Denote the resulting probability space by $T(n, p)$ and observe that $T(n, \frac{1}{2})$ is the usual notion of a random tournament. We note that there are other models of random graphs where the probability of an edge depends on the order of vertex labels (see, e.g., [2]).

Given $G \sim T(n, p)$, define the typical density of H in G , denoted by $d(H, p)$, to be the expected density of H in G . Notice that $d(H) = d(H, \frac{1}{2})$. Using Chebyshev's inequality, it is easy to prove that $S = \{H \in \mathcal{T}_h \mid d(H, p) > d(H)\}$ is dominant (see the proof of Lemma 2.3 below). However, recall that we would like to obtain explicit constructions of large dominant sets and for this we need to pinpoint some explicit range of p that ensures that S is large. To this end, it is beneficial

¹Unless stated otherwise, all logarithms are in base 2.

to observe that $d(H, p)$ is, in fact, a polynomial in p . Indeed, each order of the vertices of H corresponds to a term in $d(H, p)$ of the form $p^k(1-p)^{\binom{h}{2}-k}$ where k is the number of edges of H pointing from a lower ordered vertex to a higher one. So, for instance, for $H = T_3$ we have that $d(T_3, p) = p^3 + (1-p)^3 + 2p^2(1-p) + 2p(1-p)^2 = 1 - p + p^2$ while for $H = C_3$ we have $d(C_3, p) = p^2(1-p) + p(1-p)^2 = p - p^2$. Since, by symmetry, $d(H, p) = d(H, 1-p)$ it is more convenient to work with the following definition.

Definition 2.1. *The bias polynomial of H is $B(H, x) = d(H, x + \frac{1}{2})$.*

The following simple lemma lists some obvious properties of the bias polynomial.

Lemma 2.2. *Let $B(H, x)$ be the bias polynomial of a tournament H with h vertices.*

1. $B(H, x)$ is an even polynomial. Equivalently, each term of $B(H, x)$ is a constant multiple of x to an even power.
2. $B(H, 0) = d(H)$, $B(T_h, \pm\frac{1}{2}) = 1$ and otherwise $B(H, \pm\frac{1}{2}) = 0$.
3. 0 is a local extremum of $B(H, x)$. It is a local minimum if and only if the coefficient of the lowest order term of $B(H, x) - d(H)$ is positive.
4. $\sum_{H \in \mathcal{T}_h} B(H, x) = 1$.

Proof. Property 1 follows since $B(H, x) = d(H, x + \frac{1}{2}) = d(H, \frac{1}{2} - x) = B(H, -x)$. Property 2 follows since $B(H, 0) = d(H, \frac{1}{2}) = d(H)$. Property 3 follows since $B(H, x)$ is an even polynomial and the condition for local minimum follows since this is the case when the derivative at zero changes sign from negative to positive. Property 4 follows from the fact that for every $0 \leq p \leq 1$, $\sum_{H \in \mathcal{T}_h} d(H, p) = 1$. \square

Let $\mathcal{F}(h, x) = \{H \in \mathcal{T}_h \mid B(H, x) > d(H)\}$. While $\mathcal{F}(h, 0) = \emptyset$ and $\mathcal{F}(h, \pm\frac{1}{2}) = \{T_h\}$, we will prove that for certain $x = x(h)$, $\mathcal{F}(h, x)$ is large. For this to be of use, we need the following.

Lemma 2.3. *For every $x \in (0, \frac{1}{2})$, $\mathcal{F}(h, x)$ is dominant.*

Proof. Fix $0 < x < \frac{1}{2}$. Let

$$\beta = \min_{H \in \mathcal{F}(h, x)} \frac{B(H, x)}{d(H)} - 1.$$

Observe that $\beta > 0$ since by the definition of $\mathcal{F}(h, x)$ we have $B(H, x) > d(H)$ for every $H \in \mathcal{F}(h, x)$. We prove that for all sufficiently large n , there is an n -vertex tournament G such that $d_H(G) \geq (1 + \beta/2)d(H)$ holds for all $H \in \mathcal{F}(h, x)$, thus obtaining that $\mathcal{F}(h, x)$ is dominant.

Let $p = x + \frac{1}{2}$ and consider $G \sim \mathcal{T}(n, p)$. Let $H \in \mathcal{F}(h, x)$ and notice that $B(H, x) = d(H, p)$ is the expected density of H in G . Recall that $c_H(G)$ denotes the number of H -copies in G . So,

the expected value of $c_H(G)$ is $\binom{n}{h}B(H, x) = \Theta(n^h)$. We may consider each h -set of vertices of G as an indicator random variable for the event that the corresponding h -set induces a copy of H , thus $c_H(G)$ is the sum of these $\binom{n}{h}$ variables, each with success probability $B(H, x)$. But also notice that two indicator variables corresponding to disjoint h -sets are independent. Hence, the variance of $c_H(G)$ is only $O(n^{2h-1})$. By the second moment method (see [3]), the probability that $c_H(G)$ is smaller than its expected value by more than $\binom{n}{h}\frac{\beta}{2}d(H)$ is $O(n^{-1})$. Since n is chosen sufficiently large, we may assume that $n \gg |\mathcal{F}(h, x)|$. Hence there exists an n -vertex tournament G such that for all $H \in \mathcal{F}(h, x)$ it holds that $c_H(G) \geq \binom{n}{h}B(H, x) - \binom{n}{h}\frac{\beta}{2}d(H)$ and equivalently $d_H(G) \geq B(H, x) - \frac{\beta}{2}d(H)$. Finally, notice that by the definition of β ,

$$d_H(G) \geq B(H, x) - \frac{\beta}{2}d(H) \geq d(H) + \frac{\beta}{2}d(H) = \left(1 + \frac{\beta}{2}\right)d(H).$$

□

2.2 Minimum feedback arc set and dominant families

For a tournament H , a *feedback arc set* of H is a set of edges covering every directed cycle. Equivalently, it is a spanning subgraph of H whose complement is acyclic. Let $a(H)$ denote the cardinality of a smallest feedback arc set of H . While it is straightforward that $a(H) \leq \frac{1}{2}\binom{h}{2}$ and that $a(H) = 0$ if and only if $H = T_h$, determining the precise value is NP-Hard in general [1]. Spencer [16], improving earlier results of Erdős and Moon [10], proved that $a(H) \leq \frac{1}{2}\binom{h}{2} - \Theta(h^{3/2})$. We will prove that the set of all tournaments whose $a(H)$ value is slightly below this upper bound is dominant.

Let $\mathcal{A}(h, t)$ denote the set of all tournaments having $a(H) \leq \frac{1}{2}\binom{h}{2} - t$.

Theorem 2.4. $\mathcal{A}(h, h^{3/2}\sqrt{\ln h})$ is dominant for all $h \geq 30$.

Proof. We will prove that for all $h \geq 30$ it holds that $\mathcal{A}(h, h^{3/2}\sqrt{\ln h}) \subseteq \mathcal{F}(h, (\ln h/h)^{1/2})$ and hence the result will follow by Lemma 2.3. Let $x = (\ln h/h)^{1/2}$ and let $H \in \mathcal{A}(h, h^{3/2}\sqrt{\ln h})$. We must prove that $H \in \mathcal{F}(h, x)$, namely that $B(H, x) > d(H)$. Recalling that $d(H) = h!2^{-\binom{h}{2}}/aut(H)$, we must prove that $B(H, x) > h!2^{-\binom{h}{2}}/aut(H)$.

Let the vertices of H be labeled with $[h] = \{1, \dots, h\}$. For a permutation $\pi \in S_h$, let $f(\pi)$ (the “forward” edges) denote the number of edges (u, v) of H with $\pi(u) < \pi(v)$ and let $b(\pi) = \binom{h}{2} - f(\pi)$ be the “backward” edges. Then we have that

$$B(H, x) = \frac{1}{aut(H)} \sum_{\pi \in S_h} \left(\frac{1}{2} + x\right)^{f(\pi)} \left(\frac{1}{2} - x\right)^{b(\pi)}. \quad (1)$$

So it suffices to prove that

$$\sum_{\pi \in S_h} \left(\frac{1}{2} + x\right)^{f(\pi)} \left(\frac{1}{2} - x\right)^{b(\pi)} > h! 2^{-\binom{h}{2}}.$$

There are $h!$ terms on the left-hand side of the last inequality but some (in fact, most) of them are smaller than $2^{-\binom{h}{2}}$ as it is likely that for many permutations π it holds that $f(\pi)$ and $b(\pi)$ are very close, or $b(\pi)$ is larger than $f(\pi)$. But, on the other hand, we do know that for some permutation, $f(\pi)$ is considerably larger than $b(\pi)$. Indeed, since $H \in \mathcal{A}(h, h^{3/2}\sqrt{\ln h})$, there is a minimum feedback arc set of H of size at most $\frac{1}{2}\binom{h}{2} - h^{3/2}\sqrt{\ln h}$. But recall that this means that there is an acyclic spanning subgraph of H with at least $\frac{1}{2}\binom{h}{2} + h^{3/2}\sqrt{\ln h}$ edges. As each acyclic digraph has an ordering π of its vertices where all edges of the digraph are forward, we have that there exists π_0 such that $f(\pi_0) \geq \frac{1}{2}\binom{h}{2} + h^{3/2}\sqrt{\ln h}$ and consequently $b(\pi_0) \leq \frac{1}{2}\binom{h}{2} - h^{3/2}\sqrt{\ln h}$. It therefore suffices to prove that

$$\left(\frac{1}{2} + x\right)^{f(\pi_0)} \left(\frac{1}{2} - x\right)^{b(\pi_0)} > h! 2^{-\binom{h}{2}}$$

or equivalently that

$$(1 + 2x)^{f(\pi_0) - b(\pi_0)} (1 - 4x^2)^{b(\pi_0)} > h!.$$

Indeed, this holds since

$$\begin{aligned} & (1 + 2x)^{f(\pi_0) - b(\pi_0)} (1 - 4x^2)^{b(\pi_0)} \\ & > \left(1 + \frac{2\sqrt{\ln h}}{\sqrt{h}}\right)^{2h^{3/2}\sqrt{\ln h}} \left(1 - \frac{4\ln h}{h}\right)^{h^2/4} \\ & > e^{-2h \ln h} e^{3h \ln h} = h^h \end{aligned}$$

where the last inequality holds for all $h \geq 30$. □

It is important to stress that $\mathcal{A}(h, h^{3/2}\sqrt{\ln h})$, while large, is *not* a constant proportion of the family \mathcal{T}_h , as proved by Spencer [17] and de la Vega [9]. But on the other hand $\mathcal{A}(h, h^{3/2}\sqrt{\ln h})$ does contain, say, quasi-random tournaments. Indeed, by one of the equivalent notions of quasi-random tournaments proved by Chung and Graham [6], there is a quasi-random sequence of tournaments $\{H_h\}$ where H_h has h vertices such that $H_h \in \mathcal{A}(h, h^{3/2}\sqrt{\ln h})$.

tournament	bias polynomial	in \mathcal{B}_h
T_4	$\frac{3}{8} + 2x^2 + 2x^4$	\checkmark
C_4	$\frac{3}{8} - 2x^2 + 2x^4$	
D	$\frac{1}{8} - 2x^4$	
D^t	$\frac{1}{8} - 2x^4$	

Table 1: Tournaments on four vertices and their bias polynomials.




Figure 1: The tournaments on 5 vertices.

2.3 The bias subset

Property 3 of Lemma 2.2 states that we can partition \mathcal{T}_h into two subsets: those tournaments H for which 0 is a local minimum of $B(H, x)$ and those for which 0 is a local maximum of $B(H, x)$.

Definition 2.5. *The bias subset $\mathcal{B}_h \subset \mathcal{T}_h$ consists of the tournaments $H \in \mathcal{T}_h$ for which 0 is a local minimum of $B(H, x)$.*

For example, it is easy to verify that $B(T_3, x) = \frac{3}{4} + x^2$ while $B(C_3, x) = \frac{1}{4} - x^2$. Hence, $\mathcal{B}_3 = \{T_3\}$. The following is a corollary of Lemma 2.3.

Proposition 2.6. *\mathcal{B}_h is dominant.*

Proof. For each $H \in \mathcal{B}_h$, let $\alpha_H > 0$ be the largest real such that $B(H, x)$ is monotone increasing in $(0, \alpha_H)$. Such an interval exists since 0 is a local minimum of $B(H, x)$. Notice that if $H \neq T_H$ then it must be that $0 < \alpha_H \leq \frac{1}{2}$ since by Lemma 2.2, $B(H, \frac{1}{2}) = 0$ and $B(H, 0) = d(H) > 0$. If $H = T_h$ then it may be that $\alpha_H > \frac{1}{2}$ (in fact, it may be infinity) so if this occurs, just redefine

tournament	bias polynomial	in \mathcal{B}_h
T_5	$\frac{15}{128} + \frac{25}{16}x^2 + 6x^4 + 7x^6 + 2x^8$	✓
H_5^2	$\frac{5}{128} + \frac{5}{16}x^2 - \frac{1}{2}x^4 - 5x^6 - 2x^8$	✓
H_5^3	$\frac{15}{128} + \frac{5}{16}x^2 - 4x^4 + 3x^6 + 2x^8$	✓
H_5^4	$\frac{5}{128} + \frac{5}{16}x^2 - \frac{1}{2}x^4 - 5x^6 - 2x^8$	✓
H_5^5	$\frac{15}{128} - \frac{5}{16}x^2 + \frac{1}{2}x^4 - 3x^6 - 6x^8$	
H_5^6	$\frac{15}{128} + \frac{5}{16}x^2 - 4x^4 + 3x^6 + 2x^8$	✓
H_5^7	$\frac{5}{128} + \frac{5}{16}x^2 - \frac{1}{2}x^4 - 5x^6 - 2x^8$	✓
H_5^8	$\frac{15}{128} - \frac{5}{16}x^2 - \frac{5}{2}x^4 + 5x^6 + 10x^8$	
H_5^9	$\frac{15}{128} - \frac{15}{16}x^2 + 2x^4 - x^6 + 2x^8$	
H_5^{10}	$\frac{5}{128} - \frac{5}{16}x^2 + x^4 - 3x^6 + 6x^8$	
H_5^{11}	$\frac{15}{128} - \frac{15}{16}x^2 + x^4 + 7x^6 - 14x^8$	
H_5^{12}	$\frac{3}{128} - \frac{5}{16}x^2 + \frac{3}{2}x^4 - 3x^6 + 2x^8$	

Table 2: Tournaments on five vertices and their bias polynomials.

$\alpha_{T_H} = \frac{1}{2}$. Now define $\alpha_h = \min\{\frac{1}{2}\alpha_H \mid H \in \mathcal{B}_h\}$. As α_h is a minimum of a finite set of positive reals, each no larger than $\frac{1}{4}$, we have that $0 < \alpha_h \leq \frac{1}{4}$. By Lemma 2.3, $\mathcal{F}(h, \alpha_h)$ is dominant. As $\mathcal{B}_h \subseteq \mathcal{F}(h, \alpha_h)$, the proposition follows. \square

Note that \mathcal{B}_h is explicitly constructed, as for each tournament H one merely needs to compute the bias polynomial $B(H, x)$ as given in (1) and check whether the coefficient of the lowest order term of $B(H, x) - d(H) = B(H, x) - B(H, 0)$ is positive. In Tables 1 and 2 we list the bias polynomials of \mathcal{T}_4 and \mathcal{T}_5 respectively. In particular, we obtain that $\mathcal{B}_5 = \{T_5, H_5^2, H_5^3, H_5^4, H_5^6, H_5^7\}$ which is half of the total of 12 tournaments on 5 vertices. In Table 3 we list for all $3 \leq h \leq 9$ the size of \mathcal{B}_h and the ratio of \mathcal{B}_h and \mathcal{T}_h . In particular, we have that $|\mathcal{B}_9| = 79229$ which constitutes more than 41% of the total number of tournaments on 9 vertices². The following conjecture, if true, will give a dominant subset that is at least an absolute constant fraction of \mathcal{T}_h .

Conjecture 2.7. *There exists an absolute constant $c > 0$ such that for all $h \geq 3$, $|\mathcal{B}_h| \geq c|\mathcal{T}_h|$.*

²Source code of our program is available at <https://www.dropbox.com/s/y9zovepfr1hg1nt/dominant-tour.zip?dl=0>

h	3	4	5	6	7	8	9
$ \mathcal{T}_h $	2	4	12	56	456	6880	191536
$ \mathcal{B}_h $	1	1	6	25	199	2769	79229
$ \mathcal{B}_h / \mathcal{T}_h $	0.5	0.25	0.5	0.446...	0.436...	0.402...	0.413...

Table 3: The sizes of \mathcal{T}_h and \mathcal{B}_h and their ratio, for small h .

3 Tournaments with a common subgraph

For a tournament H^* with at most h vertices, let $\mathcal{T}_h(H^*)$ denote the set of all elements of \mathcal{T}_h that contain H^* as a sub-tournament. Our main result in this section follows.

Theorem 3.1. *For all sufficiently large h , if H^* contain at least $5 \log h$ vertices then $\mathcal{T}_h(H^*)$ is dominant.*

Proof. We assume that h is sufficiently large and that the number of vertices of H^* is k where $h > k \geq 5 \log h$. We define a probability space of n -vertex tournaments (hereafter we assume that n is a multiple of h , as this assumption does not affect the theorem's statement). Assume that the vertices of H^* are labeled with $[k]$. Consider vertex set $[n]$ partitioned into $k + 1$ subsets V_1, \dots, V_{k+1} . For $i = 1, \dots, k$, set V_i has n/h vertices and set V_{k+1} consists of the remaining $n - kn/h$ vertices. For all $1 \leq i < j \leq k$, the edges between V_i and V_j are all directed from V_i to V_j if $(i, j) \in E(H^*)$ or all directed from V_j to V_i if $(j, i) \in E(H^*)$. Observe that each transversal of V_1, \dots, V_k induces a copy of H^* . The remaining edges, which in particular include the edges having at least one endpoint in V_{k+1} , are oriented randomly, uniformly and independently. Denote the resulting probability space by $T(n, h, H^*)$. We prove that for a small positive $\beta = \beta(h)$ it holds that for each $H \in \mathcal{T}_h(H^*)$, its expected density in $G \sim T(n, h, H^*)$ is at least $(1 + \beta)d(H)$. By the second moment method, exactly as in the proof of Lemma 2.3, this implies that $\mathcal{T}_h(H^*)$ is dominant.

Let, therefore, $H \in \mathcal{T}_h(H^*)$ be labeled with vertex set $[h]$ such that the sub-tournament of H induced by $[k]$ is label-isomorphic to H^* . Recall that $d(H) = h!2^{-\binom{h}{2}}/aut(H)$. Let P denote the set of all $(h - k)!$ permutations of $[h]$ that are stationary on $[k]$, let $Aut(H)$ denote the automorphism group of H and let $Q \leq Aut(H) \cap P$ be the sub-group of $Aut(H)$ consisting of the permutations of $[h]$ that are stationary on $[k]$. Observe that $1 \leq |Q| \leq aut(H)$.

Suppose now that $G \sim T(n, h, H^*)$. Consider a random injection f from $[h]$ to $[n]$. We call f good if $f(i) \in V_i$ for $i = 1, \dots, k$ and $f(i) \in V_{k+1}$ for $i = k + 1, \dots, h$. By the sizes of the V_i 's we

have that f is good with probability

$$\frac{1}{h^k} \prod_{i=k+1}^h \left(\frac{n - kn/h - i + k + 1}{n - i + 1} \right) \geq \frac{1}{(eh)^k}.$$

Given that f is good, the probability that its image induces a copy of H is

$$\frac{(h-k)!}{|Q|} 2^{\binom{k}{2} - \binom{h}{2}}$$

since $\binom{h}{2} - \binom{k}{2}$ is the number of edges with an endpoint in V_{k+1} . Hence, $\mathbb{E}[d_H(G)]$ (the expectation of $d_H(G)$) satisfies

$$\begin{aligned} \mathbb{E}[d_H(G)] &\geq \frac{1}{(eh)^k} \cdot \frac{(h-k)!}{|Q|} 2^{\binom{k}{2} - \binom{h}{2}} \\ &\geq \frac{1}{(eh^2)^k} \cdot \frac{h! 2^{-\binom{h}{2}}}{\text{aut}(H)} 2^{\binom{k}{2}} \\ &= d(H) \frac{1}{(eh^2)^k} 2^{\binom{k}{2}} \end{aligned}$$

So, to prove the existence of $\beta = \beta(h)$ it suffices to prove that $2^{(k-1)/2} > eh^2$. Indeed this holds as $k \geq 5 \log h$ and because h is sufficiently large. \square

4 Concluding remarks and some open problems

We list a few open problems and conjectures concerning dominant families. An h -vertex tournament H is *highly dominant* if every maximal dominant subset of \mathcal{T}_h contains H . The proposition shows that there are highly dominant tournaments.

Proposition 4.1. *T_h is highly dominant for all $h \geq 3$.*

Proof. Let $\mathcal{F} \subset \mathcal{T}_h$ be dominant. Hence, there exists $\beta = \beta(\mathcal{F})$ and $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$, there exists a tournament G with n vertices such that $d_H(G) \geq (1 + \beta)d(H)$ for each $H \in \mathcal{F}$. For $n \geq n_0$ let G_n be tournament satisfying $d_H(G_n) \geq (1 + \beta)d(H)$ for each $H \in \mathcal{F}$. Fix some $H \in \mathcal{F}$. As $d_H(G_n) \geq (1 + \beta)d(H)$ for all $n \geq n_0$, it follows from the result of Chung and Graham [6] that $\{G_n\}$ is *not* a quasi-random sequence, as it violates property $P_1(h)$ there. But on the other hand, it follows from exercise 10.44(b) of [14] and also from [8] that T_h is quasi-random forcing, implying that for our sequence, there exists $\epsilon > 0$ and $n_1 \geq n_0$ such that for all $n \geq n_1$, $d_{T_h}(G_n) \geq (1 + \epsilon)d(T_h)$. This implies that $\{T_h\} \cup \mathcal{F}$ is dominant. \square

Problem 4.2. *Determine all highly dominant tournaments. In particular, are there non-transitive highly dominant tournaments?*

It is very easy to show that for every positive integer $k \geq 2$, there is a minimum integer $f(k)$ such that for all $h \geq f(k)$, every k -subset of \mathcal{T}_h is dominant. The following proposition gives an upper bound for $f(k)$.

Proposition 4.3. $f(k) \leq (1 + o_k(1)) \log k$.

Proof. Fix $\epsilon > 0$ and assume throughout the proof that k is sufficiently large. Let $h \geq (1 + \epsilon) \log k$. Let $r = h \lceil \sqrt{hk} \rceil$. Consider a complete graph M on r vertices. Take r/h pairwise vertex-disjoint copies of K_h (namely, a K_h -factor of M), remove the edges of this factor from M and repeat taking factors. After taking t factors we have already taken tr/h pairwise edge-disjoint copies of K_h and the spanning subgraph of M consisting of the edges not yet taken is regular of degree $r - 1 - t(h - 1)$. By the Hajnal-Szemerédi Theorem [11] we can do so as long as $r - 1 - t(h - 1) \geq r - r/h$ so we can have $t \geq r/h^2$. Thus, we can find in M at least $r^2/h^3 \geq k$ pairwise edge-disjoint copies of K_h . Now suppose that the vertices of M are $[r]$ and that a set of k pairwise edge-disjoint copies of K_h in M is $\mathcal{R} = \{X_1, \dots, X_k\}$ and $V(X_i) = \{x_{i,1}, \dots, x_{i,h}\}$.

Now suppose that $\mathcal{F} = \{H_1, \dots, H_k\} \subset \mathcal{T}_h$. We must prove that \mathcal{F} is dominant. We assume that the vertices of each H_i are labeled with $[h]$. Suppose that n is an integer multiple of r . Consider vertex sets V_1, \dots, V_r each of size n/r . We construct a random tournament with n vertices as follows. For each $i = 1, \dots, k$, and for each pair j, j' of distinct indices from $[h]$, we orient all edges from $V_{x_{i,j}}$ to $V_{x_{i,j'}}$ if $(j, j') \in E(H_i)$ else we orient all edges from $V_{x_{i,j'}}$ to $V_{x_{i,j}}$ if $(j', j) \in E(H_i)$. Notice that the orientations are well-defined as the elements of \mathcal{R} are pairwise edge-disjoint. The remaining edge of G (those having two endpoints in the same part V_i or those between V_i and V_j where i, j are not both in some element of \mathcal{R}) are oriented arbitrarily.

Fix some $H_i \in \mathcal{F}$. Then, $d_H(G)$ is at least the probability that a randomly chosen h -set of G is a transversal of $V_{x_{i,1}}, \dots, V_{x_{i,h}}$, as any such transversal induces a copy of H_i in G . But the probability that a randomly chosen h -set of G is such is at least $h!/r^h$, so $d_H(G) \geq h!/r^h$. It therefore remains to prove that

$$\frac{h!}{r^h} > d(H) = \frac{h! 2^{-\binom{h}{2}}}{\text{aut}(H)}$$

so it suffices to prove that $r^h < 2^{\binom{h}{2}}$ or, equivalently, $2r^2 < 2^h$. Indeed, this holds since $r = h \lceil \sqrt{hk} \rceil$ and since $h \geq (1 + \epsilon) \log k$. \square

Problem 4.4. *Determine some small values of $f(k)$. In particular, determine $f(2)$.*

Let $g(h)$ denote the maximum size of a dominant subset of \mathcal{T}_h . Of course, we do not expect to obtain an exact formula for $g(h)$, as there is no such exact formula for $|\mathcal{T}_h|$. But perhaps good asymptotic values could be obtained.

Problem 4.5. *Provide good estimates for $g(h)$.*

References

- [1] Noga Alon. Ranking tournaments. *SIAM Journal on Discrete Mathematics*, 20(1):137–142, 2006.
- [2] Noga Alon, Michael Krivelevich, and Benny Sudakov. Large nearly regular induced subgraphs. *SIAM Journal on Discrete Mathematics*, 22(4):1325–1337, 2008.
- [3] Noga Alon and Joel Spencer. *The probabilistic method*. John Wiley & Sons, 2004.
- [4] Matija Bucić, Eoin Long, Asaf Shapira, and Benny Sudakov. Tournament quasirandomness from local counting. arXiv 1910.09936, 2019.
- [5] Timothy F. N. Chan, Andrzej Grzesik, Daniel Král, and Jonathan A. Noel. Cycles of length three and four in tournaments. arXiv 1902.00572, 2019.
- [6] Fan R. K. Chung and Ronald L. Graham. Quasi-random tournaments. *Journal of Graph Theory*, 15(2):173–198, 1991.
- [7] Leonardo N. Coregliano, Roberto F. Parente, and Cristiane M. Sato. On the maximum density of fixed strongly connected subtournaments. *Electronic Journal of Combinatorics*, 26(1):P44, 2019.
- [8] Leonardo N. Coregliano and Alexander A. Razborov. On the density of transitive tournaments. *Journal of Graph Theory*, 85(1):12–21, 2017.
- [9] W. Fernandez de la Vega. On the maximum cardinality of a consistent set of arcs in a random tournament. *Journal of Combinatorial Theory, Series B*, 35(3):328–332, 1983.
- [10] Paul Erdős and J. W. Moon. On sets of consistent arcs in a tournament. *Canadian Mathematical Bulletin*, 8:269–271, 1965.
- [11] A. Hajnal and E. Szemerédi. Proof of a conjecture of Erdős. In *Combinatorial theory and its applications, II (Proc. Colloq., Balatonfüred, 1969)*, pages 601–623. North-Holland, Amsterdam, 1970.
- [12] Robert Hancock, Adam Kabela, Daniel Král, Taísa Martins, Roberto Parente, Fiona Skerman, and Jan Volec. No additional tournaments are quasirandom-forcing. arXiv 1912.04243, 2019.
- [13] Nati Linial and Avraham Morgenstern. On the number of 4-cycles in a tournament. *Journal of Graph Theory*, 83(3):266–276, 2016.

- [14] László Lovász. *Combinatorial Problems and Exercises*, volume 361. American Mathematical Society, 2007.
- [15] Alexander A. Razborov. Flag algebras. *The Journal of Symbolic Logic*, 72(4):1239–1282, 2007.
- [16] Joel Spencer. Optimal ranking of tournaments. *Networks*, 1(2):135–138, 1971.
- [17] Joel Spencer. Optimally ranking unrankable tournaments. *Periodica Mathematica Hungarica*, 11(2):131–144, 1980.
- [18] Richard Stearns. The voting problem. *The American Mathematical Monthly*, 66(9):761–763, 1959.