

Vector clique decompositions

Raphael Yuster *

Abstract

Let \mathcal{F}_k be the set of all graphs on k vertices. For a graph G , a k -decomposition is a set of induced subgraphs of G , each of which is isomorphic to an element of \mathcal{F}_k , such that each pair of vertices of G is in exactly one element of the set. It is a fundamental result of Wilson that for all $n = |V(G)|$ sufficiently large, G has a k -decomposition if and only if G is k -divisible, namely $k - 1$ divides $n - 1$ and $\binom{k}{2}$ divides $\binom{n}{2}$.

Let $\mathbf{v} \in \mathbb{R}^{|\mathcal{F}_k|}$ be indexed by \mathcal{F}_k . For a k -decomposition L of G , let $\nu_{\mathbf{v}}(L) = \sum_{F \in \mathcal{F}_k} \mathbf{v}_F d_{L,F}$ where $d_{L,F}$ is the fraction of elements of L that are isomorphic to F . Let $\nu_{\mathbf{v}}(G) = \max_L \nu_{\mathbf{v}}(L)$ and $\nu_{\mathbf{v}}(n) = \min\{\nu_{\mathbf{v}}(G) : |V(G)| = n\}$ ¹. It is not difficult to prove that the sequence $\nu_{\mathbf{v}}(n)$ has a limit so let $\nu_{\mathbf{v}} = \lim_{n \rightarrow \infty} \nu_{\mathbf{v}}(n)$. Replacing k -decompositions with their fractional relaxations, one obtains the (polynomial time computable) fractional analogue $\nu_{\mathbf{v}}^*(G)$ and the corresponding fractional values $\nu_{\mathbf{v}}^*(n)$ and $\nu_{\mathbf{v}}^*$. Our first main result is that for each $\mathbf{v} \in \mathbb{R}^{|\mathcal{F}_k|}$

$$\nu_{\mathbf{v}} = \nu_{\mathbf{v}}^* .$$

Furthermore, there is a polynomial time algorithm that produces a decomposition L of a k -decomposable graph such that $\nu_{\mathbf{v}}(L) \geq \nu_{\mathbf{v}} - o_n(1)$.

A similar result holds when \mathcal{F}_k is the family of all tournaments on k vertices and when \mathcal{F}_k is the family of all edge-colorings of K_k .

We use these results to obtain new and improved bounds on several decomposition results. For example, we prove that every n -vertex tournament which is 3-divisible (namely $n = 1, 3 \pmod{6}$) has a triangle decomposition in which the number of directed triangles is less than $0.0222n^2(1 + o(1))$ and that every 5-decomposable n -vertex graph has a 5-decomposition in which the fraction of cycles of length 5 is $o_n(1)$.

MSC codes: 05C70, 05C35

1 Introduction

The problem of decomposing a large graph G into pairwise edge-disjoint copies of a given graph F has been extensively studied and dates back to a result of Kirkman from 1847 [13], who proved

*Department of Mathematics, University of Haifa, Haifa 31905, Israel. Email: raphy@math.haifa.ac.il. This research was supported by the Israel Science Foundation (grant No. 1082/16).

¹If n is not such that graphs on n vertices have a k -decomposition, one can synthetically define $\nu_{\mathbf{v}}(n) = \nu_{\mathbf{v}}(m)$ where $m < n$ is the largest integer such that graphs on m vertices are k -decomposable.

that K_n has a K_3 -decomposition whenever $n \equiv 1, 3 \pmod{6}$. These divisibility requirements are necessary as in any decomposition of a graph into triangles, the degree of each vertex must be even and the number of edges must be divisible by 3.

More generally, for a graph G to have an F -decomposition, it must trivially hold that the gcd of the degree sequence of G , denoted by $\gcd(G)$ is divisible by $\gcd(F)$ and that the number of edges of G , denoted by $e(G)$, is divisible by $e(F)$. We therefore say that G is F -divisible if these two necessary conditions hold.

The F -decomposition problem for $G = K_n$ was completely solved (for large n) by Wilson [16, 17, 18, 19]. He proved that whenever n is sufficiently large and K_n is F -divisible (which simply means that $n - 1$ is divisible by $\gcd(F)$ and $\binom{n}{2}$ is divisible by $e(F)$), then it has an F -decomposition. Recently, this result has been generalized by Keevash [11] to the complete uniform hypergraph setting [11]. See also Glock et al. [6] for another proof.

Another equivalent way to state Wilson's Theorem is the following. Suppose \mathcal{F} is the set of all spanning subgraphs of F . An \mathcal{F} -decomposition of a graph G is a set of subgraphs of G , each of which is isomorphic to an element of \mathcal{F} , such that each pair of vertices of G is in exactly one element of the set (and if this pair is an edge of G , then it is also an edge of that element). So, Wilson's theorem asserts that for $n \geq n_0(F)$, if K_n is F -divisible, then any graph G with n vertices has an \mathcal{F} -decomposition. This equivalent statement, leads, however, to a wider set of questions as clearly, if G has an \mathcal{F} -decomposition, it has many (in particular, since any vertex permutation may lead to a distinct \mathcal{F} -decomposition). Thus, we can ask about the quality of the various \mathcal{F} -decompositions with respect to the distribution of the members of \mathcal{F} in it.

More formally, for a vector $\mathbf{v} \in \mathbb{R}^{|\mathcal{F}|}$ indexed by \mathcal{F} and an \mathcal{F} -decomposition L of G , let $D_{\mathbf{v}}(L) = \sum_{H \in \mathcal{F}} \mathbf{v}_H |L_H|$ where L_H is the subset of L whose elements are isomorphic to H . It will be slightly more convenient to normalize this quantity by defining $d_{L,H} = |L_H|/|L|$ to be the density of H in L and defining $\nu_{\mathbf{v}}(L) = \sum_{H \in \mathcal{F}} \mathbf{v}_H d_{L,H}$, observing that $\nu_{\mathbf{v}}(L) = D_{\mathbf{v}}(L)/|L|$ and that $|L| = \binom{n}{2}/e(F)$.

The quality of the decomposition is thus measured by $\nu_{\mathbf{v}}(G) = \max_L \nu_{\mathbf{v}}(L)$ where the maximum is over all \mathcal{F} -decompositions of G . We call $\nu_{\mathbf{v}}(G)$ the *optimal* \mathcal{F} -decomposition of G with respect to \mathbf{v} . So, when K_n has an F -decomposition, $\nu_{\mathbf{v}}(G)$ is well-defined for every $\mathbf{v} \in \mathbb{R}^{|\mathcal{F}|}$ and every graph G with n vertices.

Notice that given F , we may consider additional structures underlined by F other than the spanning subgraphs of F . For instance, we may define \mathcal{F} to be all possible orientations of F and then an \mathcal{F} -decomposition is defined for *tournaments* G and $\nu_{\mathbf{v}}(G)$ is defined analogously. Likewise, we may define \mathcal{F} to be all edge colorings of F with colors from a given set of colors. In this case, an \mathcal{F} -decomposition is defined for edge colorings G of K_n and $\nu_{\mathbf{v}}(G)$ is defined analogously.

In what follows, we will state our results for the case of $F = K_k$, although our results do carry over quite seamlessly to certain more general F . We prefer this approach as it seems to be the most interesting case (in fact, already for some questions arising in the case $k = 3$), yet it captures all details of the general proof and since all our applications involve the case where $F = K_k$. So, let \mathcal{F}_k denote the set of all graphs on k vertices, let \mathcal{T}_k denote the set of all tournaments on k vertices,

and for a color set C , let \mathcal{C}_k denote the set of all edge colorings of K_k with colors from C . Let G be an n -vertex graph. If K_n is K_k -divisible (i.e. if $k - 1$ divides $n - 1$ and $\binom{k}{2}$ divides $\binom{n}{2}$), then G is called k -divisible. Similarly, G is k -decomposable if K_n has a K_k -decomposition. In these terms, Wilson's Theorem asserts that for all n sufficiently large, G is k -decomposable if and only if it is k -divisible.

We first observe that computing $\nu_{\mathbf{v}}(G)$ is easy for some vectors, while NP-Hard for some others. Indeed, let $\mathbf{v} \in \mathbb{R}^{|\mathcal{F}_k|}$ be a constant vector, all entries equal to c . In this case, once we know that G is k -decomposable (which we can determine in polynomial time by Wilson's Theorem), we have that $\nu_{\mathbf{v}}(G) = c$ as any k -decomposition has this optimal weight. On the other hand, consider the case $k = 3$ and the vector which assigns K_3 the weight 1 and assigns the other graphs on three vertices, the weight zero. Suppose G is 3-divisible. Now, if G and its complement each have a K_3 -decomposition, then we would have $\nu_{\mathbf{v}}(G) = e(G)/\binom{n}{2}$. Otherwise, we would have $\nu_{\mathbf{v}}(G) < e(G)/\binom{n}{2}$. But determining whether a graph and its complement have a K_3 -decomposition is NP-Complete (see [3, 9])².

Another minor observation is that if $\mathbf{w} \in \mathbb{R}^{|\mathcal{F}_k|}$ is obtained from \mathbf{v} by dilation and translation with a constant vector, namely, $\mathbf{w} = c\mathbf{v} + d\mathbf{1}$ for some $c > 0$, then $\nu_{\mathbf{w}}(G) = c\nu_{\mathbf{v}}(G) + d$. For this reason, it may sometimes be convenient to assume that the smallest coordinate of \mathbf{v} is 0 and the largest coordinate is 1 (or that $\mathbf{v} = \mathbf{1}$). Notice also that by dilation and translation with a constant vector, one can transform \mathbf{v} to a nonnegative vector whose coordinate sum is 1, namely a probability distribution on \mathcal{F}_k .

Given a vector $\mathbf{v} \in \mathbb{R}^{|\mathcal{F}_k|}$, the extremal graph-theoretic question of interest is how small can $\nu_{\mathbf{v}}(G)$ be³. Thus, let $\nu_{\mathbf{v}}(n)$ denote the minimum of $\nu_{\mathbf{v}}(G)$ taken over all graphs G with n vertices such that K_n is k -decomposable. To formally extend this sequence to all n , one can synthetically define $\nu_{\mathbf{v}}(n) = \nu_{\mathbf{v}}(m)$ such that $m \leq n$ is the largest integer such that K_m is k -decomposable (trivially K_1 is k -decomposable). It is not difficult to prove that the sequence $\nu_{\mathbf{v}}(n)$ converges (as shown later in this paper), but, as noted earlier, in most cases it is difficult, and possibly intractable, to determine the limit. So, let $\nu_{\mathbf{v}} = \lim_{n \rightarrow \infty} \nu_{\mathbf{v}}(n)$.

To state our first result, we need to recall the notion of a fractional k -decomposition. Let $\binom{G}{k}$ denote the set of $\binom{n}{k}$ k -vertex induced subgraphs of an n -vertex graph G . For a pair of vertices x, y of G , let $\binom{G}{k, x, y}$ be the k -vertex induced subgraphs of G that contain both x and y . A *fractional k -decomposition* is a function $f : \binom{G}{k} \rightarrow [0, 1]$ such that for each pair of vertices x, y , $\sum_{H \in \binom{G}{k, x, y}} f(H) = 1$. Clearly, a k -decomposition is also a fractional k -decomposition whose image is $\{0, 1\}$. Observe that every graph with $n \geq k$ vertices has a fractional k -decomposition, regardless of being k -divisible. For $\mathbf{v} \in \mathbb{R}^{|\mathcal{F}_k|}$ indexed by \mathcal{F}_k and a fractional k -decomposition f

²In fact it is proved that deciding if a graph has a K_3 -decomposition is NP-Complete, but it is straightforward to reduce this problem to the problem of whether a graph and its complement each have a K_3 -decomposition.

³Maximizing $\nu_{\mathbf{v}}(G)$ is trivial. It is just the largest coordinate of \mathbf{v} , as if K_n is k -decomposable, we can replace each copy of K_k in a K_k -decomposition of K_n with a copy of H where \mathbf{v}_H is the maximum coordinate of \mathbf{v} and the obtained graph G has $\nu_{\mathbf{v}}(G) = \mathbf{v}_H$.

of G , let $D_{\mathbf{v}}(f) = \sum_{H \in \mathcal{F}_k} \mathbf{v}_H f(H)$ where $f(H)$ is the sum of the values of f on elements of $\binom{G}{k}$ that are isomorphic to H . As before, it will be slightly more convenient to consider the normalized value $\nu_{\mathbf{v}}^*(f) = D_{\mathbf{v}}(f) \binom{k}{2} / \binom{n}{2}$. We therefore define the *optimal* fractional k -decomposition of G with respect to \mathbf{v} by $\nu_{\mathbf{v}}^*(G) = \max_f \nu_{\mathbf{v}}^*(f)$ where the maximum is taken over all fractional k -decompositions of G . We define $\nu_{\mathbf{v}}^*(n)$ to be the minimum of $\nu_{\mathbf{v}}^*(G)$ taken over all graphs G with n vertices. It is easy to verify that the sequence $\nu_{\mathbf{v}}^*(n)$ is non-decreasing and is upper bounded by the largest coordinate of \mathbf{v} , thus let $\nu_{\mathbf{v}}^* = \lim_{n \rightarrow \infty} \nu_{\mathbf{v}}^*(n)$. By the previous remark, we always have $\nu_{\mathbf{v}}^*(G) \geq \nu_{\mathbf{v}}(G)$, and consequently $\nu_{\mathbf{v}}^*(n) \geq \nu_{\mathbf{v}}(n)$ and $\nu_{\mathbf{v}}^* \geq \nu_{\mathbf{v}}$. The following is our first main result. We state it also for the analogous versions of tournaments and edge-colored graphs.

Theorem 1 *Let $k \geq 3$ be a given integer.*

1. *Let $\mathbf{v} \in \mathbb{R}^{|\mathcal{F}_k|}$ be a given vector indexed by \mathcal{F}_k . Then, $\nu_{\mathbf{v}} = \nu_{\mathbf{v}}^*$.*
2. *Let $\mathbf{v} \in \mathbb{R}^{|\mathcal{T}_k|}$ be a given vector indexed by \mathcal{T}_k . Then, $\nu_{\mathbf{v}} = \nu_{\mathbf{v}}^*$.*
3. *Let C be a finite set of colors and let $\mathbf{v} \in \mathbb{R}^{|\mathcal{C}_k|}$ be a given vector indexed by \mathcal{C}_k . Then, $\nu_{\mathbf{v}} = \nu_{\mathbf{v}}^*$.*

In all cases, if G has n vertices such that K_n is k -decomposable, then a k -decomposition L of G satisfying $\nu_{\mathbf{v}}(L) \geq \nu_{\mathbf{v}} - o_n(1)$ can be constructed in polynomial time.

Note that the first case (that of \mathcal{F}_k) is equivalent to the special instance of the third case when the color set is $C = \{\text{red}, \text{blue}\}$. Indeed, for a graph in \mathcal{F}_k we can color its edges blue and its non-edges red thereby obtaining \mathcal{C}_k , and when considering an n -vertex graph G and an optimal \mathcal{F}_k -decomposition of it, we can equivalently consider the optimal \mathcal{C}_k -decomposition of the blue-red coloring of K_n where the edges of G are colored blue and its non-edges are colored red. It therefore suffices to prove only cases 2 and 3 of Theorem 1.

The proof of Theorem 1 consists of two main ingredients. We first use a result from [22] which can, in particular, be formulated as follows. Given a family \mathcal{F} of graphs, and given a fractional \mathcal{F} -decomposition of G (assuming there is one), one can find a set P of subgraphs of G such that each element of P is isomorphic to an element of \mathcal{F} and any pair of vertices of G is in at most one element of P (if this pair is an edge of G , then it is also an edge in the element of P in which it appears). Furthermore, the number of pairs that are not covered by P is $o(|V(G)|^2)$. So, assuming G is dense, P is a packing of elements of \mathcal{F} in G such that almost all pairs of vertices of G are packed and in this sense, it is an ‘‘almost’’ \mathcal{F} -decomposition. The result in [22] extended an earlier result of Haxell and Rödl [8] where \mathcal{F} is a single graph. Both results are actually more general, as they show that any fractional packing (which may be far from a decomposition) can be converted to an integral packing with relatively small loss. If we apply this result for $\mathcal{F} = \mathcal{F}_k$ we are close to proving the first part of Theorem 1, but there are two caveats.

Since we now have a vector $\mathbf{v} \in \mathbb{R}^{|\mathcal{F}_k|}$ associated, even if we start with an optimal fractional decomposition attaining $\nu_{\mathbf{v}}^*(G)$ it could be that the obtained integral ‘‘almost decomposition’’ distributes the weights to the elements of \mathcal{F}_k in a way that decreases the total weight significantly

below $\nu_{\mathbf{v}}^*(G)$. However, fortunately, the proof from [22] (implicitly) shows that we can almost maintain the correct distribution.

The second (and more difficult) problem is that the obtained almost decomposition needs now to be modified to a full decomposition, and *without* affecting much the total weight, staying close to $\nu_{\mathbf{v}}^*(G)$. To this end, we use a fundamental result of Barber et al. [2] based on the method of iterative absorption which enables us to achieve this goal, but with a price. To achieve their setting, we need, in fact, to first decompose a graph with high minimum degree to edge disjoint copies of some K_m (here m is huge compared to k , but fixed), and apply the aforementioned result of [22] to each element of this K_m -decomposition separately (more precisely, to the subgraph of G induced by the vertices of that element). We then need to sparsify our obtained packing in order to achieve a setting suitable for the application of [2]. The second and third part of Theorem 1 are obtained using analogues of [22] for tournaments and edge-colored graphs.

Theorem 1 provides a convenient mechanism to study certain natural decomposition problems, as it is sometimes much easier to obtain bounds for the fractional problem. In fact, in many cases, we can glue optimal fractional decompositions of small graphs into good fractional decompositions for arbitrary large graphs. We next give two very natural applications, but one may construct additional.

Theorem 2 *Let $n \equiv 1, 3 \pmod{6}$. Any tournament on n vertices has a triangle decomposition where the number of directed triangles in the decomposition is only $0.0222n^2(1 + o(1))$.*

Let $\mathcal{F}^* \subset \mathcal{F}_k$. Consider the vector $\mathbf{v} \in \mathbb{R}^{|\mathcal{F}_k|}$ which assigns 0 to the elements of \mathcal{F}^* and 1 to the elements of $\mathcal{F}_k \setminus \mathcal{F}^*$. We say that \mathcal{F}^* is *essentially avoidable* if $\nu_{\mathbf{v}} = 1$. In other words, for every k -decomposable graph G , there is a k -decomposition of G which almost completely avoids using elements from \mathcal{F}^* (i.e. the fraction of elements of this decomposition which are isomorphic to elements of \mathcal{F}^* is $o_n(1)$). If $\mathcal{F}^* = \{H\}$, we say that H is essentially avoidable. A result from [23], together with the proof of Theorem 1 implies the following.

Theorem 3

1. C_5 is essentially avoidable. More generally, if k is odd and $\mathcal{F}^* \subset \mathcal{F}_k$ is the family of all graphs H on k vertices such that both H and its complement are Eulerian, then \mathcal{F}^* is essentially avoidable.
2. Almost all graphs are essentially avoidable. Namely, if $\mathcal{U}_k \subset \mathcal{F}_k$ is the set of all graphs on k vertices that are not essentially avoidable, then $|\mathcal{U}_k| = o(|\mathcal{F}_k|)$.

For the first nontrivial case $k = 3$, it is possible to determine $\nu_{\mathbf{v}}$ for every binary vector and certain additional types of vectors $\mathbf{v} \in \mathbb{R}^{|\mathcal{F}_3|}$ (these vectors are four dimensional as $|\mathcal{F}_3| = 4$). However, even for $k = 3$, there are still some types of vectors for which we do not know $\nu_{\mathbf{v}}$. For the case $k = 4$ the situation is even more involved as we still do not know $\nu_{\mathbf{v}}$ even for all binary vectors. We elaborate more on this in Section 4 which considers the small cases $k = 3, 4$.

It also seems plausible to try and evaluate the asymptotic behavior of $\nu_{\mathbf{v}}(G)$, namely, the asymptotic value of $\nu_{\mathbf{v}}(G)$ when $G \sim \mathcal{G}(n, p)$ is a random graph. In this case, it turns out that the problem can be completely solved, and the asymptotic value efficiently computed, for all \mathbf{v} and all constant $0 < p < 1$, as we prove in Section 5.

Our road-map follows. Section 2 contains the proof of Theorem 1. Our demonstrative applications, Theorem 2 and Theorem 3 are the theme of Section 3. Section 4 focuses on small cases. Section 5 analyzes $\nu_{\mathbf{v}}(G)$ when $G \sim \mathcal{G}(n, p)$.

2 Integer and fractional vector valued decompositions

As noted in the introduction, it suffices to prove the second and third parts of Theorem 1. In this section we mostly prove the third part (the edge-coloring case). The proof of the second part (the tournament case) follows along the same lines and requires only minor modifications, which are outlined in the last subsection of this section.

2.1 From fractional decomposition to a similarly distributed integer packing

Let C be a finite set of colors and recall that \mathcal{C}_k is the set of all edge colorings of K_k with colors from C . Suppose G is a graph whose edges are colored by C . We call such G a C -colored graph and note that here we do not assume that G is complete (so non-edges of G correspond to non-colored pairs). Let $\binom{G}{K_k}$ denote the set of K_k -subgraphs of G and for $H \in \mathcal{C}_k$, let $\binom{G}{H} \subseteq \binom{G}{K_k}$ be the set of K_k -subgraphs of G that are color-isomorphic to H . More formally, for each $X \in \binom{G}{H}$ there is a bijection $b : V(X) \rightarrow V(H)$ such that $(u, v) \in E(X)$ and $(b(u), b(v)) \in E(H)$ have the same color.

We can naturally extend the notion of a fractional K_k -decomposition to graphs that are not necessarily complete, such as G above. For an edge $e \in E(G)$, let $\binom{G}{K_k, e} \subseteq \binom{G}{K_k}$ be the set of K_k -subgraphs of G that contain e . We say that a function $f : \binom{G}{K_k} \rightarrow [0, 1]$ is a fractional K_k -decomposition if $\sum_{X \in \binom{G}{K_k, e}} f(X) = 1$ holds for each $e \in E(G)$. Notice that a necessary (though not sufficient) requirement for G to have a fractional K_k -decomposition is that each edge of G belongs to at least one K_k -subgraph of G .

Now, suppose f is a fractional K_k -decomposition of G and that G is C -colored. For $H \in \mathcal{C}_k$, let $f(G, H) = \sum_{X \in \binom{G}{H}} f(X)$. Since f is a fractional K_k -decomposition, we have that

$$\sum_{H \in \mathcal{C}_k} f(G, H) = \frac{|E(G)|}{\binom{k}{2}}. \quad (1)$$

The following lemma follows implicitly from a generalization of the proof of the main result of [22].

Lemma 2.1 *Let C be a finite set of colors, let $k \geq 3$ be an integer and let $\epsilon > 0$. There exists $n_0 = n_0(k, C, \epsilon)$ such that the following holds. Suppose G is C -colored and has $n > n_0$ vertices. Let f be a fractional K_k -decomposition of G . Then for every $H \in \mathcal{C}_k$ there is a set P_H of induced*

subgraphs of G that are color-isomorphic to H , such that $|P_H| \geq f(G, H) - \epsilon n^2$. Furthermore any two elements of $P = \cup_{H \in \mathcal{C}_k} P_H$ intersect in at most one vertex.

Since the main result of [22] is not proved for the edge-colored case (it is only for uncolored graphs) and since in any case the (rather short) proof there implies Lemma 2.1 only implicitly, we present the proof of Lemma 2.1 in Subsection 2.4.

Notice that P in Lemma 2.1 is, in particular, a packing of G with pairwise edge-disjoint copies of K_k . As the following corollary shows, if we take an optimal fractional decomposition with respect to some $\mathbf{v} \in \mathbb{R}^{|\mathcal{C}_k|}$ and apply Lemma 2.1 to it, we obtain an integral packing of G with elements of \mathcal{C}_k that is close to an optimal \mathcal{C}_k -decomposition with respect to \mathbf{v} . To be more formal, for $\mathbf{v} \in \mathbb{R}^{|\mathcal{C}_k|}$ indexed by \mathcal{C}_k and a fractional K_k -decomposition f of a C -colored graph G , let $D_{\mathbf{v}}(f) = \sum_{H \in \mathcal{C}_k} \mathbf{v}_H f(G, H)$. As before, after normalizing we define $\nu_{\mathbf{v}}^*(f) = D_{\mathbf{v}}(f) \binom{k}{2} / |E(G)|$ and define the *optimal* fractional K_k -decomposition of G with respect to \mathbf{v} by $\nu_{\mathbf{v}}^*(G) = \max_f \nu_{\mathbf{v}}^*(f)$ where the maximum is taken over all fractional K_k -decompositions of G . If G has no fractional K_k -decomposition, then define $\nu_{\mathbf{v}}^*(G) = 0$.

Corollary 2.2 *Let C be a finite set of colors, let $k \geq 3$ be an integer, let $\mathbf{v} \in \mathbb{R}^{|\mathcal{C}_k|}$, and let $\gamma > 0$. There exists $N_{2.2} = N_{2.2}(k, C, \gamma, \mathbf{v})$ such that the following holds for all C -colored graphs G with $n > N_{2.2}$ vertices which have a fractional K_k -decomposition. For every $H \in \mathcal{C}_k$ there is a set P_H of induced subgraphs of G that are color-isomorphic to H , such that any two elements of $P = \cup_{H \in \mathcal{C}_k} P_H$ intersect in at most one vertex. Furthermore,*

$$(a) \quad |P| \geq \frac{|E(G)| - \gamma n^2}{\binom{k}{2}}.$$

$$(b) \quad \sum_{H \in \mathcal{C}_k} \mathbf{v}_H |P_H| \geq \frac{|E(G)|}{\binom{k}{2}} \nu_{\mathbf{v}}^*(G) - \gamma n^2.$$

Proof. Let $s = \sum_{H \in \mathcal{C}_k} \mathbf{v}_H$. Define $\epsilon = \gamma / (\binom{k}{2} |\mathcal{C}_k|)$ if $s < 1$ else set $\epsilon = \gamma / (\binom{k}{2} |\mathcal{C}_k| s)$. Let $N_{2.2}(k, C, \gamma, \mathbf{v}) = n_0(k, C, \epsilon)$ where the latter is the constant from Lemma 2.1. Let G be a C -colored graph having $n > N_{2.2}$ vertices. If G has no fractional K_k -decomposition, then there is nothing to prove, so assume that f is an optimal fractional K_k -decomposition of G with respect to \mathbf{v} , thus $\nu_{\mathbf{v}}^*(f) = \nu_{\mathbf{v}}^*(G)$. By Lemma 2.1, for every $H \in \mathcal{C}_k$ there is a set P_H of induced subgraphs of G that are color-isomorphic to H such that $|P_H| \geq f(G, H) - \epsilon n^2$ and any two elements of $P = \cup_{H \in \mathcal{C}_k} P_H$ intersect in at most one vertex. Now,

$$\begin{aligned} \sum_{H \in \mathcal{C}_k} \mathbf{v}_H |P_H| &\geq \sum_{H \in \mathcal{C}_k} \mathbf{v}_H (f(G, H) - \epsilon n^2) \\ &= \left(\sum_{H \in \mathcal{C}_k} \mathbf{v}_H f(G, H) \right) - \epsilon n^2 s \\ &= \frac{|E(G)|}{\binom{k}{2}} \nu_{\mathbf{v}}^*(G) - \epsilon n^2 s \\ &\geq \frac{|E(G)|}{\binom{k}{2}} \nu_{\mathbf{v}}^*(G) - \gamma n^2 \end{aligned}$$

which proves (b). To see (a) we just use (1) and

$$|P| = \sum_{H \in \mathcal{C}_k} |P_H| \geq \left(\sum_{H \in \mathcal{C}_k} f(G, H) \right) - \epsilon n^2 |\mathcal{C}_k| \geq \frac{|E(G)| - \gamma n^2}{\binom{k}{2}}.$$

■

2.2 From packing to decomposition

The following lemma is a major ingredient of the proof of Theorem 1. Recall that an *equitable partition* of a graph G into q parts $\mathcal{P} = \{W_1, \dots, W_q\}$ is a partition of $V(G)$ such that $||W_i| - |W_j|| \leq 1$ for all $1 \leq i < j \leq q$. The *Turán graph* with q parts, denoted by $T(n, q)$ is the complete q -partite graph on n vertices where the parts form an equitable partition.

Lemma 2.3 *Let $k \geq 3$ be an integer. Then there exists $q_{2.3}(k)$ such that for all $q \geq q_{2.3}$ there exist $N_{2.3}(q, k)$ and $\gamma = \gamma_{2.3}(q, k)$ such that the following holds for all $n > N_{2.3}$ for which K_n is k -divisible. Let G be a complete graph on n vertices, let \mathcal{P} be an equitable partition of G into q parts and let $G[\mathcal{P}]$ be the $T(n, q)$ spanning subgraph of G formed by the parts of \mathcal{P} . Suppose P is a packing of $G[\mathcal{P}]$ with pairwise edge-disjoint copies of K_k such that at most γn^2 edges of $G[\mathcal{P}]$ are uncovered by elements of P . Then, there is a sub-packing $P' \subseteq P$ such that $|P| - |P'| \leq 8\sqrt{\gamma}n^2$ and there is a K_k -decomposition of G that contains P' .*

The proof of Lemma 2.3 mainly follows from the proof of the main result of Barber et al. [2]. We prove it in Subsection 2.3. We will also need the following result which states that a graph with large enough minimum degree has a fractional K_m -decomposition.

Lemma 2.4 [1, 4, 21] *For every integer $m \geq 3$, there exists $\alpha = \alpha(m) < 1$ such that every graph on n vertices and minimum degree at least αn has a fractional K_m -decomposition.* ■

The first bound for α was given in [21] who proved that $\alpha \leq 1 - 1/(9m^{10})$. This was later improved in [4] to $1 - 2/(9m^2(m-1)^2)$ and in [1] to $1 - 1/(10^4 m^{3/2})$. It is worth noting that recently, an even stronger version of Lemma 2.4 has been proved by Barber et al. [2]. In particular, they have proved that if n is sufficiently large, and an n -vertex graph with minimum degree at least αn is K_k -divisible, then it has a K_m -decomposition (with roughly the same α as the one required for the fractional K_m -decomposition). However, using this stronger version for Lemma 2.4 will not make a difference in our arguments that follow. As mentioned above, we will, however, need to use the result from [2] later in a subtler setting in order to prove Lemma 2.3.

Next, we need the following simple lemma.

Lemma 2.5 *The sequence $\nu_{\mathbf{v}}^*(n)$ is non-decreasing and bounded from above, hence the limit $\nu_{\mathbf{v}}^*$ exists. In particular, for every $\epsilon > 0$, there exists m such that $\nu_{\mathbf{v}}^*(m) \geq \nu_{\mathbf{v}}^* - \epsilon$.*

Proof. Let s be the maximum coordinate of \mathbf{v} . Let f be a fractional K_k -decomposition of $G = K_n$. Then, f has $\nu_{\mathbf{v}}^*(f) \leq s$. So, the sequence $\nu_{\mathbf{v}}^*(n)$ is bounded from above by s . Next, we show that $\nu_{\mathbf{v}}^*(n) \geq \nu_{\mathbf{v}}^*(n-1)$. Let G be a C -colored complete graph on n vertices. For each $v \in V(G)$, let G_v be the induced subgraph of G on $V(G) \setminus v$ and let f_v be an optimal fractional K_k -decomposition of G_v with respect to \mathbf{v} . So, by definition $\nu_{\mathbf{v}}^*(f_v) \geq \nu_{\mathbf{v}}^*(n-1)$.

Next, define a fractional K_k -decomposition f of G as follows. For each induced k -vertex subgraph X of G , let

$$f(X) = \frac{1}{n-2} \sum_{v \in V(G) \setminus V(X)} f_v(X).$$

It is easy to verify that the sum of the weights corresponding to each pair of vertices is precisely 1 so f is indeed a fractional K_k -decomposition of G and that

$$\nu_{\mathbf{v}}^*(f) = \frac{1}{n} \sum_{v \in V(G)} \nu_{\mathbf{v}}^*(f_v) \geq \nu_{\mathbf{v}}^*(n-1).$$

Hence, $\nu_{\mathbf{v}}^*(G) \geq \nu_{\mathbf{v}}^*(n-1)$ implying that $\nu_{\mathbf{v}}^*(n) \geq \nu_{\mathbf{v}}^*(n-1)$ and that the sequence is non-decreasing. ■

The following lemma immediately implies the third part of Theorem 1.

Lemma 2.6 *Let C be a finite set of colors, let $k \geq 3$ be an integer, let $\mathbf{v} \in \mathbb{R}^{|C_k|}$ be indexed by C_k , and let $\epsilon > 0$. Then there exist $N_{2.6} = N_{2.6}(\epsilon, \mathbf{v})$ such that the following holds. Let G be a C -colored complete graph which is k -divisible and with $n > N_{2.6}$ vertices. Then, G is K_k -decomposable and, furthermore $\nu_{\mathbf{v}}(G) \geq (\nu_{\mathbf{v}}^* - \epsilon)(1 - \epsilon) - \epsilon$.*

Proof. First notice that the lemma indeed implies the third part of Theorem 1 since on the one hand we always have $\nu_{\mathbf{v}}(n) \leq \nu_{\mathbf{v}}^*(n) \leq \nu_{\mathbf{v}}^*$ and on the other hand, the lemma shows that for every $\epsilon > 0$, if n is sufficiently large, then $\nu_{\mathbf{v}}(n) \geq (\nu_{\mathbf{v}}^* - \epsilon)(1 - \epsilon) - \epsilon$. Hence the limit $\nu_{\mathbf{v}}$ exists and equals $\nu_{\mathbf{v}}^*$.

We next establish some constants that are required for the proof and for the definition of $N_{2.6}$. Let $\epsilon > 0$ be given as in the statement of the lemma. Let $m \geq k$ be the smallest integer such that $\nu_{\mathbf{v}}^*(m) \geq \nu_{\mathbf{v}}^* - \epsilon/2$. Notice that m exists by Lemma 2.5. Let $\alpha = \alpha(m)$ be the constant from Lemma 2.4. Let $q = \lceil \max\{2/(1 - \alpha), 5/\epsilon, q_{2.3}(k)\} \rceil$. Let s be the maximum of 1 and the maximum coordinate of \mathbf{v} . Let $\gamma = \min\{\epsilon^2/(1024s^2k^4), \gamma_{2.3}(q, k)\}$. Let $N_{2.6} = \max\{N_{2.2}(k, C, \gamma, \mathbf{v}), N_{2.3}(q, k)\}$. Let $n > N_{2.6}$.

Let G be a C -colored complete graph on n vertices which is k -divisible. Consider some arbitrary equitable partition \mathcal{P} of G into q parts. Let $G[\mathcal{P}]$ denote the spanning subgraph of G consisting of all edges whose endpoints are in distinct parts. Notice that $G[\mathcal{P}]$ is no longer complete, but we still view $G[\mathcal{P}]$ as a C -colored graph, where the edges of $G[\mathcal{P}]$ retain their colors. Clearly, the minimum degree of $G[\mathcal{P}]$ satisfies $\delta(G[\mathcal{P}]) \geq \lfloor n - n/q \rfloor$ since in $G[\mathcal{P}]$ each vertex is adjacent to all other vertices but those in its part. By our choice of q we have that $\delta(G[\mathcal{P}]) \geq \alpha n$. Hence, by Lemma 2.4, $G[\mathcal{P}]$ has a fractional K_m -decomposition, call it g .

Recall that $\binom{G[\mathcal{P}]}{K_m}$ denotes the set of all K_m -subgraphs of $G[\mathcal{P}]$. So, $g : \binom{G[\mathcal{P}]}{K_m} \rightarrow [0, 1]$ is such that for each edge of $G[\mathcal{P}]$, the sum of the values of g over all elements of $\binom{G[\mathcal{P}]}{K_m}$ that contain the edge is 1. We now define, for each $X \in \binom{G[\mathcal{P}]}{K_m}$, a fractional K_k -decomposition, denoted by f_X . We take f_X to be an optimal fractional K_k -decomposition of X with respect to \mathbf{v} (notice that f_X exists since X is a complete \mathcal{C} -colored graph and $|X| = m \geq k$). Thus, $\nu_{\mathbf{v}}^*(f_X) = \nu_{\mathbf{v}}^*(X)$.

We next define a fractional K_k -decomposition of $G[\mathcal{P}]$ denoted by f , as follows. Let Y be some K_k -subgraph of $G[\mathcal{P}]$. Let

$$f(Y) = \sum_{X \in \binom{G[\mathcal{P}]}{K_m}, V(X) \supset V(Y)} f_X(Y)g(X). \quad (2)$$

Notice that f is indeed a fractional K_k -decomposition of $G[\mathcal{P}]$ since f_X is such for every $X \in \binom{G[\mathcal{P}]}{K_m}$ and since g is a fractional K_m -decomposition of $G[\mathcal{P}]$.

We next estimate $D_{\mathbf{v}}(f) = \sum_{H \in \mathcal{C}_k} \mathbf{v}_H f(G[\mathcal{P}], H)$. By (2) we have:

$$\begin{aligned} D_{\mathbf{v}}(f) &= \sum_{X \in \binom{G[\mathcal{P}]}{K_m}} g(X)D_{\mathbf{v}}(f_X) \\ &= \sum_{X \in \binom{G[\mathcal{P}]}{K_m}} g(X)\nu_{\mathbf{v}}^*(f_X) \frac{\binom{m}{2}}{\binom{k}{2}} \\ &= \sum_{X \in \binom{G[\mathcal{P}]}{K_m}} g(X)\nu_{\mathbf{v}}^*(X) \frac{\binom{m}{2}}{\binom{k}{2}} \\ &\geq \sum_{X \in \binom{G[\mathcal{P}]}{K_m}} g(X)\nu_{\mathbf{v}}^*(m) \frac{\binom{m}{2}}{\binom{k}{2}} \\ &= \nu_{\mathbf{v}}^*(m) \frac{\binom{m}{2}}{\binom{k}{2}} \frac{|E(G[\mathcal{P}])|}{\binom{m}{2}} \\ &\geq \left(\nu_{\mathbf{v}}^* - \frac{\epsilon}{2} \right) \frac{|E(G[\mathcal{P}])|}{\binom{k}{2}}. \end{aligned}$$

Since $\nu_{\mathbf{v}}^*(f) = D_{\mathbf{v}}(f) \binom{k}{2} / |E(G[\mathcal{P}])|$ we obtain from the last inequality that

$$\nu_{\mathbf{v}}^*(G[\mathcal{P}]) \geq \nu_{\mathbf{v}}^*(f) \geq \nu_{\mathbf{v}}^* - \frac{\epsilon}{2}.$$

We now apply Corollary 2.2 to the graph $G[\mathcal{P}]$, which we can do since it has $n > N_{2.6} \geq N_{2.2}(k, \mathcal{C}, \gamma, \mathbf{v})$ vertices and since $G[\mathcal{P}]$ has a fractional K_k -decomposition. By the corollary, we obtain that for every $H \in \mathcal{C}_k$ there is a set P_H of induced subgraphs of $G[\mathcal{P}]$ that are color-isomorphic to H , such that any two elements of $P = \cup_{H \in \mathcal{C}_k} P_H$ intersect in at most one vertex. Furthermore,

$$\sum_{H \in \mathcal{C}_k} \mathbf{v}_H |P_H| \geq \frac{|E(G[\mathcal{P}])|}{\binom{k}{2}} \nu_{\mathbf{v}}^*(G[\mathcal{P}]) - \gamma n^2$$

and

$$|P| \geq \frac{|E(G[\mathcal{P}])| - \gamma n^2}{\binom{k}{2}}. \quad (3)$$

But recall that $E(G[\mathcal{P}])$ consists of all $\binom{n}{2}$ edges of G except those which have both of their endpoints in the same part of \mathcal{P} . Thus, $|E(G[\mathcal{P}])| \geq \binom{n}{2} - q \binom{\lceil n/q \rceil}{2} \geq \binom{n}{2} - n^2/q$. Also, we have already proved that $\nu_{\mathbf{v}}^*(G[\mathcal{P}]) \geq \nu_{\mathbf{v}}^* - \frac{\epsilon}{2}$. We therefore obtain using $q \geq 5/\epsilon$ that

$$\begin{aligned} \sum_{H \in \mathcal{C}_k} \mathbf{v}_{\mathbf{H}} |P_H| &\geq \frac{\binom{n}{2} - n^2/q}{\binom{k}{2}} \left(\nu_{\mathbf{v}}^* - \frac{\epsilon}{2} \right) - \gamma n^2 \\ &\geq \frac{\binom{n}{2}}{\binom{k}{2}} \left(1 - \frac{\epsilon}{2} \right) \left(\nu_{\mathbf{v}}^* - \frac{\epsilon}{2} \right) - \gamma n^2. \end{aligned} \quad (4)$$

Now, recall that each element $X \in P$ is also an induced K_k -subgraph of our complete graph G . Let G' denote the spanning subgraph of G consisting of all edges that are not covered by elements of P . Clearly, G' is K_k -divisible since both G and the complement of G' (which is the edge-disjoint union of K_k 's) are K_k -divisible. Now, suppose first that it was possible to find a K_k -decomposition of G' . Hence, in this case, there is a K_k -decomposition of G that contains P . We would therefore obtain from (4) that

$$\nu_{\mathbf{v}}(G) \geq \frac{\sum_{H \in \mathcal{C}_k} \mathbf{v}_{\mathbf{H}} |P_H|}{\binom{n}{2} / \binom{k}{2}} \geq \left(1 - \frac{\epsilon}{2} \right) \left(\nu_{\mathbf{v}}^* - \frac{\epsilon}{2} \right) - \frac{\epsilon}{2}. \quad (5)$$

Unfortunately, we have no guarantee that G' has a K_k -decomposition. Suppose, however, that it was possible to modify P just a bit, say, by removing just a few of the elements of P so that after this change, the corresponding remainder graph G' would have a K_k -decomposition. Then, almost the same bound for $\nu_{\mathbf{v}}(G)$ would apply, assuming that $\sum_{H \in \mathcal{F}_k} \mathbf{v}_{\mathbf{H}} |P_H|$ did not change much after the modification. Fortunately, this is possible, as a consequence of Lemma 2.3, as follows. We can apply Lemma 2.3 since by (3) P covers all but at most γn^2 edges of $G[\mathcal{P}]$. The lemma shows that there is a sub-packing $P' \subseteq P$ such that $|P| - |P'| \leq 8\sqrt{\gamma} n^2$ and there is a K_k -decomposition P^* of G that contains P' . Recall that s is the maximum of 1 and the maximum coordinate of \mathbf{v} . We therefore have by (5) that:

$$\nu_{\mathbf{v}}(G) \geq \left(1 - \frac{\epsilon}{2} \right) \left(\nu_{\mathbf{v}}^* - \frac{\epsilon}{2} \right) - \frac{\epsilon}{2} - \frac{8s\sqrt{\gamma}n^2}{\binom{n}{2} / \binom{k}{2}} \geq (1 - \epsilon)(\nu_{\mathbf{v}}^* - \epsilon) - \epsilon$$

where we have used that $\sqrt{\gamma} \leq \epsilon / (32sk^2)$. ■

Lemma 2.6 can be implemented in polynomial time as claimed in the statement of Theorem 1. Namely the K_k -decomposition P^* in the lemma can be constructed in time which is polynomial in $n = |V(G)|$. To see this, we first observe that Lemma 2.1 can be implemented in polynomial time (i.e. constructing the packing P in that lemma), as proved in [22]. This implies that Corollary 2.2 can be implemented in polynomial time, since finding the optimal fractional K_k -decomposition of

G with respect to \mathbf{v} denoted by f in the proof of the corollary can be found in polynomial time using linear programming (the number of variables is $O(n^k)$ as the number of K_k and the number of constraints is only $O(n^2)$ as the number of edges). Once we obtain the packing P of Corollary 2.2, we apply Lemma 2.3 which constructs P^* in polynomial time, as Lemmas 2.7 and 2.8 in Subsection 2.3 can be implemented in polynomial time as proved in [2].

2.3 Proof of Lemma 2.3

The proof of Lemma 2.3 is based on the proof of the main result of [2] (Theorem 1.3 there). In fact, we will only need to use a special case of that result, for the case of the small graph being K_k and for the case of the host graph being $G = K_n$ although most of the arguments in [2] are still required even for this special case, which is not surprising since this special case implies Wilson's decomposition theorem for the case of K_k . To achieve the setting in [2] we require some definitions taken from there.

For a graph G , a positive integer q and a real $\delta > 0$, a (q, δ) -partition of G is an equitable partition $\mathcal{P} = \{V_1, \dots, V_q\}$ of $V(G)$ such that for each $1 \leq i \leq q$ and each $v \in V(G)$, $d_G(v, V_i) \geq \delta|V_i|$. Here $d_G(v, V_i)$ denotes the number of neighbors of v in V_i . Notice that if $G = K_n$, then G trivially has a (q, δ) -partition, but a straightforward probabilistic argument shows that this also holds if G is just an n -vertex graph with minimum degree slightly larger than δn and n is sufficiently large (Proposition 7.3 in [2]). For an equitable partition \mathcal{P} into q parts, recall that $G[\mathcal{P}]$ denotes the q -partite subgraph of G induced by the parts of \mathcal{P} .

For an equitable partition \mathcal{P} into q parts, a *refinement* of \mathcal{P} is obtained by taking an equitable partition into q parts of each part of \mathcal{P} . Notice that a refinement is an equitable partition into q^2 parts.

Let \mathcal{P}_1 be an equitable partition of $V(G)$ and for each $2 \leq i \leq \ell$ let \mathcal{P}_i be a refinement of \mathcal{P}_{i-1} . We say that $\mathcal{P}_1, \dots, \mathcal{P}_\ell$ is a (q, δ, m) -partition sequence of G if the following hold.

- (i) \mathcal{P}_1 is a (q, δ) -partition of G .
- (ii) For each $2 \leq i \leq \ell$ and each $V \in \mathcal{P}_{i-1}$, $\mathcal{P}_i[V]$ is a (q, δ) -partition of $G[V]$.
- (iii) Each part of \mathcal{P}_ℓ is of size m or $m - 1$.

Once again, if $G = K_n$, then it trivially has a (q, δ, m) -partition sequence, but also if n is sufficiently large it is easy to prove that a graph with n vertices and minimum degree slightly larger than δn (say minimum degree at least $(\delta + \epsilon)n$) has a (q, δ, m) -partition sequence, where m is bounded by a constant depending only on q and ϵ (Lemma 7.4 in [2]).

The first major ingredient in the proof of Theorem 1.3 in [2] is that of the existence of an *absorber*. Informally, an absorber A^* of a K_k -divisible graph G is a K_k -divisible spanning subgraph of G with small maximum degree which has the following property. Suppose we take an "almost K_k -decomposition" of the spanning subgraph G' obtained from G after removing the edges of A^* . Let H^* be the leftover edges of G' uncovered by the almost decomposition. Note that H^* is also K_k -divisible. Then A^* has the property that $A^* \cup H^*$ has a K_k -decomposition (and hence so does

G). Of course, in order to obtain such an A^* we need to make sure that the set of possible H^* is small (in particular, if one can guarantee that H^* has no more than $O(n)$ edges, this will limit the number of possibilities for H^*). The formal definition of such an absorber is given in Lemma 8.1 there, which is stated here for the special case of K_k . Note that some of the notations have been changed to adjust to the notations in the present paper.

Lemma 2.7 *Let $k \geq 3$ and $\epsilon > 0$. Then there exists $m_{2.7}(\epsilon, k)$ such that the following holds for all $m \geq m_{2.7}$. There exists $N_{2.7}(\epsilon, k, m)$ such that for all $n > N_{2.7}$ the following holds. Set $\delta := 1 - 1/(3k) + \epsilon$, $t := \lceil n/m \rceil$ and let G be a graph with n vertices. Let $\mathcal{P} = \{V_1, \dots, V_t\}$ be an equitable partition of $V(G)$ so that each part has size m or $m - 1$. Suppose that $\delta(G[\mathcal{P}]) \geq \delta n$ and $\delta(G[V_i]) \geq \delta |V_i|$ for each $1 \leq i \leq t$. Then G contain a K_k -divisible subgraph A^* such that:*

- (i) $\Delta(A^*[\mathcal{P}]) \leq \epsilon^2 n$ and $\Delta(A^*[V_i]) < k$ for each $1 \leq i \leq t$.
- (ii) If H^* is a K_k -divisible graph on $V(G)$ that is edge-disjoint from A^* and has $E(H^*[\mathcal{P}]) = \emptyset$, then $H^* \cup A^*$ has a K_k -decomposition. ■

For a subgraph X of G let $G - X$ denote the spanning subgraph of G obtained by removing the edges of X . In order to apply Lemma 2.7 one first needs to decompose $G - (A^* \cup H^*)$. This is the other major ingredient in [2], which appears as Lemma 10.1 there. The following is a version of Lemma 10.1 for the special case of K_k and with an addendum that follows from its proof.

Lemma 2.8 *Let $k \geq 3$ and $\epsilon > 0$. Then there exists $q_{2.8}(k, \epsilon)$ such that the following holds for all $q \geq q_{2.8}$. There exists $\gamma_{2.8}(q, \epsilon, k)$ such that the following holds for all $\gamma \leq \gamma_{2.8}$, for all $m \geq m_{2.8}(\gamma)$ and for every K_k -divisible graph G on n vertices. Define $\delta := \max\{\alpha(k), 1 - 1/(3k)\}$ where $\alpha(k)$ is the constant from Lemma 2.4. Suppose $\mathcal{P}_1, \dots, \mathcal{P}_\ell$ is a $(q, \delta + \epsilon, m)$ -partition sequence of G . Then there exists a subgraph H^* of $\cup_{V \in \mathcal{P}_\ell} G[V]$ such that $G - H^*$ has a K_k -decomposition P^* . Furthermore, if P is packing of $G[\mathcal{P}_1]$ covering all but at most $2\gamma n^2$ edges of $G[\mathcal{P}_1]$, then there exists such a P^* such that $|P^* \setminus P| \leq 6\sqrt{\gamma} n^2$. ■*

We note that the ‘‘Furthermore’’ part does not appear in the statement of Lemma 10.1 in [2], but immediately follows from its proof. Indeed, the first part of the proof (Lemma 10.6 there) proceeds as follows. Take any packing P of $G[\mathcal{P}_1]$ that covers all but at most $2\gamma n^2$ edges of G . By removing at most $6\sqrt{\gamma} n^2$ elements from P you obtain a packing P' such that the subgraph H of G consisting of the edges of G that are uncovered by P' has some nice properties (stated as (G1) and (G2) in Lemma 10.6). From there onwards the proof Lemma 10.1 proceeds by iteratively improving P' in $\ell - 1$ steps where in step i one obtains an almost optimal packing in each part of \mathcal{P}_i which covers also the remaining uncovered edges between parts of the previous partition \mathcal{P}_{i-1} until obtaining P^* of Lemma 2.8. In particular, P^* still retains almost all of the element of the initial packing P , but at most $6\sqrt{\gamma} n^2$ elements.

Proof of Lemma 2.3: Let $k \geq 3$ be an integer. Define the following constants.

- (i) $\delta := \max\{\alpha(k), 1 - 1/(3k)\}$ where $\alpha(k)$ is the constant from Lemma 2.4.

- (ii) $\epsilon = (1 - \delta)/10$ and $\epsilon' = \epsilon/3$.
- (iii) $q_{2.3}(k) = \max\{2/(1 - \delta - \epsilon'), q_{2.8}(k, \epsilon')\}$ and let $q \geq q_{2.3}$.
- (iv) $\gamma = \gamma_{2.3}(q, k) = \gamma_{2.8}(q, \epsilon', k)$.
- (v) $m = \max\{m_{2.8}(\gamma), m_{2.7}(\gamma, k)\}$.
- (vi) $N_{2.3}(q, k) = N_{2.7}(\gamma, k, m)$.

Now let $n > N_{2.3}$ such that K_n is k -divisible. Let G be a complete graph on n vertices, let \mathcal{P} be an equitable partition of G into q parts and let $G[\mathcal{P}]$ be the $T(n, q)$ spanning subgraph of G formed by the parts of \mathcal{P} . Suppose P is a packing of $G[\mathcal{P}]$ with pairwise edge-disjoint copies of K_k such that at most γn^2 edges of $G[\mathcal{P}]$ are uncovered by elements of P .

Suppose $\mathcal{P}_1, \dots, \mathcal{P}_\ell$ is a $(q, \delta + \epsilon, m)$ -partition sequence of G where $\mathcal{P}_1 = \mathcal{P}$. Observe that such a $(q, \delta + \epsilon, m)$ -partition exists since G is a complete graph, since $\mathcal{P}_1 = \mathcal{P}$ is an equitable partition into q parts, and since $\delta + \epsilon < 1$.

Let $G_1 = G[\mathcal{P}] = G[\mathcal{P}_1]$ and let $G_{\ell+1} = G - G[\mathcal{P}_\ell]$. So, $G_{\ell+1}$ consists of all edges with both endpoints in the same part of \mathcal{P}_ℓ . Consider now the graph $H = G_1 \cup G_{\ell+1}$ (i.e. the spanning subgraph of G consisting of all the edges of G_1 and $G_{\ell+1}$) and consider the partition \mathcal{P}_ℓ of H . First observe that the minimum degree $\delta(H[\mathcal{P}_\ell])$ is at least $n - \lceil n/q \rceil \geq (\delta + \epsilon')n$ where we have used here that $q \geq 2/(1 - \delta - \epsilon')$. Similarly, for each $V \in \mathcal{P}_\ell$ we have $\delta(H[V]) = |V| - 1 \in \{m - 1, m - 2\}$. So, $\delta(H[V]) \geq (\delta + \epsilon')|V|$ since $\delta + \epsilon' < 1$. We may therefore apply Lemma 2.7 where H plays the role of G , γ plays the role of ϵ and \mathcal{P}_ℓ plays the role of \mathcal{P} .

By Lemma 2.7, H contains a K_k -divisible subgraph A^* such that:

- (i) $\Delta(A^*[\mathcal{P}_\ell]) \leq \gamma^2 n$ and $\Delta(A^*[V]) < k$ for each $V \in \mathcal{P}_\ell$.
- (ii) If H^* is a K_k -divisible graph on $V(G) = V(H)$ that is edge-disjoint from A^* and $E(H^*[\mathcal{P}_\ell]) = \emptyset$, then $H^* \cup A^*$ has a K_k -decomposition.

Observe that (i) and (ii) imply also that $\Delta(A^*) < \gamma^2 n + k$. Let $G' = G - A^*$. Thus, G' is also K_k -divisible. Note that for each $V \in \mathcal{P}_1$ and each $v \in V(G)$ we have $d_{G'}(v, V) \geq d_G(v, V) - \Delta(A^*) \geq (|V| - 1) - (\gamma^2 n + k - 1) \geq (\delta + \epsilon')|V|$. So, \mathcal{P}_1 is a $(q, \delta + \epsilon')$ -partition of G' . Note also that by (i) we have that $\Delta(A^* - A^*[\mathcal{P}_1]) < k$ so $\mathcal{P}_1, \dots, \mathcal{P}_\ell$ is also a $(q, \delta + \epsilon', m)$ -partition sequence of G' . Recall also that the packing P covered at most γn^2 edges of $G[\mathcal{P}]$. Let $P'' \subset P$ be the elements of P which are entirely in G' . Hence, each element of $P \setminus P''$ contain an edge of A^* . Since $\Delta(A^*) < \gamma^2 n + k$, we have that the number of edges of A^* is at most $\gamma^2 n^2 + nk$. It follows that P'' covers all elements of $G'(\mathcal{P})$ but at most $\gamma n^2 + \binom{k}{2}(\gamma^2 n^2 + nk) < 2\gamma n^2$.

We can therefore apply Lemma 2.8 to G' playing the role of G , ϵ' playing the role of ϵ , and P'' playing the role of P in that lemma. By Lemma 2.8 we obtain a subgraph H^* of $\cup_{V \in \mathcal{P}_\ell} G'[V]$ such that $G' - H^*$ has a K_k -decomposition P^* . Furthermore, $|P^* \setminus P''| \leq 6\sqrt{\gamma}n^2$. But now, by (ii) $A^* \cup H^*$ has a K_k -decomposition, so together with P^* this forms a K_k -decomposition of $G = K_n$ containing all but at most $6\sqrt{\gamma}n^2$ elements of P'' thus all but at most $6\sqrt{\gamma}n^2 + 2\gamma n^2 \leq 8\sqrt{\gamma}n^2$ elements of P . ■

2.4 Proof of Lemma 2.1

As noted earlier, Lemma 2.1 follows implicitly from the main result in [22]. That result is stated in terms of uncolored graphs, while here we need the colored version. Thus, we reproduce the arguments in the proof of [22] where the lemmas there whose proofs remain identical or for which the colored version is an immediate extension are only restated in their colored version without proof, but with reference to the original lemma in [22].

We first need to recall the edge-colored version of the Szemerédi's regularity lemma [15]. Let $G = (V, E)$ be a C -colored graph, and let A and B be two disjoint subsets of $V(G)$. If A and B are non-empty and $c \in C$, let $E_c(A, B)$ denote the set of edges between them that are colored c . The c -density between A and B is defined as

$$d_c(A, B) = \frac{|E_c(A, B)|}{|A||B|}.$$

For $\gamma > 0$ the pair (A, B) is called γ -regular if for every $X \subseteq A$ and $Y \subseteq B$ satisfying $|X| \geq \gamma|A|$ and $|Y| \geq \gamma|B|$ we have

$$|d_c(X, Y) - d_c(A, B)| \leq \gamma \quad \text{for all } c \in C.$$

An equitable partition of the set of vertices V of a C -colored graph G into the classes V_1, \dots, V_m is called γ -regular if all but at most $\gamma \binom{m}{2}$ of the pairs (V_i, V_j) are γ -regular. The regularity lemma (colored version) states the following:

Lemma 2.9 *Let C be a finite set of colors and let $\gamma > 0$. There is an integer $M(\gamma, C) > 0$ such that for every C -colored graph G of order $n > M$ there is a γ -regular partition of the vertex set of G into m classes, for some $1/\gamma < m < M$. ■*

The proof of Lemma 2.9 is completely analogous to the proof of the original regularity lemma.

For an edge (x, y) of a C -colored graph, let $c(x, y)$ denote its color. Let H be a C -colored graph with $V(H) = \{1, \dots, k\}$, $k \geq 3$. Let W be a C -colored k -partite graph with vertex classes V_1, \dots, V_k . A subgraph J of W with $V(J) = \{v_1, \dots, v_k\}$ is *partite-color-isomorphic* to H if $v_i \in V_i$ for $i = 1, \dots, k$ and the map $i \rightarrow v_i$ is a *color preserving isomorphism* from H to J . Namely, $(i, j) \in E(H)$ if and only if $(v_i, v_j) \in E(J)$ and in case they are both edges, then $c(i, j) = c(v_i, v_j)$.

The following is a standard counting lemma whose proof follows from the definition of γ -regularity. It is analogous to Lemma 2.2 of [22].

Lemma 2.10 *Let C be a finite set of colors, let $k \geq 3$ be a positive integer, and let δ and ζ be positive reals. There exist $\gamma = \gamma(\delta, \zeta, k, C)$ and $T = T(\delta, \zeta, k, C)$ such that the following holds. Let H be a C -colored graph with $V(H) = \{1, \dots, k\}$ and let W be a C -colored k -partite graph with vertex classes V_1, \dots, V_k where $|V_i| = t > T$ for $i = 1, \dots, k$. Furthermore, for each $(i, j) \in E(H)$, (V_i, V_j) is a γ -regular pair with $d_{c(i,j)}(V_i, V_j) \geq \delta$ and for each $(i, j) \notin E(H)$, $E(V_i, V_j) = \emptyset$. Then, there exists a spanning subgraph W' of W , consisting of at least $(1 - \zeta)|E(W)|$ edges such that the following holds. For an edge $e \in E(W')$, let $\text{count}(e)$ denote the number of subgraphs of W' that*

are partite-color-isomorphic to H and that contain e . Then, for all $e \in E(W')$, if $e \in E(V_i, V_j)$, then

$$\left| \text{count}(e) - t^{k-2} \frac{\prod_{(s,p) \in E(H)} d_{c(s,p)}(V_s, V_p)}{d_{c(i,j)}(V_i, V_j)} \right| < \zeta t^{k-2}.$$

■

We need the result of Frankl and Rödl [5] on near perfect matchings of uniform hypergraphs. Recall that if x, y are two vertices of a hypergraph then $\deg(x)$ denotes the degree of x and $\deg(x, y)$ denotes the number of edges that contain both x and y . We use the version of the Frankl and Rödl Theorem due to Pippenger.

Lemma 2.11 *For an integer $r \geq 2$ and a real $\beta > 0$ there exists $\mu = \mu(r, \beta) > 0$ so that: If the r -uniform hypergraph L on q vertices has the following properties for some d :*

(i) $(1 - \mu)d < \deg(x) < (1 + \mu)d$ holds for all vertices,

(ii) $\deg(x, y) < \mu d$ for all distinct x and y ,

then L has a matching of size at least $(q/r)(1 - \beta)$.

■

Let C be a finite set of colors, let $k \geq 3$ be an integer and let $\epsilon > 0$. Let $\delta = \beta = \epsilon/4$. Let $\mu = \mu(\binom{k}{2}, \beta)$ be as in Lemma 2.11. Let $\zeta = \mu\delta^{k^2}/2$. Let $\gamma = \gamma(\delta, \zeta, k, C)$ and $T = T(\delta, \zeta, k, C)$ be as in Lemma 2.10. Let $M = M(\gamma\epsilon/(25k^2), C)$ be as in Lemma 2.9. Finally, we shall define $n_0 = n_0(k, C, \epsilon)$ to be a sufficiently large constant, depending on the above chosen parameters, and for which the inequalities stated in the proof below hold.

Fix an n -vertex C -colored graph G with $n > n_0$ vertices and assume that G has a fractional K_k -decomposition $f : \binom{G}{K_k} \rightarrow [0, 1]$. We apply Lemma 2.9 to G and obtain a γ' -regular partition with m' parts, where $\gamma' = \gamma\epsilon/(25k^2)$ and $1/\gamma' < m' < M$. Denote the parts by $U_1, \dots, U_{m'}$. Notice that the size of each part is either $\lfloor n/m' \rfloor$ or $\lceil n/m' \rceil$. For simplicity we may and will assume that n/m' is an integer, as this assumption does not affect the asymptotic nature of the result. Similarly, we assume that $25k^2/\epsilon$ and $n/(25m'k^2/\epsilon)$ are integers.

We randomly partition each U_i into $25k^2/\epsilon$ equal parts of size $n/(25m'k^2/\epsilon)$ each. All m' partitions are independent. We now have $m = 25m'k^2/\epsilon$ refined vertex classes, denoted V_1, \dots, V_m . Suppose $V_i \subset U_s$ and $V_j \subset U_t$ where $s \neq t$. We claim that if (U_s, U_t) is a γ' -regular pair, then (V_i, V_j) is a γ -regular pair. Indeed, if $X \subseteq V_i$ and $Y \subseteq V_j$ have $|X|, |Y| \geq \gamma n/(25m'k^2/\epsilon)$, then $|X|, |Y| \geq \gamma' n/m'$ and so $|d_c(X, Y) - d_c(U_s, U_t)| \leq \gamma'$ for each $c \in C$. Also $|d_c(V_i, V_j) - d_c(U_s, U_t)| \leq \gamma'$. Thus, $|d_c(X, Y) - d_c(V_i, V_j)| \leq 2\gamma' \leq \gamma$.

Let X be some K_k -subgraph of G . We call X *good* if its k vertices belong to distinct vertex classes of the refined partition. Since the probability that two vertices of X belong to the same vertex class of the refined partition is less than $\epsilon/(25k^2)$, the probability that X is not good is at most $\binom{k}{2}\epsilon/(25k^2) < \epsilon/50$. Since f is a fractional K_k -decomposition, the sum of its values is $|f| = |E(G)|/\binom{k}{2} < n^2$. Hence, if f^{**} is the restriction of f to good elements (the non-good elements having $f^{**}(X) = 0$), then the expected sum of the values of f^{**} is at least $|f|(1 - \epsilon/50)$.

We therefore *fix* a partition V_1, \dots, V_m for which $|f^{**}| \geq |f|(1 - \epsilon/50)$. Notice that f^{**} is no longer a fractional k -decomposition; it is merely a fractional K_k -packing of G (i.e. for each edge of G , the sum of the values of f^{**} on the elements of $\binom{G}{K_k}$ that contain the edge is at most 1). Furthermore, for each $H \in \mathcal{C}_k$ we have that

$$f^{**}(G, H) \geq f(G, H) - (|f| - |f^{**}|) \geq f(G, H) - \frac{\epsilon}{50}|f| \geq f(G, H) - \frac{\epsilon}{50}n^2.$$

Let G^* be the spanning subgraph of G consisting of the following edges: An edge $(u, v) \in E(G)$ is in $E(G^*)$ if and only if $u \in V_i, v \in V_j, i \neq j, (V_i, V_j)$ is a γ -regular pair, and $d_{c(u,v)}(V_i, V_j) \geq \delta$. (thus, we discard edges inside classes, between non regular pairs, or if the color of the edge is sparse in the pair to which it belongs). Let f^* be the restriction of f^{**} to copies of K_k in G^* . We claim that $|f^*| > |f^{**}| - 0.6\delta n^2$. Indeed, by considering the number of discarded edges we get (using $\delta \gg \gamma' \geq 1/m'$)

$$\begin{aligned} |f^{**}| - |f^*| &\leq |E(G) - E(G^*)| \\ &< \gamma' \binom{m'}{2} \frac{n^2}{m'^2} + \binom{m'}{2} (\delta + \gamma') \frac{n^2}{m'^2} + m' \binom{n/m'}{2} \\ &< 0.6\delta n^2. \end{aligned}$$

In particular, for each $H \in \mathcal{C}_k$ we have that

$$f^*(G^*, H) \geq f^{**}(G, H) - (|f^{**}| - |f^*|) \geq f(G, H) - \frac{\epsilon}{50}n^2 - 0.6\delta n^2 \geq f(G, H) - \frac{\epsilon}{10}n^2.$$

Let R denote the m -vertex *multigraph* whose vertices are $\{1, \dots, m\}$ and a pair (i, j) with color c is an edge of R if and only if (V_i, V_j) is a γ -regular pair and $d_c(i, j) \geq \delta$. Notice that R is indeed a multigraph but any two multiple edges have distinct colors. We define a fractional K_k -packing f' of R as follows. Let X be a subgraph of R that is color-isomorphic to some $H \in \mathcal{C}_k$ and assume that the vertices of X are $\{u_1, \dots, u_k\}$ where u_i plays the role of vertex i in H . We define $f'(X)$ to be the sum of the values of f^* taken over all subgraphs of $G^*[V_{u_1}, \dots, V_{u_k}]$ which are partite-color-isomorphic to H , divided by n^2/m^2 (and where the isomorphism is $i \rightarrow u_i$). Notice that $|f'| = m^2|f^*|/n^2$ since every K_k -subgraph of G^* contributes its weight (divided by n^2/m^2) to the sum of the weights of f' . Likewise

$$f'(R, H) = f^*(G^*, H) \frac{m^2}{n^2}.$$

We use f' to define a random partition of $E(G^*)$. Our parts correspond to the copies of elements of $\binom{R}{K_k}$. We denote the partition by $\mathcal{Q} = \{Q_X : X \in \binom{R}{K_k}\}$. Let $X \in \binom{R}{K_k}$ and assume that X contains the edge (i, j) of $E(R)$ and that the color of the edge is c . Each $e \in E_c(V_i, V_j)$ (which, by the definition of R , must be an edge of G^*) is chosen to be in Q_X with probability $f'(X)/d_c(V_i, V_j)$. The choices made by distinct edges of G^* are independent. Notice that this random coloring is legal (in the sense that the sum of probabilities is at most one) since the sum of $f'(X)$ taken over

all possible X containing the edge (i, j) of $E(R)$ whose color is c is at most $d_c(V_i, V_j)$. Notice also that some edges of G^* might stay unassigned to a part in our random partitioning (as maybe an edge (i, j) of $E(R)$ whose color is c does not belong to any X). In this case, we can assign such unassigned edges of G^* to some “spare part”, denoted Q_0 , so that $\mathcal{Q} = \{Q_X : X \in \binom{R}{K_k}\} \cup \{Q_0\}$ is indeed a partition of $E(G^*)$.

Let X be a subgraph of R that is color-isomorphic to some $H \in \mathcal{C}_k$, and assume that $f'(X) > m^{1-k}$ (we need this assumption in the lemmas below). Without loss of generality, assume that the vertices of X are $\{1, \dots, k\}$ where $i \in V(X)$ plays the role of $i \in V(H)$. Let $W_X = G^*[V_1, \dots, V_k]$. Notice that W_X is a subgraph of G^* which satisfies the conditions in Lemma 2.10, since $t = n/m > n_0\epsilon/(25k^2M) > T$ (here we assume $n_0 > 25k^2MT/\epsilon$). Let W'_X be the spanning subgraph of W_X whose existence is guaranteed in Lemma 2.10. Let Z_X denote the spanning subgraph of W'_X consisting only of the edges that belong to the part Q_X . Notice that Z_X is a random subgraph of W'_X . For an edge $e \in E(Z_X)$, let $S_X(e)$ denote the set of subgraphs of Z_X that contain e and that are partite-color-isomorphic to H . Put $s_X(e) = |S_X(e)|$. the proof of the following two lemmas are identical to the proofs of Lemmas 3.1 and Lemma 3.2 in [22], respectively.

Lemma 2.12 *With probability at least $1 - m^3/n$, for all $e \in E(Z_X)$,*

$$\left| s_X(e) - t^{k-2} f'(X) \binom{k}{2}^{-1} \right| < \mu f'(X) \binom{k}{2}^{-1} t^{k-2}.$$

■

Lemma 2.13 *With probability at least $1 - 1/n$,*

$$|E(Z_X)| > (1 - 2\zeta) \binom{k}{2} \frac{n^2}{m^2} f'(X).$$

■

Since R contains at most $O(m^k)$ copies of K_k , we have that with probability at least $1 - O(m^k/n) - O(m^{k+3}/n) > 0$ (here we assume again that n_0 is sufficiently large) *all* copies X of K_k in R with $f'(X) > m^{1-k}$ satisfy the statements of Lemma 2.12 and Lemma 2.13. We therefore fix a partition \mathcal{Q} for which Lemma 2.12 and Lemma 2.13 hold for all such X .

Let $H \in \mathcal{C}_k$. Let X be a copy of K_k in R with $f'(X) > m^{1-k}$ that is partite-color-isomorphic to H . We construct an r -uniform hypergraph L_X as follows. The vertices of L_X are the edges of Z_X . The edges of L_X correspond to the edge sets of the subgraphs of Z_X that are partite-color-isomorphic to H . We claim that this hypergraph satisfies the conditions of Lemma 2.11. Indeed, let q denote the number of vertices of L_X . Let $d = t^{k-2} f'(X) \binom{k}{2}^{-1}$. Notice that by Lemma 2.12 *all* vertices of L_X have their degrees between $(1 - \mu)d$ and $(1 + \mu)d$. Also notice that the co-degree of any two vertices of L_X is at most t^{k-3} as two edges cannot belong, together, to more than t^{k-3} subgraphs of L_X that are partite-color-isomorphic to H . Also observe that for n_0 sufficiently large,

$\mu d > t^{k-3}$. By Lemma 2.11 we have a set \mathcal{S}_X of at least $(q/\binom{k}{2})(1-\beta)$ pairwise edge-disjoint subgraphs of Z_X that are partite-color-isomorphic to H . In particular, by Lemma 2.13,

$$|\mathcal{S}_X| \geq (1-\beta)(1-2\zeta)\frac{n^2}{m^2}f'(X) > (1-2\beta)f'(X)\frac{n^2}{m^2}.$$

Now, let \mathcal{X}_H be the set of all subgraphs of R that are partite-color-isomorphic to H . By definition, $f'(R, H) = \sum_{X \in \mathcal{X}_H} f'(X)$. Since trivially $|\mathcal{X}_H| \leq m^k$, the total contribution of the elements $X \in \mathcal{X}_H$ with $f'(X) \leq m^{1-k}$ to the sum is at most m . Hence,

$$\begin{aligned} \left| \bigcup_{X \in \mathcal{X}_H, f'(X) > m^{1-k}} \mathcal{S}_X \right| &\geq (1-2\beta)\frac{n^2}{m^2} \sum_{X \in \mathcal{X}_H, f'(X) > m^{1-k}} f'(X) \\ &\geq (1-2\beta)\frac{n^2}{m^2} (f'(R, H) - m) \\ &= (1-2\beta)\frac{n^2}{m^2} \left(f^*(G^*, H)\frac{m^2}{n^2} - m \right) \\ &= (1-2\beta)f^*(G^*, H) - (1-2\beta)\frac{n^2}{m} \\ &\geq (1-2\beta) \left(f(G, H) - \frac{\epsilon}{10}n^2 \right) - (1-2\beta)\frac{n^2}{m} \\ &\geq f(G, H) - \epsilon n^2. \end{aligned}$$

As the \mathcal{S}_X are pairwise disjoint for distinct X , we have obtained a set P_H of induced subgraphs of G that are color-isomorphic to H , such that $|P_H| \geq f(G, H) - \epsilon n^2$. Notice further that for distinct $H \in \mathcal{C}_k$, the corresponding sets P_H are disjoint. ■

2.5 Tournaments

The proof of the tournament case of Theorem 1 is almost identical to the proof of the edge-colored case presented in this section. One just needs to prove the following analogue of Lemma 2.1 which is the following Lemma 2.14. Recall that an *orientation* is a directed simple graph without cycles of length 2. A K_k -subgraph of an orientation G is a k -vertex tournament subgraph of G . We similarly define a K_k -decomposition and a fractional K_k -decomposition of an orientation.

Lemma 2.14 *Let $k \geq 3$ be an integer and let $\epsilon > 0$. There exists $n_0 = n_0(k, \epsilon)$ such that the following holds. Suppose G is an orientation with $n > n_0$ vertices. Let f be a fractional K_k -decomposition of G . Then for every $H \in \mathcal{T}_k$ there is a set P_H of induced subgraphs of G that are isomorphic to H , such that $|P_H| \geq f(G, H) - \epsilon n^2$. Furthermore any two elements of $P = \bigcup_{H \in \mathcal{T}_k} P_H$ intersect in at most one vertex.*

The proof of Lemma 2.14 is completely analogous to the proof of Lemma 2.14 where instead of using the colored version of Szemerédi's regularity lemma (Lemma 2.9) we use the directed version of the lemma. We refer to [14] which contains this directed version of the main result of [22] and

therefore implies Lemma 2.14. We therefore obtain the following corollary, whose proof is analogous to that of corollary 2.2.

Corollary 2.15 *Let $k \geq 3$ be an integer, let $\mathbf{v} \in \mathbb{R}^{|\mathcal{T}_k|}$, and let $\gamma > 0$. There exists $N_{2.15} = N_{2.15}(k, \gamma, \mathbf{v})$ such that the following holds for all orientations G with $n > N_{2.15}$ vertices which have a fractional K_k -decomposition. For every $H \in \mathcal{T}_k$ there is a set P_H of induced subgraphs of G that are isomorphic to H , such that any two elements of $P = \cup_{H \in \mathcal{T}_k} P_H$ intersect in at most one vertex. Furthermore,*

$$(a) \quad |P| \geq \frac{|E(G)| - \gamma n^2}{\binom{k}{2}}.$$

$$(b) \quad \sum_{H \in \mathcal{T}_k} \mathbf{v}_H |P_H| \geq \frac{|E(G)|}{\binom{k}{2}} \nu_{\mathbf{v}}^*(G) - \gamma n^2.$$

Finally, we need the analogue of Lemma 2.6 for the tournament setting. The lemma is proved in exactly the same way using Lemmas 2.3 and 2.4 (which stay intact; recall that they do not depend on the setting, whether it is tournaments or edge colored graphs) and using the straightforward Lemma 2.5 (whose statement stays intact, but in its proof G is a tournament instead of a C -colored complete graph). We therefore obtain the following lemma which immediately implies the second part of Theorem 1.

Lemma 2.16 *Let $k \geq 3$ be an integer, let $\mathbf{v} \in \mathbb{R}^{|\mathcal{T}_k|}$ be indexed by \mathcal{T}_k , and let $\epsilon > 0$. Then there exist $N_{2.16} = N_{2.16}(\epsilon, \mathbf{v})$ such that the following holds. Let G be a tournament which is k -divisible and with $n > N_{2.16}$ vertices. Then, G is K_k -decomposable and, furthermore $\nu_{\mathbf{v}}(G) \geq (\nu_{\mathbf{v}}^* - \epsilon)(1 - \epsilon) - \epsilon$.*

3 Applications

3.1 Triangles in tournaments

Our first application of Theorem 1 concerns the simplest case $k = 3$ for tournaments. Note that here we have $\mathcal{T}_3 = \{T_3, C_3\}$ where T_3 denotes the transitive triangle and C_3 denoted the directed (cyclic) triangle. Recall from the introduction that the various possibilities for $\nu_{\mathbf{v}}$ for $\mathbf{v} \in \mathbb{R}^{|\mathcal{T}_3|}$ reduce to the cases where the smallest coordinate of \mathbf{v} is 0 and the largest coordinate is 1 (if \mathbf{v} is the constant vector, then trivially $\nu_{\mathbf{v}}$ equals that constant). We furthermore see that the case of $\mathbf{v}(T_3) = 0$ is trivial since the sequence of transitive n -vertex tournaments shows that $\nu_{\mathbf{v}} = 0$ in this case. Hence the *only* vector for which $\nu_{\mathbf{v}}$ is nontrivial to evaluate is the one which assigns $\mathbf{v}(T_3) = 1$ and $\mathbf{v}(C_3) = 0$.

Conjecture 1 *Let $\mathbf{v} \in \mathbb{R}^{|\mathcal{T}_3|}$ where $\mathbf{v}(T_3) = 1$ and $\mathbf{v}(C_3) = 0$. Then $\nu_{\mathbf{v}} = 1$.*

In [20] it was conjectured that every tournament can be packed with $\lceil n(n-1)/6 - n/3 \rceil = (1 - o(1))\binom{n}{2} / \binom{3}{2}$ edge-disjoint copies of T_3 (and, if true, this conjectured value is shown there to be optimal). However, notice that even if the conjecture in [20] is true, this by no means implies that

$\nu_{\mathbf{v}} = 1$, since we have no guarantee that a very large T_3 -packing is part of a triangle decomposition (notice also that a triangle decomposition exists whenever $n \equiv 1, 3 \pmod{6}$, by Kirkman's Theorem).

Here we prove the following theorem which implies Theorem 2.

Theorem 4 *Let $\mathbf{v} \in \mathbb{R}^{|T_3|}$ where $\mathbf{v}(T_3) = 1$ and $\mathbf{v}(C_3) = 0$. Then, $\nu_{\mathbf{v}} \geq \frac{85}{98}$. In particular, for all $n \equiv 1, 3 \pmod{6}$, every tournament on n vertices has a triangle decomposition where the number of C_3 in the decomposition is at most $\frac{13}{98 \cdot 6} n^2 (1 + o_n(1))$.*

Proof. By Theorem 1, it suffices to prove that $\nu_{\mathbf{v}}^* \geq \frac{85}{98}$. A computer assisted proof (outlined below) yields that $\nu_{\mathbf{v}}^*(14) = \frac{78}{91}$. As proved in Corollary 2.7 in [10]⁴ following an iterative improvement argument appearing first in Lemma 2.2 of [12], $\nu_{\mathbf{v}}^*(n) \geq (\nu_{\mathbf{v}}^*(r)(r-1) + 1)/r - o_n(1)$. So, plugging in the case $r = 14$, $\nu_{\mathbf{v}}^*(14) = \frac{78}{91}$ and taking the limit yields $\nu_{\mathbf{v}}^* \geq \frac{85}{98}$.

So, it remains to show that $\nu_{\mathbf{v}}^*(14) \geq \frac{78}{91}$. Let us recall that this means that for *every* tournament G on 14 vertices, there is a fractional triangle decomposition f such that $\nu_{\mathbf{v}}^*(f) \geq \frac{78}{91}$, or, equivalently, that $D_{\mathbf{v}}(f) \geq \frac{78}{91} \cdot \binom{14}{2} / \binom{3}{2} = 26$. In our case, since $\mathbf{v}(T_3) = 1$ and $\mathbf{v}(C_3) = 0$, this means that the sum of the values of f on all T_3 copies of G is at least 26. As a side note, we observe that any 14 vertex tournament G_0 that is obtained by taking three disjoint sets of vertices A, B, C with $|A| = |B| = 5$ and $|C| = 4$ and orienting all edges from A to B , from B to C and from C to A (the orientations of edges with both endpoints in the same part is arbitrary), has the property that each of its T_3 copies contains an edge with both endpoints in the same part. So, for any fractional triangle decomposition f , the sum of the values of f on all T_3 copies of such a G_0 is at most the number of edges with both endpoints in the same part which is $\binom{5}{2} + \binom{5}{2} + \binom{4}{2} = 26$. Thus, we always have $\nu_{\mathbf{v}}^*(14) \leq \frac{78}{91}$.

The naive computational approach would therefore be as follows. Generate all 14-vertex tournaments G (say, up to isomorphism). For each such G , write down the linear programming problem which finds a fractional triangle decomposition which maximizes the sum of the values it assigns to the T_3 elements of G , and verify that this maximum, denoted by $D^*(G)$ is always at least 26. This naive approach is infeasible since the number of (pairwise non-isomorphic) tournaments on 14 vertices is more than any computer can handle (already the number of 14-vertex strongly connected tournaments on 14 vertices is 28304491788158056 by the OEIS), moreover running a (rather large) linear programming instance on each. Instead we take the following significantly better approach.

We call a tournament G on $r+1$ vertices an *extension* of a tournament G' on r vertices, if G' is a subgraph of G . Notice that a tournament on r vertices has at most 2^r extensions as can be seen by adding a new vertex and considering all possible orientations of its r incident edges. The following simple lemma is immediate from the proof of Lemma 2.5 (the construction of f there).

Lemma 3.1 *Let G be a tournament with $r+1$ vertices. If $D^*(G) < t$, then it is an extension of some G' with $D^*(G') < t \cdot \frac{r-1}{r+1}$. ■*

⁴That corollary is used in [10] for fractional triangle packings but it is identical for fractional triangle decompositions.

r	Size of M_r	Threshold value	Below threshold	Lowest value
10	9733056	12.86	16	12
11	16384	15.72	256	15
12	524288	18.86	2048	18
13	8388608	22.3	98304	22
14	805306368	26	0	26

Table 1: The procedure for verifying that $D^*(G) \geq 26$ for all 1-vertex tournaments.

For $r \geq 3$ and a real t , let $\mathcal{T}_r(t)$ denote the set of all r -vertex tournaments G with $D^*(G) < t$. So, our goal is to prove that $\mathcal{T}_{14}(26) = \emptyset$. By lemma 3.1, it suffices to check all extensions of $\mathcal{T}_{13}(26 \cdot \frac{12}{14}) \subseteq \mathcal{T}_{13}(22.3)$. In turn, it suffices to check all extensions of $\mathcal{T}_{12}(18.86)$. In turn, it suffices to check all extensions of $\mathcal{T}_{11}(15.72)$. In turn, it suffices to check all extensions of $\mathcal{T}_{10}(12.86)$. So, we start by generating all non-isomorphic tournaments on 10 vertices. There are known lists of such tournaments, see <https://users.cecs.anu.edu.au/~bdm/data/digraphs.html>. There are only 9733056 such tournaments. We denote this set by M_{10} . For each $G \in M_{10}$, we run the corresponding linear program to compute $D^*(G)$. If $D^*(G) \geq 12.86$ then, as shown earlier, we are not worried, as we are not missing anything by not checking extensions of such G . However, if $D^*(G) < 12.86$, we say that G is below the threshold, so we generate all 2^{10} extensions of G (we don't mind generating isomorphic tournaments, as the time required to check isomorphisms would be larger). Doing it for all G on 10 vertices which are below the threshold, yields a multiset of tournaments on 11 vertices, call it M_{11} . Notice that by the above, we know that if some G on 14 vertices has $D^*(G) < 26$, then it contains an element of M_{11} as a subgraph. Now, for each tournament $G \in M_{11}$, we run the corresponding linear program. If $D^*(G) < 15.72$, we generate all 2^{11} extensions of G . This yields a multiset of tournaments on 12 vertices, call it M_{12} . For each tournament $G \in M_{12}$, if $D^*(G) < 18.86$, we generate all 2^{12} extensions of G . This yields a multiset of tournaments on 13 vertices, M_{13} . For each tournament $G \in M_{13}$, if $D^*(G) < 22.3$, we generate all 2^{13} extensions of G . This yields a multiset of tournaments on 14 vertices, M_{14} . Finally, we check all tournaments in M_{14} to verify that $D^*(G) \geq 26$ for each of them. This procedure is summarized in Table 1. The table also lists for each $r = 10, \dots, 14$ the size of the (multi)set M_r , the number of elements of M_r that are below the threshold, which means that $|M_{r+1}|$ is precisely 2^r times larger than this amount. We also list the lowest value of $D^*(G)$ encountered during the search.

The code of the program that performs the procedure above can be found in <https://github.com/raphaelyuster/Vector-valued-decompositions>. The program runs in fewer than five days on standard personal computer equipment. The program uses the well-established linear programming package `lp-solve` which has a very efficient and simple to use api, see the `lp-solve` package homepage can be found at <https://sourceforge.net/projects/lpsolve>. Let us note that the linear programming instances are easy to generate. Suppose G is a tournament on r vertices. Generate a variable for each triple of vertices of G , so there are $\binom{r}{3}$ variables. To compute $D^*(G)$ one

should maximize the sum of the variables that correspond to triples that induce a T_3 . The constraints are: For each pair of vertices i, j of G , the sum of the variables that correspond to triples that contain both i, j should be precisely 1. Hence there are $\binom{r}{2}$ such constraints. Furthermore, we require that all variables are nonnegative. So there are $\binom{r}{3}$ such constraints. This completes the proof of Theorem 4.

3.2 Essentially avoidable graphs

In order to prove Theorem 3 we need to extend the notion of $\nu_{\mathbf{v}}^*(f)$ to fractional packings. Formally, for $\mathbf{v} \in \mathbb{R}^{|\mathcal{C}_k|}$ indexed by \mathcal{C}_k and a fractional K_k -packing f of a C -colored graph G , let $D_{\mathbf{v}}(f) = \sum_{H \in \mathcal{C}_k} \mathbf{v}_H f(G, H)$. After normalizing we define $\nu_{\mathbf{v}}^*(f) = D_{\mathbf{v}}(f) \binom{k}{2} / |E(G)|$. Since the result and proof in [22] applies to fractional packings, so do Lemma 2.1 and Corollary 2.2. Restated for fractional packings, Corollary 2.2 becomes:

Corollary 3.2 *Let C be a finite set of colors, let $k \geq 3$ be an integer, let $\mathbf{v} \in \mathbb{R}^{|\mathcal{C}_k|}$, and let $\gamma > 0$. There exists $N_{3.2} = N_{3.2}(k, C, \gamma, \mathbf{v})$ such that the following holds for all C -colored graphs G with $n > N_{3.2}$ vertices. Suppose f is a fractional K_k -packing of G . Then for every $H \in \mathcal{C}_k$ there is a set P_H of induced subgraphs of G that are color-isomorphic to H , such that any two elements of $P = \cup_{H \in \mathcal{C}_k} P_H$ intersect in at most one vertex. Furthermore,*

$$(a) \quad |P| \geq \frac{(\sum_{H \in \mathcal{C}_k} f(G, H)) - \gamma n^2}{\binom{k}{2}} .$$

$$(b) \quad \sum_{H \in \mathcal{C}_k} \mathbf{v}_H |P_H| \geq \frac{|E(G)|}{\binom{k}{2}} \nu_{\mathbf{v}}^*(f) - \gamma n^2 .$$

We say that a fractional K_k -packing f of a C -colored graph G is δ -close to a fractional decomposition if $\sum_{H \in \mathcal{C}_k} f(G, H) \geq |E(G)| - \delta n^2$. We say that a binary vector $\mathbf{v} \in \mathbb{R}^{|\mathcal{C}_k|}$ is nice if for every $\delta > 0$, if n is sufficiently large, then every C -colored complete graph G on n vertices has a fractional packing f that is δ -close to fractional decomposition and with $\nu_{\mathbf{v}}^*(f) \geq 1 - \delta$. With these definitions, together with Corollary 3.2, Lemma 2.3 and Lemma 2.4, the following lemma is proved in the same way Lemma 2.6 is proved.

Lemma 3.3 *Let C be a finite set of colors, let $k \geq 3$ be an integer, let $\mathbf{v} \in \mathbb{R}^{|\mathcal{C}_k|}$ be a nice vector indexed by \mathcal{C}_k , and let $\epsilon > 0$. Then there exist $N_{3.3} = N_{3.3}(\epsilon, \mathbf{v})$ such that the following holds. Let G be a C -colored complete graph which is k -divisible and with $n > N_{3.3}$ vertices. Then, G is K_k -decomposable and $\nu_{\mathbf{v}}(G) \geq 1 - \epsilon$. ■*

Let $C = \{\text{red}, \text{blue}\}$. A binary vector $\mathbf{v} \in \mathbb{R}^{|\mathcal{C}_k|}$ therefore corresponds to a characteristic vector of a subset $\mathcal{F}^* \subset \mathcal{F}_k$, where $\mathbf{v}_H = 1$ if and only if the blue edges of H correspond to a graph form \mathcal{F}^* .

Proof of Theorem 3. Let $k \geq 5$ be odd and $\mathcal{F}^* \subset \mathcal{F}_k$ be the family of all graphs H on k vertices such that both H and its complement are Eulerian. Let \mathbf{v} be the corresponding characteristic

vector of $\mathcal{F}_k \setminus \mathcal{F}^*$. Theorem 3 of [23] implies that \mathbf{v} is nice. By Lemma 3.3, if G is a k -divisible graph on $n > N_{3,3}$ vertices, then $\nu_v(G) \geq 1 - \epsilon$. Thus, \mathcal{F}^* is essentially avoidable.

For a graph H on k vertices, let \mathbf{v} be the characteristic vector of $\mathcal{F}_k \setminus \{H\}$. Let $\mathcal{U}_k \subseteq \mathcal{F}_k$ be the set of graphs H whose corresponding characteristic vector of $\mathcal{F}_k \setminus \{H\}$ is not nice. By Lemma 3.3, this implies that each $H \notin \mathcal{U}_k$ is essentially avoidable. Theorem 2 of [23] implies that $|\mathcal{U}_k| = o(|\mathcal{F}_k|)$. Hence the second part of the theorem follows. \blacksquare

4 Small k

We start this section by considering $\nu_{\mathbf{v}}$ for $\mathbf{v} \in \mathbb{R}^{|\mathcal{F}_3|}$ which is the first nontrivial case. By Theorem 1 it suffices to determine $\nu_{\mathbf{v}}^*$ and as noted in the introduction our problem is reduced to vectors whose smallest coordinate is 0 and whose largest coordinate is 1. We call such vectors *normalized*.

Observe that $\mathcal{F}_3 = \{K_3, P_3, Q_3, I_3\}$ where P_3 denotes the path on three vertices, $I_3 = K_3^c$ is the independent set on 3 vertices and $Q_3 = P_3^c$. We will use the convention of writing $\mathbf{v} = (\mathbf{v}(K_3), \mathbf{v}(P_3), \mathbf{v}(Q_3), \mathbf{v}(I_3))$. The following proposition determines $\nu_{\mathbf{v}}$ for a significant amount of normalized vectors, which include in particular all binary vectors.

Proposition 4.1 *Let $\mathbf{v} \in \mathbb{R}^{|\mathcal{F}_3|}$ be a normalized vector.*

1. $\frac{1}{4} \min\{\mathbf{v}(K_3), \mathbf{v}(I_3)\} \leq \nu_{\mathbf{v}} \leq \min\{\mathbf{v}(K_3), \mathbf{v}(I_3)\}$.
2. *If $\mathbf{v} = (1, 0, \beta, \alpha)$ or $\mathbf{v} = (\alpha, \beta, 0, 1)$, then $\nu_{\mathbf{v}} = \frac{\alpha}{4}$.*

Proof. For the first part of the proposition, consider first $G = K_n$. Here each 3-vertex subgraph is a K_3 so we obtain $\nu_{\mathbf{v}}^*(G) = \mathbf{v}(K_3)$. Similarly, for $G = I_n$ we have $\nu_{\mathbf{v}}^*(G) = \mathbf{v}(I_3)$. Hence, $\nu_{\mathbf{v}}^*(n) \leq \min\{\mathbf{v}(K_3), \mathbf{v}(I_3)\}$ so $\nu_{\mathbf{v}} = \nu_{\mathbf{v}}^* \leq \min\{\mathbf{v}(K_3), \mathbf{v}(I_3)\}$.

We recall a theorem of Goodman [7] who proved that in any n -vertex graph, $\frac{1}{4} \binom{n}{3} (1 - o_n(1))$ of the sets of 3 vertices induce either a K_3 or an I_3 . Hence, the fractional decomposition f which assigns a value of $1/(n-2)$ to each 3-set of vertices has $\nu_{\mathbf{v}}^*(f) \geq (1 - o_n(1)) \frac{1}{4} \{\min\{\mathbf{v}(K_3), \mathbf{v}(I_3)\}$. Hence, $\nu_{\mathbf{v}} = \nu_{\mathbf{v}}^* \geq \frac{1}{4} \{\min\{\mathbf{v}(K_3), \mathbf{v}(I_3)\}$.

For the second part of the proposition, note first that for $\mathbf{v} = (1, 0, \beta, \alpha)$ or $\mathbf{v} = (\alpha, \beta, 0, 1)$, the aforementioned lower bound implies that $\nu_{\mathbf{v}} = \nu_{\mathbf{v}}^* \geq \frac{\alpha}{4}$.

For the upper bound, we consider first the case $\mathbf{v} = (1, 0, \beta, \alpha)$. Let G be the complete balanced bipartite graph on n vertices, where the sides are A, B with $|A| = \lfloor n/2 \rfloor$ and $|B| = \lceil n/2 \rceil$. We assume that $n \equiv 1, 3 \pmod{6}$ so that there is a 3-decomposition of G . Notice that this implies that n is odd and that $|A||B|$ is even. Consider some 3-decomposition L of G . As the $|A||B| = (n^2 - 1)/4$ edges of G must be packed, there exist precisely $|A||B|/2 = (n^2 - 1)/8$ elements of L that are isomorphic to P_3 . These elements also contain $(n^2 - 1)/8$ pairs of vertices with both endpoints in the same part, so L has precisely $n(n-1)/6 - (n^2 - 1)/8 = (n^2 - 4n + 3)/24$ elements isomorphic to I_3 . This proves that

$$\nu_{\mathbf{v}}(n) \leq \alpha \frac{n^2 - 4n + 3}{24} \cdot \frac{3}{\binom{n}{2}} = \frac{\alpha}{4} \cdot \frac{n-3}{n}$$

proving that $\nu_{\mathbf{v}} \leq \frac{\alpha}{4}$. The case $\mathbf{v} = (\alpha, \beta, 0, 1)$ is proved analogously by taking complements. ■

Observe that Proposition 4.1 determines $\nu_{\mathbf{v}}$ for all binary vectors. It is always 0 unless $\mathbf{v}(K_3) = \mathbf{v}(I_3) = 1$ in which case it is $\frac{1}{4}$ except for the trivial case $\mathbf{v} = (1, 1, 1, 1)$ where we have $\nu_{\mathbf{v}} = 1$. Still, Proposition 4.1 does not cover all possible normalized vectors, so we raise the following problem.

Problem 1 Determine $\nu_{\mathbf{v}}$ for all (normalized vectors) $\mathbf{v} \in \mathbb{R}^{|\mathcal{F}_3|}$.

Moving to the next case $k = 4$, we do not know the value of $\nu_{\mathbf{v}}$ even for all binary vectors. Notice that $|\mathcal{F}_4| = 11$ as there are 11 distinct 4-vertex graphs. While the all-1 vector trivially has $\nu_{\mathbf{v}} = 1$, we do not know of a single normalized vector for which $\nu_{\mathbf{v}} = 1$. So, a realistic open problem is the following.

Problem 2 Determine the normalized vectors $\mathbf{v} \in \mathbb{R}^{|\mathcal{F}_4|}$ for which $\nu_{\mathbf{v}} = 1$.

Small examples suggest that it is plausible that the binary vector which assigns 1 to all graphs in \mathcal{F}_4 except C_4 and assigns 0 to C_4 has $\nu_{\mathbf{v}} = 1$. Finally, note that for $k = 5$ we know of a normalized binary vector which has $\nu_{\mathbf{v}} = 1$. Indeed by Theorem 3, the vector which assigns 1 to all graphs in \mathcal{F}_5 except C_5 and assigns 0 to C_5 has $\nu_{\mathbf{v}} = 1$.

5 The random graph

In this section we asymptotically determine $\nu_{\mathbf{v}}(G)$ for almost all k -decomposable graphs G and for all $\mathbf{v} \in \mathbb{R}^{|\mathcal{F}_k|}$. The main result in this section is stated for the Erdős-Rényi random graph probability space $\mathcal{G}(n, p)$ where $0 < p < 1$ is a constant. Recall that a property which holds for almost all $G \sim \mathcal{G}(n, \frac{1}{2})$ is referred to as a property that holds for almost all graphs.

Recall that an n -vertex graph $G \sim \mathcal{G}(n, p)$ is obtained by independently deciding for each pair of vertices whether it is an edge with probability p . Now, suppose n is such that graphs with n vertices are k -decomposable (recall that by Wilson's Theorem this holds for all n sufficiently large such that K_n is K_k -divisible). Then, for $G \sim \mathcal{G}(n, p)$ we have that $\nu_{\mathbf{v}}(G)$ is a random variable, and hence our ultimate goal would be to show that $\nu_{\mathbf{v}}(G)$ converges in distribution to a constant, and determine this constant. Indeed this is the main result in this section.

To define the constant to which $\nu_{\mathbf{v}}(G)$ converges in distribution, we set up a small (constant size) linear program. Let $\mathbf{v} \in \mathbb{R}^{|\mathcal{F}_k|}$ and consider the linear program $LP(\mathbf{v}, p)$ defined as follows.

$$\begin{aligned} \mathbf{max} \quad & \sum_{H \in \mathcal{F}_k} \mathbf{v}_H x_H \\ \mathbf{s.t.} \quad & \sum_{H \in \mathcal{F}_k} \left(e(H) - p \binom{k}{2} \right) x_H = 0, \\ & \sum_{H \in \mathcal{F}_k} x_H = 1, \\ & x_H \geq 0 \quad \forall H \in \mathcal{F}_k. \end{aligned}$$

Clearly $LP(\mathbf{v}, p)$ is feasible since setting $x_{K_k} = p$ and $x_{I_k} = (1 - p)$ and setting all other variables to 0, all constraints are satisfied. Therefore, let $s(\mathbf{v}, p)$ denote the optimal solution of $LP(\mathbf{v}, p)$. Our main theorem follows. Notice that when we write $n \rightarrow \infty$ we only consider n such that graphs with n vertices are k -decomposable.

Theorem 5 *Let $\mathbf{v} \in \mathbb{R}^{|\mathcal{F}_k|}$ and let $0 < p < 1$. For every $\epsilon > 0$, $G \sim \mathcal{G}(n, p)$ satisfies*

$$\lim_{n \rightarrow \infty} \Pr [|\nu_{\mathbf{v}}(G) - s(\mathbf{v}, p)| < \epsilon] = 1.$$

Proof. We first prove that $\Pr [\nu_{\mathbf{v}}(G) \geq s(\mathbf{v}, p) + \epsilon] = o_n(1)$. To this end, we don't even need to assume that G is a random graph. All that suffices is to assume that $e(G) = p\binom{n}{2} \pm o(n^2)$, which trivially holds with probability $1 - o_n(1)$ for $G \sim \mathcal{G}(n, p)$. So, assume that G is an n -vertex, k -decomposable graph with $e(G) = p\binom{n}{2} \pm o(n^2)$. We prove that $\nu_{\mathbf{v}}(G) \leq s(\mathbf{v}, p) + \epsilon$. Take an optimal k -decomposition L of G with respect to \mathbf{v} . For $H \in \mathcal{F}_k$, let L_H be the subset of L whose elements are isomorphic to H and let $y_H = |L_H|/|L|$. Observe that $y_H \geq 0$ and that $\sum_{H \in \mathcal{F}_k} y_H = 1$. Next, observe that $e(H)|L_H|$ is the total number of edges of G in all the elements of L_H , and since L is a decomposition, we have that $\sum_{H \in \mathcal{F}_k} e(H)|L_H| = e(G)$ and so $\sum_{H \in \mathcal{F}_k} e(H)y_H = e(G)/|L|$. But since $|L| = \binom{n}{2}/\binom{k}{2}$ and since $e(G) = p\binom{n}{2} \pm o(n^2)$ we have that $\sum_{H \in \mathcal{F}_k} (e(H) - p\binom{k}{2})y_H = o_n(1)$. Hence, there exist z_H such that $|z_H - y_H| = o_n(1)$ for all $H \in \mathcal{F}_k$ such that z_H form a feasible solution of $LP(\mathbf{v}, p)$ and such that for all n sufficiently large, $\sum_{H \in \mathcal{F}_k} \mathbf{v}_H(y_H - z_H) \leq \epsilon$. As the z_H form a feasible solution we get that

$$\nu_{\mathbf{v}}(G) = \nu_{\mathbf{v}}(L) = \sum_{H \in \mathcal{F}_k} \mathbf{v}_H y_H \leq \epsilon + \sum_{H \in \mathcal{F}_k} \mathbf{v}_H z_H \leq s(\mathbf{v}, p) + \epsilon.$$

We next prove that $\Pr [\nu_{\mathbf{v}}(G) \geq s(\mathbf{v}, p) - \epsilon] = 1 - o_n(1)$. We will assume for simplicity that p is rational. This can be assumed since for given n, ϵ, \mathbf{v} , the function $\Pr [\nu_{\mathbf{v}}(G) \geq s(\mathbf{v}, p) - \epsilon]$ where $G \sim \mathcal{G}(n, p)$ is continuous in p .

Now, as p is rational, so is $s(\mathbf{v}, p)$ and there is an optimal solution $\mathbf{x} = \{x_H : H \in \mathcal{F}_k\}$ where all the x_H are rational. By taking a common denominator d , we denote $x_H = a_H/d$ where the a_H are nonnegative integers not exceeding d . It will be convenient to view $G \sim \mathcal{G}(n, p)$ as an edge colored K_n where the blue edges are the edges of G and the red edges are the non-edges of G and similarly view the elements of \mathcal{F}_K as blue-red edge colored K_k .

We construct a gadget blue-red edge-colored graph D as follows. D consists of d edge-disjoint copies of K_k (any such graph D suffices). For each $H \in \mathcal{F}_k$ precisely a_H of the K_k comprising D are color-isomorphic to H . Notice that D has precisely $d\binom{k}{2}$ edges, where $\sum_{H \in \mathcal{F}_k} a_H e(H)$ of them are colored blue and the others are colored red. But observe that since the x_H form a feasible solution to $LP(\mathbf{v}, p)$, this also means that the number of blue edges of D is $pd\binom{k}{2}$ and the number of red edges is $(1 - p)d\binom{k}{2}$.

Let $r > 1$ be the smallest integer such that K_r has a D -decomposition. By Wilson's Theorem, r exists. Let R be a blue-red edge coloring of K_r obtained by taking a D -decomposition of K_r , and

coloring each element of this decomposition such that it is color isomorphic to D . Observe that the number of blue edges of R is $p\binom{r}{2}$ and the number of red edges is $(1-p)\binom{r}{2}$.

Now we consider $G \sim \mathcal{G}(n, p)$ (recall that G is viewed as a blue-red edge-colored K_n). We construct an $\binom{r}{2}$ uniform hypergraph M as follows. The vertices of M are the $\binom{r}{2}$ edges of G . The edges of M are all the K_r -subgraphs of G that are color-isomorphic to R . We observe some properties of M which stem from the fact that $G \sim \mathcal{G}(n, p)$. What is the degree of a blue vertex of M , or, stated equivalently, what is the number of copies of R in G that contain a given blue edge? For an r -set of vertices of G , let q denote the probability that it induces R . It doesn't really matter what q is, but nevertheless it is easy to compute it: $q = p^{\binom{r}{2}}(1-p)^{\binom{r}{2}-\binom{r}{2}}r!/aut(R)$ where $aut(R)$ is cardinality of the color-preserving automorphism group of R . For a given pair of vertices u, v and for an additional set W of $r-2$ vertices, what is the probability that $W \cup \{u, v\}$ induces R and that (u, v) is blue? Since only a p fraction of edges of R are blue, and given that $W \cup \{u, v\}$ induces R , (u, v) is equally likely to be any edge of R , the probability that $W \cup \{u, v\}$ induces R and that (u, v) is blue is precisely pq . Now, given that (u, v) is blue, the probability of an additional subset W of $r-2$ vertices to induce together with u, v a copy of R is, by conditional expectation, precisely $pq/p = q$. Hence, the expected degree of a blue vertex of M is precisely $q\binom{n-2}{r-2}$. Similarly given that (u, v) is red, the probability of an additional subset W of $r-2$ vertices to induce together with u, v a copy of R is, by conditional expectation, precisely $(1-p)q/(1-p) = q$ so the the expected degree of a red vertex of M is also precisely $q\binom{n-2}{r-2}$. Since the degree of a vertex of M (i.e. edge of G) is a random variable which is the sum of $\binom{n-2}{r-2}$ indicator random variables and each variable only depends on $O(n^{r-3})$ other variables, we have by Janson's inequality that for all n sufficiently large, the probability that all vertices of M have their degrees $q\binom{n-2}{r-2} + o(n^{r-2})$ is $1 - o_n(1)$. Another (trivial) property of M is that the co-degree of any two vertices of M , or equivalently, the number of copies of R in G that contain two distinct given edges is $O(n^{r-3})$.

Given these properties of M we can now apply Lemma 2.11 (the Frankl-Rödl hypergraph matching theorem) which states that with probability $1 - o_n(1)$, M has a matching covering all but $o(V(M))$ of the vertices of M . In other words, with probability $1 - o(1)$, there is a packing of G with pairwise edge-disjoint copies of R , such that the number of unpacked edges is $o(n^2)$. But now recall that each copy of R decomposes into D and each copy of D contains, for each $H \in \mathcal{F}_k$, precisely a_H pairwise edge-disjoint K_k subgraph that are color-isomorphic to H . But since the x_H are an optimal solution to $LP(\mathbf{v}, p)$, we get that with probability $1 - o_n(1)$, there is a k -packing P of G such that $\nu_{\mathbf{v}}(P) \geq s(\mathbf{v}, p) - o_n(1)$. We can now only slightly modify P to obtain a k -decomposition L using Lemma 2.3 precisely in the same way shown in Lemma 2.6 where $\nu_{\mathbf{v}}(L) \geq \nu_{\mathbf{v}}(P) - o_n(1)$. Thus, $\nu_{\mathbf{v}}(L) \geq s(\mathbf{v}, p) - o_n(1)$ with probability $1 - o_n(1)$, implying that for every $\epsilon > 0$, $\Pr[\nu_{\mathbf{v}}(G) \geq s(\mathbf{v}, p) - \epsilon] = 1 - o_n(1)$.

Combining now the two parts of the proof we obtain that $\Pr[|\nu_{\mathbf{v}}(G) - s(\mathbf{v}, p)| < \epsilon] = 1 - o_n(1)$, implying the theorem. \blacksquare

Since $s(\mathbf{v}, p)$ can be solved in constant time for every $\mathbf{v} \in \mathbb{R}^{|\mathcal{F}_k|}$, we can view Theorem 5 as saying that the asymptotic value of $\nu_{\mathbf{v}}(G)$ is determined for almost all graphs (using $p = \frac{1}{2}$).

We end this section with an example of a nontrivial case already for $k = 3$. Using the notation of the previous section, consider the vector $\mathbf{v} \in \mathbb{R}^{|\mathcal{F}_3|}$ defined by $\mathbf{v}(K_3) = 1$, $\mathbf{v}(P_3) = \frac{1}{2}$, $\mathbf{v}(Q_3) = \frac{1}{2}$, $\mathbf{v}(I_3) = 0$ and assume $p = \frac{1}{2}$. Putting $x_3 = x_{K_3}$, $x_2 = x_{P_3}$, $x_1 = x_{Q_3}$, $x_0 = x_{I_3}$, the linear program $LP(\mathbf{v}, \frac{1}{2})$ becomes:

$$\begin{aligned} \mathbf{max} \quad & \frac{1}{2}x_1 + \frac{1}{2}x_2 + x_3 \\ \mathbf{s.t.} \quad & -\frac{3}{2}x_0 - \frac{1}{2}x_1 + \frac{1}{2}x_2 + \frac{3}{2}x_3 = 0, \\ & x_0 + x_1 + x_2 + x_3 = 1, \\ & x_i \geq 0 \quad \forall i \in \{0, 1, 2, 3\}. \end{aligned}$$

The optimal solution here is $s(\mathbf{v}, \frac{1}{2}) = \frac{5}{8}$ with $x_1 = \frac{3}{4}$, $x_3 = \frac{1}{4}$, $x_0 = x_2 = 0$. Mimicking the proof of Theorem 5, we construct a gadget blue-red edge colored graph D consisting of four edge disjoint triangles. One triangle is completely blue (this corresponds to one copy of K_3), the other three triangles each have two red edges and one blue edge (this corresponds to three copies of Q_3). We observe that D has 12 edges, 6 of which are blue and 6 are red. As in the proof of Theorem 5, a random graph $G \sim \mathcal{G}(n, \frac{1}{2})$ where the non-edges are colored red and the edges are colored blue almost surely almost decomposes to D . So, as in the theorem, this implies that we have a decomposition L of G into triangles where the number of blue triangles is roughly $n(n-1)/24$ and the number of triangles with two red edges and one blue edge is roughly $n(n-1)/8$. This yields that $\nu_{\mathbf{v}}(L) = \frac{5}{8}(1 - o_n(1))$.

References

- [1] B. Barber, D. Kühn, A. Lo, R. Montgomery, and D. Osthus. Fractional clique decompositions of dense graphs and hypergraphs. *Journal of Combinatorial Theory, Series B*, 127:148–186, 2017.
- [2] B. Barber, D. Kühn, A. Lo, and D. Osthus. Edge-decompositions of graphs with high minimum degree. *Advances in Mathematics*, 288:337–385, 2016.
- [3] D. Dor and M. Tarsi. Graph decomposition is NPC- a complete proof of Holyer’s conjecture. In *Proceedings of the twenty-fourth annual ACM symposium on Theory of computing*, pages 252–263, 1992.
- [4] P. Dukes. Rational decomposition of dense hypergraphs and some related eigenvalue estimates. *Linear Algebra and its Applications*, 436(9):3736–3746, 2012.
- [5] P. Frankl and V. Rödl. Near perfect coverings in graphs and hypergraphs. *European Journal of Combinatorics*, 6(4):317–326, 1985.

- [6] S. Glock, D. Kühn, A. Lo, and D. Osthus. The existence of designs via iterative absorption. arXiv:1611.06827, 2016.
- [7] W. Goodman. On sets of acquaintances and strangers at any party. *The American Mathematical Monthly*, 66(9):778–783, 1959.
- [8] P. E. Haxell and V. Rödl. Integer and fractional packings in dense graphs. *Combinatorica*, 21(1):13–38, 2001.
- [9] I. Holyer. The NP-Completeness of some edge-partition problems. *SIAM Journal on Computing*, 10(4):713–717, 1981.
- [10] M. Kabiya and R. Yuster. Packing transitive triples in a tournament. *Annals of Combinatorics*, 12(3):291–306, 2008.
- [11] P. Keevash. The existence of designs. arXiv:1401.3665, 2014.
- [12] P. Keevash and B. Sudakov. Packing triangles in a graph and its complement. *Journal of Graph Theory*, 47(3):203–216, 2004.
- [13] T. P. Kirkman. On a problem in combinations. *Cambridge and Dublin Mathematical Journal*, 2:191–204, 1847.
- [14] Z. Nutov and R. Yuster. Packing directed cycles efficiently. *Discrete applied mathematics*, (2):82–91, 2007.
- [15] E. Szemerédi. Regular partitions of graphs. In *Problèmes combinatoires et théorie des graphes (Colloq. Internat. CNRS, Univ. Orsay, Orsay, 1976)*, volume 260 of *Colloq. Internat. CNRS*, pages 399–401. CNRS, Paris, 1978.
- [16] R. M. Wilson. An existence theory for pairwise balanced designs I. Composition theorems and morphisms. *Journal of Combinatorial Theory, Series A*, 13(2):220–245, 1972.
- [17] R. M. Wilson. An existence theory for pairwise balanced designs II. The structure of PBD-closed sets and the existence conjectures. *Journal of Combinatorial Theory, Series A*, 13(2):246–273, 1972.
- [18] R. M. Wilson. Decomposition of complete graphs into subgraphs isomorphic to a given graph. *Congressus Numerantium*, 15:647–659, 1975.
- [19] R. M. Wilson. An existence theory for pairwise balanced designs, III: Proof of the existence conjectures. *Journal of Combinatorial Theory, Series A*, 18(1):71–79, 1975.
- [20] R. Yuster. The number of edge-disjoint transitive triples in a tournament. *Discrete Mathematics*, 287(1-3):187–191, 2004.

- [21] R. Yuster. Asymptotically optimal K_k -packings of dense graphs via fractional K_k decompositions. *Journal of Combinatorial Theory, Series B*, 95(1):1–11, 2005.
- [22] R. Yuster. Integer and fractional packing of families of graphs. *Random Structures & Algorithms*, 26(1-2):110–118, 2005.
- [23] R. Yuster. Packing without some pieces. *Journal of Combinatorics*, to appear, 2018.