The University of British Columbia



TIME: 75 MINUTES

LAST NAME:	FIRST NAME:	
STUDENT # ·	SIGNATURE	

This Examination paper consists of 9 pages (including this one). Make sure you have all 9.

INSTRUCTIONS:

No memory aids allowed. No calculators allowed. No communication devices allowed.

MARKING:

Q1	/8
$\mathbf{Q2}$	/10
$\mathbf{Q3}$	/14
Q4	/10
Q5	/8
TOTAL	/50

Q1 [8 marks]

Use Green's Theorem to evaluate

$$\int_C (\cos(x^2 - x) - y^3) dx + (x^3 - y^2) dy$$

where C is the curve

$$\vec{\mathbf{r}}(t) = 2\cos t \, \vec{\mathbf{i}} + 2\sin t \, \vec{\mathbf{j}}, \qquad -\frac{\pi}{2} \le t \le \frac{\pi}{2}$$

with direction increasing with t. Notice that C is <u>not closed</u>.

 $\mathbf{Q2}$ [10 marks]

Let $\vec{\mathbf{F}}(x,y) = P(x,y) \ \vec{\mathbf{i}} + Q(x,y) \ \vec{\mathbf{j}}$ be a vector field defined everywhere except (-2,0) and (2,0). Let C_1 be the circle $(x+2)^2 + y^2 = 1$, oriented <u>counterclockwise</u>, and let C_2 be the circle $(x-2)^2 + y^2 = 1$, oriented <u>clockwise</u>. It is given that

$$P_y = Q_x, \qquad \int_{C_1} \vec{F} \cdot d\vec{r} = 2, \qquad \int_{C_2} \vec{F} \cdot d\vec{r} = \pi .$$

Compute the following line integrals. No credit will be given for answers without an explanation. You can get partial credit by drawing all relevant curves with their correct orientation.

(a) Let C be the circle $(x-2)^2 + y^2 = 0.25$, oriented <u>counterclockwise</u>.

 $\int_C \vec{\mathbf{F}} \cdot d\vec{\mathbf{r}} =$

(b) Let C be the circle $x^2 + y^2 = 25$, oriented <u>clockwise</u>. $\int_C \vec{\mathbf{F}} \cdot d\vec{\mathbf{r}} =$

MATH 317 $\,$ MIDTERM 2 — August 6, 2015 — p. 4 of 9 $\,$

(c) Let C be the curve consisting of the four oriented line segments: (-5, 5) to (5, -5), (5, -5) to (5, 5), (5, 5) to (-5, -5), and (-5, -5) to (-5, 5). (Notice that C is not simple.)

 $\int_C \vec{\mathbf{F}} \cdot d\vec{\mathbf{r}} =$

Q3 [14 marks]

Let S be the surface parameterized by

$$\vec{\mathbf{r}}(u,v) = \cos u(\cos v + 2) \,\vec{\mathbf{i}} + \sin u(\cos v + 2) \,\vec{\mathbf{j}} + \sin v \,\vec{\mathbf{k}}$$

where $0 \le u \le 2\pi$, $0 \le v \le 2\pi$.

(a) Draw S. (Hint: Try substituting $u = 0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}$.)

(b) Below is a list of parameterized surfaces. Clearly mark those which also parameterizes S. A: $\vec{\mathbf{r}}(u,v) = \langle \sqrt{1-u^2}(2-\sin(2v)), 2u-u\sin(2v), \cos(2v) \rangle, -1 \le u \le 1, 0 \le v \le \pi$. B: $\vec{\mathbf{r}}(u,v) = \langle -\sin v(2-\cos u), \cos v(2-\cos u), \sin u \rangle, 0 \le u \le 2\pi, 0 \le v \le 2\pi$. C: $\vec{\mathbf{r}}(u,v) = \langle \sin u(2+\sqrt{1-v^2}), \cos u(2+\sqrt{1-v^2}), v \rangle, 0 \le u \le 2\pi, -1 \le v \le 1$. D: $\vec{\mathbf{r}}(u,v) = \langle \cos(2u)(\cos v - 2), \sin(2u)(\cos v - 2), \sin v \rangle, 0 \le u \le \pi, 0 \le v \le 2\pi$. E: $\vec{\mathbf{r}}(u,v) = \langle v \cos u, v \sin u, \sqrt{1-(v-2)^2} \rangle, 0 \le u \le 2\pi, 1 \le v \le 3$. MATH 317 MIDTERM 2 — August 6, 2015 — p. 6 of 9

(c) Find the area of S.

Q4 [10 marks]

Let f(u) be a function defined for any number u, and let $f' = \frac{\partial f}{\partial u}$ and $f'' = \frac{\partial^2}{\partial u^2} f$. Consider the vector field

$$\vec{\mathbf{F}}(x,y,z) = f(r)\bar{\mathbf{r}}$$

where $\vec{\mathbf{r}} = \langle x, y, z \rangle$ and $r = |\vec{\mathbf{r}}|$. In the following questions, express you final answer in terms of $\vec{\mathbf{r}}, r, f, f'$ and f''.

(a) Simplify grad f(r). (Hint: Use the chain rule.)

(b) Simplify div $\vec{\mathbf{F}}$.

MATH 317 MIDTERM 2 — August 6, 2015 — p. 8 of 9

(c) Simplify $\operatorname{curl} \vec{\mathbf{F}}$.

(d) Simplify $\nabla^2(f(r))$.

Q5 [8 marks]

Let S be the part of the cylinder $y^2 + z^2 = 4$ lying between the planes $x = 3 + \frac{1}{2}y + \frac{1}{2}z$ and $x = -\frac{1}{2}z$.

(a) Complete the following parametrization: The surface S can be parameterized as $\vec{\mathbf{r}}(u, v) = v \vec{\mathbf{i}} + 2 \cos u \vec{\mathbf{j}} + 2 \sin u \vec{\mathbf{k}}$ where $0 \le u \le 2\pi$ and $f(u) \le v \le g(u)$ with



(b) Compute the flux integral $\iint_S \vec{\mathbf{F}} \cdot d\vec{\mathbf{S}}$ where $\vec{\mathbf{F}}(x, y, z) = e^{xyz} \vec{\mathbf{i}} + \vec{\mathbf{k}}$ and S is oriented such that its normal vector points toward the inside the cylinder $y^2 + z^2 = 4$. (You may use the identities $\sin^2 \alpha = \frac{1-\cos(2\alpha)}{2}$ and $\cos^2 \alpha = \frac{1+\cos(2\alpha)}{2}$.)